arXiv:math/0102072v1 [math.DG] 8 Feb 2001

Index- and Spectral Theory for Manifolds with Generalized Fibred Cusps [*]

Boris Vaillant
Department of Mathematics, University of Bonn

February 1, 2008

Abstract

Generalizing work of W. Müller we investigate the spectral theory for the Dirac operator D on a noncompact manifold X with generalized fibred cusps

$$C(M) = M \times [A, \infty[_r, \qquad g = \, dr^2 + \phi^* g_Y + e^{-2cr} g_Z,$$

at infinity. Here $\phi : M^{h+v} \to Y^h$ is a compact fibre bundle with fibre Z and a distinguished horizontal space HM. The metric g_Z is a metric in the fibres and g_Y is a metric on the base of the fibration. We also assume that the kernel of the vertical Dirac operator at infinity forms a vector bundle over Y.

Using the "ϕ-calculus" developed by R. Mazzeo and R. Melrose we explicitly construct the meromorphic continuation of the resolvent $G(\lambda)$ of D for small spectral parameter as a special "conormal distribution". From this we deduce a description of the generalized eigensections and of the spectral measure of D.

Complementing this, we perform an explicit construction of the heat kernel $[\exp(-t\mathsf{D}^2)]$ for finite and small times t, corresponding to large spectral parameter λ. Using a generalization of Getzler's technique, due to R. Melrose, we can describe the singular terms in the heat kernel expansion for small times in the interior of the manifold as well as at spatial infinity. This then allows us to prove an index formula for D,

$$\mathrm{ind}_-(\mathsf{D}) = \frac{1}{(2\pi i)^{n/2}} \int_X \widehat{A}(R)\ \mathrm{Ch}(F^{E/S}) + \frac{1}{(2\pi i)^{(h+1)/2}} \int_Y \widehat{A}(R^Y) \widehat{\eta}(\mathsf{D}^V) + \frac{1}{2}\eta(\mathsf{D}_Y),$$

which calculates the extended L^2-index of D, in terms of the usual local expression, the family eta invariant for the family of vertical Dirac operators at infinity and the eta invariant for the horizontal "Dirac" operator at infinity.

---

[*] AMS Class.: 58G25, 58Gxx. This is the author's doctoral thesis submitted at the University of Bonn. Written with support from the DFG through funds from the SFB 256 and a fellowship from the "Graduiertenkolleg für Mathematik der Universität Bonn". Financial support was also obtained from the DAAD through a "HSP III"-scholarship.

Contents

Introduction **1**

1 The Geometry of ϕ-Manifolds **5**
- 1.1 ϕ-Manifolds . . . 5
- 1.2 Exact ϕ-Metrics . . . 8
- 1.3 ϕ-Differential Operators and the Dirac Operator D^d . . . 11
- 1.4 Geometry of the Boundary . . . 14

2 The ϕ-Calculus **20**
- 2.1 The b- and the ϕ-Blow up . . . 20
- 2.2 Mapping Properties . . . 23
- 2.3 Composition Formula . . . 26

3 Normal Operators and the Resolvent Construction **31**
- 3.1 Symbol Map . . . 31
- 3.2 Symbolic Construction . . . 32
- 3.3 Normal Operator at ff . . . 33
- 3.4 Inversion at ff . . . 35
- 3.5 Reduced Normal Operators at bf and ϕbf . . . 39
- 3.6 Inversion at ϕbf . . . 46
- 3.7 End of the Construction . . . 51
- 3.8 Consequences . . . 54

4 The Heat Kernel at Finite Time **64**
- 4.1 d-Heat Space and d-Heat Calculus . . . 64
- 4.2 Composition Formula . . . 68
- 4.3 Normal Operators and Construction of the Heat Kernel . . . 73

5 Rescaled d-Heat Calculus **79**
- 5.1 The Double Rescaling . . . 79
- 5.2 The Rescaled Calculus and Normal Operators . . . 86
- 5.3 The Rescaled Heat Kernel and the Index . . . 94
- 5.4 The Signature Operator S . . . 98

A Appendix **104**
- A.1 Clifford Algebra Conventions . . . 104
- A.2 Dirac Operator and Conformal Transformations . . . 106
- A.3 Geometry of Fibre Bundles . . . 107
- A.4 Mehler's Formula and the Simple Heat Space . . . 110
- A.5 Heat Kernels of Vertical Families . . . 113
- A.6 Conormal Functions, Pullbacks and Pushforwards . . . 116

References **120**

Introduction

In his book [Mü2] W. Müller, using methods from harmonic analysis, described the spectral decomposition and index theory of the Dirac operator over a locally symmetric space

$$\Gamma \setminus G/K$$

of rank one. This is a manifold equipped with a natural riemannian metric. Near infinity it has the structure of a “fibred cusp”

$$C(M) = M \times [A, \infty[_r,$$

where $\phi : M \to Y$ is a compact fibre bundle with fibre Z and a distinguished horizontal space HM. The metric on the cusp is given by

$$g = dr^2 + \phi^* g_Y + e^{-2cr} g_Z, \tag{1}$$

with g_Z a metric in the fibres and g_Y a metric on the base of the fibration.

In this work we investigate the spectral theory of the Dirac operator, denoted by D^d below, over a general complete manifold X, noncompact only in generalized cusps as above. We use the methods developed in [Me0], [Me3] and [MaMe3]. Following their philosophy we compactify X to a manifold with fibred boundary $\partial X = M$ by introducing $x = e^{-cr}$ as a defining function for the boundary. The metric on X will then be denoted by g_d. This is a degenerate (and singular) metric near the boundary, where it has the form

$$g_d = \left(\frac{dx}{x}\right)^2 + \phi^* g_Y + x^2 g_Z,$$

(here we have dropped the factors c^2 and c^{-2}). However, this metric is a *true* metric on the fibres of a vector bundle ${}^dTX$ whose space of sections ${}^d\mathcal{V}(X)$ can be identified with the space of sections of X into TX, possibly singular at ∂X, but of finite g_d-length.

In terms of local fibre coordinates $\mathbf{y}, \mathbf{z}$ in M and a product decomposition of X near ∂X, vector fields in ${}^d\mathcal{V}(X)$ are locally spanned over $C^\infty(X)$ by

$$x\frac{\partial}{\partial x}, \quad \frac{\partial}{\partial \mathbf{y}} = (\frac{\partial}{\partial y^1}, \ldots, \frac{\partial}{\partial y^h}), \quad \frac{1}{x}\frac{\partial}{\partial \mathbf{z}} = (\frac{1}{x}\frac{\partial}{\partial z^1}, \ldots, \frac{1}{x}\frac{\partial}{\partial z^v}).$$

It is easily seen from this representation that ${}^d\mathcal{V}(X)$ is *not* a Lie algebra, such that there is no good notion of d-differential operators etc.. This prompts the introduction of the *nondegenerate* conformal metric

$$g_\phi = \frac{1}{x^2} g_d = \left(\frac{dx}{x^2}\right)^2 + \frac{1}{x^2}\phi^* g_Y + g_Z. \tag{2}$$

The associated space of vector fields of finite g_ϕ-length,

$${}^\phi\mathcal{V}(X) := x \cdot {}^d\mathcal{V}(X) \subset \Gamma(X, TX),$$

consists of true vector fields on X and can be seen to be the space of sections of the vector bundle

$$^{\phi}TX = x\,^{d}TX.$$

It is easy to verify that it forms a Lie algebra and we can define the notions of ϕ-differential and ϕ-pseudodifferential operators. The analysis of these operators has been begun in [MaMe3], using the general framework set out in [Me0].

Now the spinor Dirac operator D^d, associated to the metric g_d, is a selfadjoint operator on $L^2(X, E, dvol_d)$. In Appendix A.2 it is shown that

$$\mathsf{D}^d = \mathbf{c}_d \circ \nabla^{E,d} = \frac{1}{x}\mathsf{D}^{\phi} + \frac{n-1}{2}\mathbf{c}_{\phi}(\frac{dx}{x^2}), \tag{3}$$

where $\mathsf{D}^{\phi} \in \operatorname{Diff}^1_{\phi}(X, E)$ is the spinor Dirac operator associated to g_{ϕ}. It follows from this – and is actually true for *any* Dirac operator D^d on a Clifford bundle associated with a metric g_d as above – that $x\mathsf{D}^d$ is an elliptic ϕ-differential operator. This will allow us to apply the analysis developed for ϕ-operators in the first part of the text to the spectral analysis of D^d.

Before describing the results obtained in this work, let us investigate the structure of the problem somewhat further. First, the restriction of the operator $x\mathsf{D}^d$ to the boundary in the usual sense of differential operators is an elliptic vertical family of first order differential operators $\mathsf{D}^{\phi,V}$. It is easy to see that D^d has compact resolvent, and therefore pure point spectrum, if this family is invertible. The other extreme case, where one allows arbitrary behavior of the null spaces of $\mathsf{D}^{\phi,V}$, turns out to be too difficult in general. In this work we make the additional *assumption*, fulfilled for instance in the case of the signature operator, that the null spaces of $\mathsf{D}^{\phi,V}$ form a vector bundle

$$\mathcal{K} \to Y, \qquad \Gamma(Y, \mathcal{K}) = \operatorname{null}(\mathsf{D}^{\phi,V}),$$

over the base Y of the boundary fibration. We will denote the projection onto this null space by $\Pi_{\circ}$ and write $B = Y \times \mathbf{R}_{+,x}$, which can be thought of as the cylindrical base of the model cusp (compactified at "infinity" $x = 0$). Then we can introduce the operator

$$I_b(\mathsf{D}^d) = "\Pi_{\circ}\mathsf{D}^d\Pi_{\circ}|_{\partial X, {}^bTX}" = \mathsf{D}_Y + \mathbf{c}_d(\frac{dx}{x})x\frac{\partial}{\partial x} : C^{\infty}(B, \mathcal{K}) \to C^{\infty}(B, \mathcal{K}). \tag{4}$$

This is described more precisely in Section 3.5. It turns out that – just as in the b-case, or the case of asymptotically cylindrical ends – the continuous spectrum of D^d is governed by this operator, with bands of continuous spectrum starting at the eigenvalues of D_Y and going out to infinity. Especially, the operator D^d is Fredholm, if and only if D_Y is invertible.

This result will be obtained as a corollary of the construction of the analytic continuation of the resolvent

$$G_{\mathsf{D}^d}(\lambda) = (\mathsf{D}^d - \lambda)^{-1}$$

of D^d described in Chapter 3. Using the concepts developed in [MaMe3], [Me0], [Me3] etc., we explicitly construct the resolvent as an integral kernel on the space X^2_{ϕ}. This is a compactification of the interior of X^2, which is adapted to the specific asymptotics of the problem at infinity. The main result, summarizing this procedure and stated as Theorem 3.25, gives a detailed account of the asymptotics of the resolvent. It is then possible to use the methods known from the b-calculus to derive the usual applications of this construction, such as a description of the spectral measure,

$$dE_{\mathsf{D}^d}(r) = \lim_{\varepsilon\to 0} \frac{1}{2\pi i}[G_{\mathsf{D}^d}(r+i\varepsilon) - G_{\mathsf{D}^d}(r-i\varepsilon)]\,dr,$$

of D^d in terms of its generalized eigenfunctions. This is done in Section 3.8.

The general framework for the constructions in Chapter 3 will be recalled in Chapter 2. Here the space X^2_ϕ is described as a blow up of X^2 and the notion of ϕ-pseudodifferential operators is introduced as a special class of distributions on X^2_ϕ. Building on the work in [MaMe3], we then analyze the mapping- and composition properties of such operators.

Chapter 1 contains an introduction to the geometry of manifolds with fibred boundaries endowed with a metric of type (2) as described above. Here we also introduce the concept of coding the asymptotic behavior of sections of a vector bundle into the structure of a new vector bundle. An instance of this has already been presented in the definition of the bundles ${}^dTX$, ${}^\phi TX$, and the general concept will be used repeatedly throughout the text. The more detailed results about the behavior near the boundary of the connections and curvatures associated to g_ϕ and g_d, obtained in the later parts of this Chapter, will however not be used seriously until Chapter 5.

The second important topic in this work is the proof of an index theorem for D^d. As a corollary to the description of the spectral measure, we can describe the large time behavior of the heat operator

$$\exp(-t\mathsf{D}^2) = \int_{\mathbf{R}} e^{-tr^2}\,dE_{\mathsf{D}^d}(r) \tag{5}$$

and prove that the extended L^2-index of D^d (is finite dimensional and) can be calculated as the large time limit of the regularized heat supertrace:

$$\mathrm{ind}_-(\mathsf{D}^d) = \lim_{T\to\infty} \mathrm{reg}_{z=0}\,\mathrm{Str}(x^z[\exp(-T(\mathsf{D}^d)^2)]).$$

This will allow us to derive a formula for the index using a variant of the McKean-Singer argument, i.e. by calculating the small time limit of the regularized heat supertrace and the difference between its large- and small-time limits. However, as can be seen from (5), the heat operator at small times involves the resolvent at large spectral parameter. Our construction in Chapter 3 does not give us sufficient control of the resolvent in that area.

Thus, in Chapter 4, we present a direct construction of the heat kernel for finite and small times. Again, this is done by introducing an appropriate compactification of the kernel space and a corresponding "heat calculus", within which the heat kernel can be constructed. Theorem 4.11 gives a certain amount of control of the asymptotics of the heat kernel at small times and near the boundary.

Chapter 5 is then devoted to a refinement of that result, providing more information about the Clifford degree of the coefficients in the asymptotic expansion. In the philosophy of Patodi, Getzler, etc. this will allow us to eliminate the divergent terms in the small time expansion(s) of the local heat supertrace. This then turns out to be sufficient to prove the index theorem. Denoting by $\eta(\mathsf{D}_Y)$ the classical eta-invariant associated to the operator D_Y and by $\widehat{\eta}(\mathsf{D}^{\phi,V}, E)$ the family eta invariant associated to the vertical family $\mathsf{D}^{\phi,V}$ (see Appendix A.5) we prove the following formula for the extended L^2-index of D in Theorem 5.29:

$$\operatorname{ind}_-(\mathsf{D}) = \frac{1}{(2\pi i)^{n/2}} \int_X \widehat{A}(R^d) \operatorname{Ch}(F^{E/S,d}) + \frac{1}{(2\pi i)^{(h+1)/2}} \int_Y \widehat{A}(R^Y)\widehat{\eta}(\mathsf{D}^{\phi,V}, E) + \frac{1}{2}\eta(\mathsf{D}_Y).$$

This formula generalizes the corresponding formula obtained by W. Müller in [Mü2]. It reduces to the usual APS (or b-) index formula in the case that there is no fibre, i.e. $Z = \{\text{pt}\}$.

In Section 5.4 we specialize our results to the case of the signature operator $\mathsf{S}^d = d_d + d_d^*$. It is a well known fact in the b-case that the extended L^2-index of this operator equals the L^2-signature. This is shown to be also true in our case. We compare our formulas with the adiabatic limit formula for the eta invariant in [BiCh1] and [Da1]. More serious applications are however left for future examination. Finally some auxiliary results and background material have been collected in the Appendix.

It is our declared purpose in this work, to give a thorough introduction to a set of methods for the spectral analysis of geometric operators associated to d-metrics. Especially, our proof of the index theorem is far from being "minimal". Remark 5.35 at the end of the text sketches several shortcuts for the proof.

Some remarks about notation: For simplicity we will assume throughout that M is *connected*, i.e. there is only one cusp. However all the results presented here generalize directly to the case of several cusps. Whenever there is a grading we use *graded* calculation rules. Thus commutators $[A, B]$ and tensor products $A \otimes B$ of graded objects are always graded. Notations like str or sdim are understood.

Acknowledgements

First, special thanks are due to both my advisors, Werner Müller and Richard Melrose. The work of Werner Müller is at the basis of this thesis. He suggested that I try to generalize his results in [Mü2] and supported my research for many years. I thank Richard Melrose for inviting me to MIT in 97/98 and taking the time to teach me his approach to the spectral analysis of operators in noncompact settings. Most of the methods used in this thesis are based on his ideas and I would not have got very far without his support.

Many more people have contributed to various parts of this work. Especially, I wish to thank my fellow students in Bonn and Cambridge, Sang Chin, Robert Lauter, Paul Loya, Sergiu Moroianu, Gorm Salomonsen, Oliver Sick and Jared Wunsch, who helped me learn the techniques used here and with whom I discussed many of the problems in detail. Some of their work (like [LaMo1], [LaMo2] and [Lo]) is closely related to mine.

1 The Geometry of ϕ-Manifolds

In this Chapter, we introduce the main geometrical notions associated to manifolds with a fibred boundary. The concepts and notation introduced in the first Section, as well as the notion of an exact ϕ-metric introduced at the beginning of Section 1.2, are fundamental to all the further parts of this work. On the other hand, most of the more detailed calculations in Sections 1.2 to 1.4 will only be used in later Chapters. They can be skipped at first reading and be referred back to as needed.

1.1 ϕ-Manifolds

Let X be a compact manifold of dimension n with connected and fibred boundary $\partial X = M$,

$$\phi : \partial X \xrightarrow{Z} Y.$$

Denote by v the dimension of the fibre Z and by h the dimension of the base Y of this fibration. We call X a *ϕ-manifold*. Fix a defining function x for the boundary ∂X.

Given these data we can define the Lie algebra of ϕ-vector fields on X by

$$^{\phi}\mathcal{V}(X) := \{V \in \Gamma(TX) \mid \quad V|_{\partial X} \quad \text{tangent to } Z \text{ and } \quad Vx = O(x^2)\}.$$

The space $^{\phi}\mathcal{V}(X)$ can be regarded as the space of sections of a vector bundle over X:

$$^{\phi}\mathcal{V}(X) \equiv \Gamma(^{\phi}TX),$$

where we call $^{\phi}TX$ the ϕ-tangent bundle to X. The dual bundle, $^{\phi}T^*X$, and all other associated tensor bundles will be distinguished by the symbol $^{\phi}$. Being isomorphic over the interior of X, the bundles $^{\phi}TX$ and TX are abstractly isomorphic over all of X. However, there is no *natural* isomorphism between them, and it will be shown that $^{\phi}TX$ carries some extra structure that TX does not have.

It will be convenient to recall that the fact that $\Gamma(^{\phi}TX)$ is a Lie algebra is equivalent to the fact that exterior derivation maps

$$d : \Gamma(^{\phi}T^*X) \to \Gamma(\Lambda^2 \, ^{\phi}T^*X).$$

Now, the inclusion $^{\phi}\mathcal{V}(X) \hookrightarrow \Gamma(TX)$ gives a natural map $^{\phi}TX \to TX$, which is not an isomorphism over the boundary, since its image consists of the vectors tangent to the fibres at the boundary. Restricted to the boundary we get a short exact sequence

$$0 \to {^{\phi}N}\partial X \to {^{\phi}TX}|_{\partial X} \to V\partial X \to 0, \tag{6}$$

where $V\partial X$ is the "vertical" bundle of vectors tangent to the fibres in the boundary. Also, this sequence is used to *define* the vector bundle $^{\phi}N\partial X$ over ∂X. The dual sequence

$$0 \to V^*\partial X \to {^{\phi}T^*X}|_{\partial X} \to {^{\phi}N^*}\partial X \to 0, \tag{7}$$

then defines the vector bundle $V^*\partial X$ over ∂X.

The existence of these subbundles over the boundary allows us to make sense of the vector bundles $[{}^\phi N\partial X]$ and $[{}^\phi V^*\partial X]$ over X (!) by defining their spaces of sections as

$$\Gamma(X, [{}^\phi N\partial X]) := \{W \in \Gamma(X, {}^\phi TX)| \quad W|_{\partial X, {}^\phi TX} \in {}^\phi N\partial X\} \tag{8}$$
$$\Gamma(X, [{}^\phi V^*\partial X]) := \{\xi \in \Gamma(X, {}^\phi T^*X)| \quad \xi|_{\partial X, {}^\phi TX} \in V^*\partial X\}. \tag{9}$$

Here the notation is intended to emphasize that restriction to the boundary is understood in the sense of elements in ${}^\phi TX$ or ${}^\phi T^*X$. Thus, the definition of these bundles really depend on the *pairs* $({}^\phi N\partial X, {}^\phi TX)$ and $(V^*\partial X, {}^\phi T^*X)$. We will include this into the notation more explicitly, whenever there is a possible ambiguity. Now the obvious inclusions of the spaces of sections give natural maps

$$[{}^\phi N\partial X] \longrightarrow {}^\phi TX, \qquad [{}^\phi V^*\partial X] \longrightarrow {}^\phi T^*X,$$

which, combined with restriction to the boundary in that sense, give

$$[{}^\phi N\partial X] \overset{|_{\partial X, {}^\phi TX}}{\longrightarrow} {}^\phi N\partial X, \qquad [{}^\phi V^*\partial X] \overset{|_{\partial X, {}^\phi T^*X}}{\longrightarrow} V^*\partial X.$$

Note that the bundle $[{}^\phi N\partial X]$ is well known as the "scattering bundle" over X (compare [Me2],[Me4]). Recalling that the "*b*-bundle" ${}^bTX$ is defined as the vector bundle whose space of sections consists of sections of TX, which are tangent to the boundary ([Me3]), we have

$$[{}^\phi N\partial X] = {}^{\mathrm{sc}}TX = x\,{}^bTX.$$

However, in our framework, sections in $[{}^\phi N\partial X]$ should rather be thought of as sections in ${}^\phi TX$ which are "asymptotically horizontal". In contrast to that, at this stage, we cannot make invariant sense of the notion of "asymptotically vertical" sections in ${}^\phi TX$!

Local Description

Let us emphasize that – once the boundary defining function x has been fixed – the constructions above are completely natural. It will be useful however, to have coordinate representations. So assume that we have chosen a collar neighborhood $C(\partial X) \cong \partial X \times [0, \varepsilon[_x$ of the boundary and a local trivialization of ϕ, so that locally near the boundary X looks like

$$U_X \cong U_Y \times Z \times [0, \varepsilon[\xrightarrow{\varphi} U_Y.$$

Then we may choose coordinates

- x : a boundary defining function of $Y \times \{0\}$ in $Y \times [0, \varepsilon[$, lifted via φ to U_X,
- $\mathbf{y} = (y_1, \dots, y_h)$: local coordinates on $U_Y \subset Y$, lifted to U_X via φ,
- $\mathbf{z} = (z_1, \dots, z_v)$: local coordinates on Z extended to U_X.

In these coordinates sections in ${}^\phi TX$ are locally spanned by

$$x^2\frac{\partial}{\partial x}, \quad x\frac{\partial}{\partial \mathbf{y}} = (x\frac{\partial}{\partial y^1}, \dots, x\frac{\partial}{\partial y^h}), \quad \frac{\partial}{\partial \mathbf{z}} = (\frac{\partial}{\partial z^1}, \dots, \frac{\partial}{\partial z^v}),$$

and the fact that ${}^\phi\mathcal{V}(X)$ is indeed a Lie algebra can be read off from this representation. Also elements of $\Gamma([{}^\phi N\partial X])$ are locally spanned by

$$x^2\frac{\partial}{\partial x}, \quad x\frac{\partial}{\partial \mathbf{y}} = (x\frac{\partial}{\partial y^1}, \dots, x\frac{\partial}{\partial y^h}), \quad x\frac{\partial}{\partial \mathbf{z}} = (x\frac{\partial}{\partial z^1}, \dots, x\frac{\partial}{\partial z^v}),$$

and sections of $[{}^\phi V\partial X]$ locally look like

$$x^3\frac{\partial}{\partial x}, \quad x^2\frac{\partial}{\partial \mathbf{y}} = (x^2\frac{\partial}{\partial y^1}, \dots, x^2\frac{\partial}{\partial y^h}), \quad \frac{\partial}{\partial \mathbf{z}} = (\frac{\partial}{\partial z^1}, \dots, \frac{\partial}{\partial z^v}).$$

A change of product structure and/or coordinates compatible with x can be regarded as a family $\Psi = (\Psi_x : \partial X \to \partial X)_{x\in[0,1]}$ of diffeomorphisms on ∂X such that Ψ_0 is a fibre bundle diffeomorphism of $\phi : \partial X \to Y$ fixing the base. Then

$$\Psi^* \frac{dy_j}{x} = \frac{dy_j}{x} + O(x^0)\langle\, d\mathbf{y},\, d\mathbf{z}\rangle, \quad \text{and} \quad \Psi_* \frac{\partial}{\partial z^j} = O(x^0)\langle \frac{\partial}{\partial \mathbf{z}}, x\frac{\partial}{\partial \mathbf{y}}\rangle,$$

which exemplifies the fact that we cannot identify "vertical" vectors within ${}^\phi TX$, for now!

Thus, as an additional piece of structure, *choose* a splitting of the Lie algebra sequence (6, 7), i.e. a decomposition

$${}^\phi TX|_{\partial X} = {}^\phi N\partial X \oplus V\partial X, \qquad {}^\phi T^*X|_{\partial X} = {}^\phi N^*\partial X \oplus V^*\partial X. \tag{10}$$

For instance, such a decomposition in the sense of Lie algebras could be obtained by choosing a product structure near the boundary as above.

The structure (10) will be *fixed once and for all*, the corresponding projections will be denoted by v and n. Among other things, this allows us to introduce the maps

$$\phi_* : {}^\phi TX|_{\partial X} \to {}^{\mathrm{sc}}TB|_Y, \qquad \phi^* : {}^{\mathrm{sc}}T^*B \to {}^\phi T^*X|_{\partial X},$$

over the boundary, with $B = Y \times [0, \infty[_x$. Also, we can now define the bundles $[{}^\phi V\partial X]$, the space of "asymptotically vertical" vectors in ${}^\phi TX$, and $[{}^\phi N^*\partial X]$ in analogy with our definition in (8, 9).

Remark 1.1 The reader should be aware of the following **pitfalls**. First

$$[{}^\phi N\partial X]^* = {}^{\mathrm{sc}}T^*X = \frac{1}{x}[{}^\phi V^*\partial X], \quad \text{thus in general} \quad [{}^\phi N\partial X]^* \neq [{}^\phi N^*\partial X]!!$$

Also, readers should convince themselves that in general

$$[{}^\phi N\partial X] \otimes [{}^\phi N\partial X] \neq [{}^\phi N\partial X \otimes {}^\phi N\partial X]_{{}^\phi TX \otimes {}^\phi TX}!!$$

The first bundle really encodes some second order behavior at the boundary of sections in ${}^\phi TX \otimes {}^\phi TX$, while the second bundle clearly only carries first order information.

The following Lemma restates the fact that we have chosen a decomposition compatible with the Lie algebra structure of ${}^\phi TX$:

Lemma 1.2 (a) $[\Gamma([{}^{\phi}V\partial X]),\Gamma([{}^{\phi}V\partial X])]\subset\Gamma([{}^{\phi}V\partial X])$

(b) $[\Gamma([{}^{\phi}N\partial X]),\Gamma([{}^{\phi}N\partial X])]\subset O(x)\Gamma([{}^{\phi}N\partial X])$

(c) $[\Gamma([{}^{\phi}N\partial X]),\Gamma({}^{\phi}TX)]\subset\Gamma([{}^{\phi}N\partial X])$. ■

Hoping that the reader has by now got used to the above way of coding the boundary (or asymptotic) behavior of sections into some bundle structure, we proceed with introducing one further modification which is going to be used in the next Section:

The vector bundle ${}^{\mathrm{e}}TX$ over X is defined (compare [Ma]) as the bundle whose space of sections is

$$\Gamma({}^{\mathrm{e}}TX)\equiv\{W\in\Gamma(TX)|\quad W|_{\partial X}\in V\partial X\}.$$

Again, this is a Lie algebra (thus exterior derivation is well defined on $\Lambda^{\mathrm{e}}T^*X$) and we have a map ${}^{\phi}TX\to{}^{\mathrm{e}}TX$. Thus, the sequence analogous to (6) is

$$0\to{}^{\mathrm{e}}N\partial X\to{}^{\mathrm{e}}TX|_{\partial X}\to V\partial X\to 0,$$

and it has a splitting map $V\partial X\to{}^{\mathrm{e}}TX$ induced by our splitting (10). In the above set of local coordinates, the bundle ${}^{\mathrm{e}}TX$ is spanned by

$$x\frac{\partial}{\partial x},\quad x\frac{\partial}{\partial\mathbf{y}}=(x\frac{\partial}{\partial y^1},\dots,x\frac{\partial}{\partial y^h}),\quad\frac{\partial}{\partial\mathbf{z}}=(\frac{\partial}{\partial z^1},\dots,\frac{\partial}{\partial z^v}).$$

We leave it to the reader to define the bundles $[{}^{\mathrm{e}}N\partial X]$, $[{}^{\mathrm{e}}V^*\partial X]$, $[{}^{\mathrm{e}}N^*\partial X]$, and $[{}^{\mathrm{e}}V\partial X]$ over X and to write down their local descriptions.

1.2 Exact ϕ-Metrics

A ϕ-metric is a fibre metric $g\in\Gamma(S^{2\phi}T^*X)$ on ${}^{\phi}TX$. We call it a *compatible* ϕ-metric when the decomposition at the boundary ${}^{\phi}TX|_{\partial X}={}^{\phi}N\partial X\oplus V\partial X$ is orthogonal. Another way to state this property is to demand that g has a decomposition

$$g\in\Gamma(S^2[{}^{\phi}N^*\partial X])+\Gamma(S^2[{}^{\phi}V^*\partial X]).$$

An *exact ϕ-metric* g_ϕ on ${}^{\phi}TX$ is a compatible ϕ-metric with a more refined decomposition of the form

$$g_\phi=\frac{d\,x^2}{x^4}+h+g_Z,\qquad\text{with}\tag{11}$$

$$\begin{aligned} h &\in \Gamma(S^2[{}^{\mathrm{e}}N^*\partial X])\subset\Gamma(S^2[{}^{\phi}N^*\partial X]),\quad h|_{\partial X,{}^{\mathrm{e}}TX}=\frac{1}{x^2}\phi^*g_B,\quad g_B\in\Gamma(S^2T^*B|_Y)\\ g_Z &\in \Gamma(S^2T^*X)\subset\Gamma(S^2[{}^{\phi}V^*\partial X]),\end{aligned}$$

where the notation $h|_{\partial X,{}^{\mathrm{e}}TX}$ means that we regard h as an element in $S^2\,{}^{\mathrm{e}}T^*X$ via the map $[{}^{\mathrm{e}}N^*\partial X]\to{}^{\mathrm{e}}T^*X$ when restricting to the boundary. We sometimes write $h=[\frac{1}{x^2}\phi^*g_B]$ to symbolize this construction. We also make use of the notation $g_{B,b}=\frac{dx^2}{x^2}+g_B$ for the underlying b-metric on B. Note that *the decomposition (11) is by no means unique.* For instance, only the $S^2V^*\partial X$-part of the restriction $g_Z|_{\partial X}$ makes invariant sense!

Product Metrics

The standard example of an exact ϕ-metric is a metric of type (2) as described in the Introduction. There we assumed that there is a fixed choice of product structure,

$$C(M) = M \times [0, \varepsilon[_x \xrightarrow{\varphi} Y \times [0, \varepsilon[_x,$$

in a collar neighborhood of the boundary ∂X and a choice of horizontal space HM. Especially, the fibration ϕ and the horizontal space of the boundary are extended to $C(M)$. Then a metric g_Y on Y can be lifted to a metric $\varphi^* g_Y$ on HM over $C(M)$. A metric, which over $C(M)$ is of the form

$$g_\phi = \left(\frac{dx}{x^2}\right)^2 + \frac{1}{x^2}\varphi^* g_Y + g_Z, \tag{12}$$

where g_Z is a metric in the fibres, will then be called a *product ϕ-metric* on X. It is easily seen that a local basis

$$V_1, \dots, V_v, \in VM \quad \text{and} \quad H_1, \dots H_h \in HM,$$

which is orthonormal w.r.t. $\phi^* g_Y + g_Z$, extends to a local basis

$$V_1, \dots, V_v, \quad xH_1, \dots xH_h, \quad x^2 \frac{\partial}{\partial x} = \mathsf{X}_\phi \quad \in {}^\phi TX$$

near ∂X, which is orthonormal w.r.t. g_ϕ and ∇^ϕ-parallel along $\frac{\partial}{\partial x}$. Using this, the connection and curvature of such a product metric can be calculated rather explicitly along the lines described in Appendix A.3.

The notion of an exact ϕ-metric is the natural generalization of metrics of this type to the "ϕ-setting". In fact, an exact ϕ-metric can be thought of as being asymptotically of the form (12). In this general setting, the techniques described above are not available to us and the geometry at the boundary is somewhat more intricate. However, we will sometimes refer to the product case to simplify our formulas.

The following Lemma lists the essential properties of (the inverse of) an exact ϕ-metric at the boundary:

Lemma 1.3 (Inverse of an Exact ϕ-Metric at ∂X)
Let $\alpha_i \in \Gamma(T^*X)$, $\beta_j \in \Gamma(T^*B|_Y)$ and choose extensions $[\frac{1}{x}\phi^*\beta_i] \in \Gamma([{}^{\mathrm{e}}N^*\partial X])$

(a) $g_\phi^{-1}(\frac{dx}{x^2}, \frac{dx}{x^2}) = 1 + \phi^* O(x^2)$

(b) $g_\phi^{-1}(\frac{dx}{x^2}, [\frac{1}{x}\phi^*\beta_1]) = \phi^* O(x)$

(c) $g_\phi^{-1}(\frac{dx}{x^2}, \alpha) = O(x^2)$

(d) $g_\phi^{-1}([\frac{1}{x}\phi^*\beta_1], [\frac{1}{x}\phi^*\beta_2]) = g_{B,b}^{-1}(\beta_1, \beta_2) = \phi^* O(x^0)$

(e) $g_\phi^{-1}([\frac{1}{x}\phi^*\beta_1], \alpha) = O(x)$.

(f) $g_\phi^{-1}(\alpha_1, \alpha_2) = O(x^0)$

Here the notation $\phi^* O(x^l)$ means that the term is $O(x^l)$ but the coefficient of x^l is of the form $\phi^* f$. ■

We will use this result to analyze the boundary behavior of the Levi-Civita connection for an exact ϕ-metric. First, we need to introduce some more notation. Denote by X_ϕ the vector field $g_\phi^{-1}(\frac{dx}{x^2}) \in \Gamma(x^2 TX)$.

Lemma 1.4 *The map $A \mapsto \widehat{A} := A - \frac{dx}{x^2}(A)\mathsf{X}_\phi$ maps $TX \to {}^bTX$ (and $xTX \to {}^{\mathrm{sc}}TX$) and has the following properties for $A, B \in TX$:*

(a) $dx(\widehat{A}) = O(x^2)$

(b) $\widehat{\widehat{A}} = \widehat{A} + O(x^0)\mathsf{X}_\phi$

(c) $g_\phi(\widehat{xA}, \widehat{xB}) = \phi^* g_B(A, B)|_{\partial X} + O(x)$.

(d) $\frac{1}{x^2}\phi^* g_B(\mathsf{X}_\phi, A) = dx(A) O(x^0)$, *where $O(x^0)$ is independent of A.* ■

The central geometric result in our setting is now stated in

Proposition 1.5 (Connection and Curvature of an Exact ϕ-Metric at ∂X) *The Levi-Civita connection for an exact ϕ-metric is a* true *connection, i.e.*

$$\nabla^\phi : \Gamma({}^\phi TX) \longrightarrow \Gamma(T^*X \otimes {}^\phi TX), \quad \text{and} \quad R^\phi \in \Gamma(\Lambda^2 T^*X, \mathrm{End}({}^\phi TX)).$$

Also, for $T, T_1, T_2 \in \Gamma({}^bTX)$, $N \in \Gamma(TX)$, $B \in \Gamma({}^\phi TX)$, $\omega \in \Gamma({}^{\mathrm{sc}}T^*B|_Y)$, $L \in \Gamma({}^{\mathrm{sc}}TB)$:

(a) $\nabla^\phi_N \frac{dx}{x^2}(B)|_{\partial X} = -\frac{1}{x^2}\phi^* g_B(\widehat{xN}, B)|_{\partial B}$

(b) $\nabla^\phi_N \mathsf{X}_\phi|_{\partial X, {}^\phi TX} = -\widehat{xN}$

(c) $\nabla^\phi_T : \Gamma([{}^\phi N\partial X]) \to \Gamma([{}^\phi N\partial X])$, *and* $\nabla^\phi_T : \Gamma([{}^\phi V\partial X]) \to \Gamma([{}^\phi V\partial X])$

(d) $\nabla^\phi_T|_{\partial X} = \mathsf{v} \circ \nabla^\phi_T \circ \mathsf{v} + \phi^* \nabla^{B,\mathrm{sc}}_T \circ \mathsf{n}$ *on* $\Gamma({}^\phi TX|_{\partial X})$

(e) $R^\phi(T_1, T_2)|_{\partial X} = \mathsf{v} \circ R^\phi(T_1, T_2) \circ \mathsf{v} \oplus \phi^* R^{B,sc}(T_1, T_2)$ *in* $\mathrm{End}({}^\phi TX|_{\partial X})$.

Especially $\nabla^\phi_T \phi^* L|_{\partial X} = 0$ in ${}^\phi TX$ and $\nabla^\phi_T \phi^* \omega|_{\partial X} = 0$ in ${}^\phi T^*X$ for vertical T.

Proof. The full proof of the mapping property of ∇^ϕ is tedious and we do not give it here. To see how to proceed, let us prove (a) in detail. First note that the expression on the RHS of (a) is well defined and independent of the ambiguities in the decomposition of g_ϕ. To prove the equality, let N, B be as above. Then, writing again $\mathsf{X}_\phi = g_\phi^{-1}(\frac{dx}{x^2})$ the Koszul formula yields

$$2\nabla^\phi_N \frac{dx}{x^2}(B) = (L_{\mathsf{X}_\phi} g_\phi)(N, B) + d\frac{dx}{x^2}(N, B) = (L_{\mathsf{X}_\phi} g_\phi)(N, B).$$

The Lie derivative of the metric can now be calculated using the decomposition of the metric and Cartan's formula:

$$\begin{aligned}
L_{\mathsf{X}_\phi}\left(\frac{dx}{x^2}\right)^2(N,B) &= \left(d\,\boldsymbol{\iota}(X_\phi)\frac{dx}{x^2}\otimes\frac{dx}{x^2}+\frac{dx}{x^2}\otimes d\,\boldsymbol{\iota}(X_\phi)\frac{dx}{x^2}\right)(N,B)\\
&= d(1+\phi^*O(x^2))(N)\frac{dx}{x^2}(B)+d(1+\phi^*O(x^2))(B)\frac{dx}{x^2}(N)=O(x)\\
L_{\mathsf{X}_\phi}g_Z(N,B) &= \mathsf{X}_\phi g_Z(N,B)-g_Z([\mathsf{X}_\phi,N],B)-g_Z(N,[\mathsf{X}_\phi,B])=O(x),
\end{aligned}$$

where we have used Lemma 1.3 (a) and (b), especially that $\mathsf{X}_\phi\in\Gamma(x^2TX)$.

The calculation of the contribution of the term h in the decomposition of g_ϕ is a little bit more involved: First, write

$$(L_{\mathsf{X}_\phi}h)(N,B)=\frac{1}{x^2}(L_{\mathsf{X}_\phi}x^2h)(N,B)-2(1+\phi^*O(x^2))h(xN,B).$$

We can assume that x^2h is (a sum of elements) of the form $[\phi^*\alpha]\otimes[\phi^*\beta]\in[x\,{}^{\mathrm{e}}N^*\partial X]^2$. Note that for elements in $[x\,{}^{\mathrm{e}}N^*\partial X]$ of this type we have $d[\phi^*\alpha],\ d[\phi^*\beta]\in\Lambda^2[x\,{}^{\mathrm{e}}N^*\partial X]$. Now

$$\begin{aligned}
\frac{1}{x^2}(L_{\mathsf{X}_\phi}x^2h)(N,B) &= \frac{1}{x^2}(L_{\mathsf{X}_\phi}[\phi^*\alpha]\otimes[\phi^*\beta])(N,B)\\
&= \frac{1}{x^2}\left[\,d[\phi^*\alpha](\mathsf{X}_\phi,N)[\phi^*\beta](B)+(N[\phi^*\alpha](\mathsf{X}_\phi))[\phi^*\beta](B)\right.\\
&\qquad\left.+\,d[\phi^*\beta](\mathsf{X}_\phi,B)[\phi^*\alpha](N)+(B[\phi^*\beta](\mathsf{X}_\phi))[\phi^*\alpha](N)\right],
\end{aligned}$$

and the first and the third summand here are $O(x)$ leaving us with

$$\begin{aligned}
\frac{1}{x^2}(L_{\mathsf{X}_\phi}x^2h)(N,B) &= \frac{1}{x^2}(N[\phi^*\alpha](\mathsf{X}_\phi))[\phi^*\beta](B)+\frac{1}{x^2}(B[\phi^*\beta](\mathsf{X}_\phi))[\phi^*\alpha](N)+O(x)\\
&= 2(Nx)[\phi^*\alpha](\frac{1}{x^2}\mathsf{X}_\phi)[\phi^*\beta](\frac{1}{x}B)+\frac{1}{x^2}[\phi^*\alpha](N)B\phi^*O(x^2)+O(x)\\
&= 2(Nx)h(\frac{1}{x^2}\mathsf{X}_\phi,xB)+O(x).
\end{aligned}$$

Summing the two results, the equation is proved.

The proof of (c) uses the same method. For $T\in\Gamma({}^bTX)$, $A\in\Gamma([{}^\phi N\partial X])$ and $V\in\Gamma([V\partial X])$ one can check that

$$2\langle\nabla^\phi_T V,A\rangle_\phi=(L_V g_\phi)(T,A),$$

and direct calculation, using the decomposition of g_ϕ as above, shows that the RHS is of order $O(x)$. Parts (d) and (e) follow easily. ■

1.3 ϕ-Differential Operators and the Dirac Operator D^d

Since $\Gamma({}^\phi TX)$ is a Lie algebra of vector fields, the associated space of ϕ-differential operators on X, $\mathrm{Diff}_\phi(X)$, can be defined in the usual way. Given vector bundles E,F this definition can be extended to ϕ-differential operators between these bundles, for example by using connections. Of interest to us will be the case of a *ϕ-Clifford bundle* , which is a hermitian vector bundle $E \to X$ with hermitian metric h^E, connection $\nabla^{E,\phi}$, and $(\nabla^{E,\phi}, \nabla^{\phi})$-parallel Clifford action

$$\mathbf{c}_\phi : \Gamma({}^{\phi}T^*X) \to \Gamma(\operatorname{End}(E)),$$

or, equivalently $\mathbf{c}_\phi \in \Gamma({}^{\phi}TX \otimes \operatorname{End}(E))$. The notational conventions used w.r.t. this Clifford action have been listed in Appendix A.1. The associated Dirac operator, given by

$$\mathsf{D}^{\phi} = \mathbf{c}_\phi \circ \nabla^{E,\phi},$$

is a ϕ-differential operator with coefficients in E. The restriction of this operator to the boundary ∂X is a vertical family of Dirac operators which we sometimes denote by $\mathsf{D}^{\phi,V}$.

Our real interest lies in "degenerate" metrics of the type $g_d = x^2 g_\phi$, where g_ϕ is an exact ϕ-metric, and the associated Dirac operator D^d. We will use the notation ^{d} to label geometric objects associated to this metric. Thus

$$g_d = x^2 g_\phi, \quad {}^{d}TX = x^{-1\phi}TX, \quad {}^{d}T^*X = x^{\phi}T^*X, \quad \mathsf{X}_d = g_d^{-1}(\frac{dx}{x})$$

and so on. Note that the sections of ${}^{d}TX$ may be singular as sections in TX and do *not* form a Lie algebra, which is why we have to consider the bundle ${}^{\phi}TX$ in the first place.

The Levi Civita connection

$$\nabla^d : \Gamma({}^{d}TX) \to \Gamma(T^*X \otimes {}^{d}TX), \quad \nabla^d : \Gamma({}^{d}T^*X) \to \Gamma(T^*X \otimes {}^{d}T^*X),$$

associated to the metric g_d, is a true connection and can be described in terms of the connection ∇^ϕ as in Appendix A.2. Thus for vectors $N \in TX$, $W \in \Gamma({}^{\phi}TX)$, and a differential form $\alpha \in \Gamma(\Lambda^{\phi}T^*X)$,

$$\begin{aligned} \nabla^d_N \frac{1}{x} W &= \frac{1}{x}\nabla^{\phi}_N W + \frac{1}{x} G(N)W,\ G(N)W := \frac{dx}{x^2}(W)xN - g_\phi(W, xN)\mathsf{X}_\phi \qquad (13) \\ \nabla^d_N\, x^{\mathsf{N}}\alpha &= x^{\mathsf{N}}\nabla^{\phi}_N\alpha + xG(N)\alpha,\ G(N)\alpha := \boldsymbol{\varepsilon}_\phi(xN)\,\boldsymbol{\iota}_\phi(\frac{dx}{x^2})\alpha - \boldsymbol{\varepsilon}(\frac{dx}{x^2})\,\boldsymbol{\iota}(xN)\alpha, \quad (14)\end{aligned}$$

where N is the counting operator. It will be useful to remember that for $N \in TX$

$$G(N) - G(\widehat{N}) = O(x^2)\operatorname{End}({}^{\phi}TX)$$

by Lemma 1.4. The map $G(N)$ is of order $O(x)$, if N is vertical at the boundary. In general $G(N)$ maps

$$G(N) : \Gamma([{}^{\phi}V\partial X]) \longrightarrow O(x)\Gamma({}^{\phi}TX), \qquad G(N) : \Gamma([{}^{\phi}N\partial X]) \longrightarrow \Gamma([{}^{\phi}N\partial X]). \qquad (15)$$

We can now introduce our basic objects of study. Given g_d as above, we consider a *d-Clifford bundle* , which is a hermitian vector bundle $E \to X$ with parallel hermitian metric h^E, connection $\nabla^{E,d} : \Gamma(E) \to \Gamma(T^*X \otimes E)$, and $(\nabla^{E,d}, \nabla^d)$-parallel Clifford action

$$\mathbf{c}_d : \Gamma({}^{d}T^*X) \to \Gamma(\operatorname{End}(E)).$$

The associated Dirac operator is

$$\mathsf{D}^d = \mathbf{c}_d \circ \nabla^{E,d}.$$

It is shown in Appendix A.2 – or can be checked by hand – that the definitions

$$\begin{aligned} \nabla_N^{E,\phi}\xi &:= \nabla_N^{E,d}\xi + \frac{1}{2}\mathbf{c}_\phi(\widehat{xN})\mathbf{c}_\phi(\frac{dx}{x^2})\xi, \qquad (16) \\ \mathbf{c}_\phi(\alpha) &= x\mathbf{c}_d(\alpha), \quad \mathbf{c}_\phi(A) = \frac{1}{x}\mathbf{c}_d(A), \quad \text{for } A \in {}^\phi TX, \alpha \in {}^\phi T^*X, \end{aligned}$$

endow the bundle E with the structure of a ϕ-Clifford bundle. The relationship between the Dirac operators D^d and D^ϕ is given by

$$\mathsf{D}^d = \mathbf{c}_d \circ \nabla^{E,d} = \frac{1}{x}\mathsf{D}^\phi + \frac{n-1}{2}\mathbf{c}_\phi(\frac{dx}{x^2}) = x^{-(n-1)/2}\frac{1}{x}\mathsf{D}^\phi x^{(n-1)/2}.$$

This will allow us to apply the analysis developed for ϕ-operators in the next Chapters to the operator D^d.

It should be emphasized that the d-Clifford structure on E and the Dirac operator D^d are regarded as our *basic objects*. For example, as explained in Appendix A.2, if D^d is the signature operator $d_d + d_d^*$ on $E = \Lambda\, {}^dT^*X$, then $\mathsf{D}^\phi = x\mathsf{D}^d$ *differs* from the conformally transformed signature operator by an endomorphism. Whenever we want to calculate explicit formulas involving the Clifford bundle we will therefore be careful to use the d-structure.

Let us end this Section with some useful remarks about the connection ∇^d and about the relationship between the curvatures R^ϕ and R^d:

Lemma 1.6 (Connection and Curvature for $\boldsymbol{g_d}$ at $\boldsymbol{\partial X}$) *Let $T \in \Gamma({}^bTX)$. Then*

(a) $\nabla_T^d \frac{dx}{x}(A)|_{\partial X} = 0$ *for* $A \in \Gamma({}^dTX)$

(b) $\nabla_T^d|_{\partial X} = \mathsf{v} \circ \nabla_T^d \circ \mathsf{v} + \phi^*\nabla_T^{B,b} \circ \mathsf{n}$ *on* $\Gamma({}^dTX|_{\partial X})$.

(c) $\frac{1}{x}\nabla_A^d \frac{dx}{x}(\frac{1}{x}B)|_{\partial X} = g_Z(A,B)$ *for* $A, B \in \Gamma({}^\phi TX)$

(d) $\nabla_\cdot^d \mathsf{X}_d|_{\partial X} = \mathsf{v}\cdot$, *the vertical projection* ${}^dTX|_{\partial X} \to {}^dV^*\partial X$

(e) *The tensor* $C = R^{{}^dT^*X} - R^{{}^\phi T^*X} \in \Gamma(\Lambda^2 T^*X \otimes \mathrm{End}({}^\phi T^*X))$ *is given on* tangent *vectors* T_1, T_2 *by*

$$C(T_1,T_2)|_{\partial X} = \boldsymbol{\varepsilon}_\phi(\widehat{xT_1})\,\boldsymbol{\iota}(\widehat{xT_2}) - \boldsymbol{\varepsilon}_\phi(\widehat{xT_2})\,\boldsymbol{\iota}(\widehat{xT_1})$$

Proof. To prove (a) we will use Lemma 1.5 (a). Thus, for $B \in \Gamma({}^\phi TX)$ we have

$$\begin{aligned} \nabla_T^d \frac{dx}{x}(\frac{1}{x}B) &= \nabla_T^\phi \frac{dx}{x^2}(B) + G(\widehat{T})\frac{dx}{x^2}(B/x) \\ &= -h(\widehat{xT},B) + g_\phi(\widehat{xT},B) - \frac{dx}{x^2}(\widehat{xT})\frac{dx}{x^2}(B) + O(x) \\ &= g_Z(\widehat{xB},T) + O(x) = O(x). \end{aligned}$$

To prove (c), we just have to observe that the $O(x)$-term in (a) is $O(x^2)$ if A, B are in $\Gamma({}^\phi TX)$. This should be clear from the proof of Lemma 1.5 (a). Part (d) is just a reformulation of (c).

To prove (e), let $A, B \in \Gamma({}^b TX)$. Then first

$$\begin{aligned} R^{{}^dT^*X}(A,B) - R^{{}^\phi T^*X}(A,B) &= [\frac{1}{x}\nabla^d_A x, \frac{1}{x}\nabla^d_B x] - \frac{1}{x}\nabla^d_{[A,B]}x - [\nabla^\phi_A, \nabla^\phi_B] + \nabla^\phi_{[A,B]} \\ &= [\nabla^\phi_A, G(B)] - [\nabla^\phi_B, G(A)] - G([A,B]) + [G(A), G(B)] \end{aligned}$$

The different terms in this expression are

$$\begin{aligned} [\nabla^\phi_A, G(B)] &= \frac{dx}{x}(A)G(B) + G(\nabla^\phi_A B) - \boldsymbol{\varepsilon}_\phi(xB)\,\boldsymbol{\iota}(\widehat{xA}) + \boldsymbol{\varepsilon}_\phi(\widehat{xA})\,\boldsymbol{\iota}(xB) && (17) \\ -[\nabla^\phi_B, G(A)] &= -\frac{dx}{x}(B)G(A) - G(\nabla^\phi_B A) + \boldsymbol{\varepsilon}_\phi(xA)\,\boldsymbol{\iota}(\widehat{xB}) - \boldsymbol{\varepsilon}_\phi(\widehat{xB})\,\boldsymbol{\iota}(xA) && (18) \\ [G(A), G(B)] &= [G(\widehat{A}), G(\widehat{B})] = \ldots = \boldsymbol{\varepsilon}_\phi(\widehat{xB})\,\boldsymbol{\iota}(\widehat{xA}) - \boldsymbol{\varepsilon}_\phi(\widehat{xA})\,\boldsymbol{\iota}(\widehat{xB}). \end{aligned}$$

Now the first summand in (17) and the last two summands in (18) give

$$\frac{dx}{x}(A)G(\widehat{B}) + \boldsymbol{\varepsilon}_\phi(xA)\,\boldsymbol{\iota}(\widehat{xB}) - \boldsymbol{\varepsilon}_\phi(\widehat{xB})\,\boldsymbol{\iota}(xA) = \boldsymbol{\varepsilon}_\phi(\widehat{xA})\,\boldsymbol{\iota}(\widehat{xB}) - \boldsymbol{\varepsilon}_\phi(\widehat{xB})\,\boldsymbol{\iota}(\widehat{xA})$$

and we get the same contribution fom the first summand in (18) and the last two summands in (17). Putting everything together gives the result.

Note that we have made extensive use of the fact that A and B are tangent to the boundary, since we needed every summand to be regular in $\mathrm{End}({}^\phi T^*X)$! ■

The expression for $C(T_1, T_2)$ only contains horizontal tensors. Therefore it maps

$$C(T_1, T_2) : [{}^\phi V^*\partial X] \longrightarrow x\,{}^\phi T^*X, \qquad C(T_1, T_2) : [{}^\phi N^*\partial X] \longrightarrow [{}^\phi N^*\partial X],$$

in general and even vanishes at the boundary whenever T_1 or T_2 are vertical. This means that $C|_{\partial X} \in \phi^*\Gamma(\Lambda^2 T^*Y \otimes \mathrm{End}({}^\phi T^*X))$. Using Lemma 1.5(e) we get the following

Corollary 1.7 *$R^d(T_1, T_2)|_{\partial X} = \mathsf{v} \circ R^\phi(T_1, T_2) \circ \mathsf{v} \oplus \phi^* R^{B,b}(T_1, T_2)$ for $T_1, T_2 \in {}^bTX$.* ■

1.4 Geometry of the Boundary

The essentials of the geometry of fibre bundles $M = \partial X \to Y$ are recalled in Appendix A.3. In this Section we want to show how the geometry of (X, g_ϕ) relates to the geometry of the boundary ∂X. To do this, a metric on ∂X has to be defined.

The first step is an extension-Lemma: Fix a normal vector field $\nu \in \Gamma(TX)$. Then any section $A \in \Gamma(\partial X, {}^\phi TX|_{\partial X})$ can be extended to the interior by parallel transport along ν. More concretely, in a small neighbourhood $U(\partial X)$ of the boundary, we write $\overline{A}$ for the extension of A into ${}^\phi TX$ such that $\frac{1}{x}\overline{A}$ is parallel w.r.t. ∇^d along ν. This choice of extension may seem a little odd at this stage, but will prove to be useful in Proposition 1.14 and Chapter 5. In any case it has the following nice properties

Lemma 1.8 (Special Extension) *Let ν, ν_1, ν_2 be normal vector fields in TX and index the associated extensions accordingly. Let $N \in \Gamma(TX)$ and $V \in \Gamma(\partial X, V\partial X)$, $A \in \Gamma(\partial X, {}^{\phi}N\partial X)$.*

(a) $\overline{A} \in \Gamma(U(\partial X), [{}^{\phi}N\partial X])$ *and* $\overline{V} \in \Gamma(U(\partial X), [{}^{\phi}V\partial X])$

(b) $\nabla^{d}_{N}\frac{1}{x}\overline{A} \in \Gamma(U(\partial X), [{}^{\mathrm{d}}N\partial X])$ *and* $\nabla^{d}_{N}\frac{1}{x}\overline{V} \in \Gamma(U(\partial X), [{}^{\mathrm{d}}V\partial X])$.

(c) $\nabla^{\phi}_{N}\overline{A} \in \Gamma(U(\partial X), [{}^{\phi}N\partial X])$ *and* $\nabla^{\phi}_{N}\overline{V} \in \Gamma(U(\partial X),\ [{}^{\phi}V\partial X])$.

(d) $\frac{1}{x}(\overline{A}^1 - \overline{A}^2) \in \Gamma(U(\partial X), [{}^{\phi}N\partial X])$ *and* $\frac{1}{x}(\overline{V}^1 - \overline{V}^2) \in \Gamma(U(\partial X), [{}^{\phi}V\partial X])$

(e) $\frac{1}{x}\overline{A}|_{\partial X} \in {}^{b}TX|_{\partial X}$ *is independent of the choice of* ν.

Proof. (a) is clear by construction, (b) and (c) follow from Lemma 1.5(c) and (15). Consider (d): Taking two extensions $\overline{V}^1, \overline{V}^2 \in \Gamma({}^{\phi}TX)$ of $V \in \Gamma(V\partial X)$ it is clear that $\overline{V}^1 - \overline{V}^2 \in O(x)\Gamma({}^{\phi}TX)$. Thus

$$\frac{1}{x}(\overline{V}^1 - \overline{V}^2) = \frac{1}{Nx}\nabla^{\phi}_{N}(\overline{V}^1 - \overline{V}^2) \in \Gamma([{}^{\phi}V\partial X])$$

by (c). ■

As a consequence, this construction yields a canonical choice of horizontal space in ${}^{b}TX|_{\partial X}$ as the span

$${}^{b}H\partial X = \langle \frac{1}{x}\overline{A}|_{\partial X, {}^{b}TX} \mid \quad A \in \Gamma(\partial X, {}^{\phi}N\partial X)\rangle.$$

The induced horizontal and vertical projections in $T\partial X = TM$ and ${}^{b}TX|_{\partial X}$ will be denoted by h and v. Using these and the boundary defining function x we can now define the maps

$$\begin{aligned}
\delta_h \quad &: \quad {}^{\phi}TX|_{\partial X} \longrightarrow {}^{b}TX|_{\partial X} \qquad A \longmapsto [\frac{1}{x}\overline{\mathsf{n}A} + \mathsf{v}A]|_{\partial X, {}^{b}TX} \\
\delta_h^{-1} \quad &: \quad {}^{b}TX|_{\partial X} \longrightarrow {}^{\phi}TX|_{\partial X} \qquad A \longmapsto x[\mathsf{h}A]_b|_{\partial X, {}^{\phi}TX} + [\mathsf{v}A]_b|_{\partial X, TX} \\
& \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad = x[A]_b|_{\partial X, {}^{\phi}TX} + [\mathsf{v}A]_b|_{\partial X, TX},
\end{aligned}$$

where $[A]_b$ is an extension of $A \in {}^{b}TX|_{\partial X}$ to ${}^{b}TX$. The mapping properties of δ_h and the justification of the "inverse"-notation are given in

Lemma 1.9 (a) $\delta_h \in \Gamma(\partial X, {}^{\phi}T^*X \otimes {}^{b}TX|_{\partial X})$

(b) $\delta_h \circ \delta_h^{-1} = \mathrm{Id}_{{}^{b}TX}$, *and* $\quad \delta_h^{-1} \circ \delta_h = \mathrm{Id}_{{}^{\phi}TX}$

(c) $\delta_h^{-1}(TM) = \langle \frac{dx}{x^2}\rangle^{\perp} \subset {}^{\phi}TX$, *and* $\quad \delta_h^{*}(T^*M) = {}^{e}T^*X$. ■

We can now simply define the metric on the fibres of ${}^bTX|_M \to M$ as

$$g_{M,b} = (\delta_h^{-1})^*(g_\phi).$$

Using the identification of TM with the kernel of $\frac{dx}{x}$ in ${}^bTX|_{\partial X}$ we can also restrict the metric $g_{M,b}$ to the metric g_M on M. However, since it involves no extra work, we prefer to stick with the bundle ${}^bTX|_M$. Thus, the Koszul formula defines the "Levi-Civita"-connection $\nabla^{M,b}$ associated to this metric. To be able to use this definition in calculations, we need a more detailed description of $g_{M,b}$. Parts (b) and (c) of the following Lemma will therefore be of particular importance:

Lemma 1.10 (Metric on the Boundary)

(a) $g_{M,b}^{-1} = \delta_h(g_\phi^{-1})$

(b) $g_{M,b}(V,T) = g_\phi(\overline{V},T)$ *for* $V \in VM$, $T \in {}^bTX$

(c) $g_{M,b}(T_1,T_2) = g_\phi(\mathsf{v}T_1,\mathsf{v}T_2) + g_d(T_1,T_2)$ *for* $T_1, T_2 \in {}^bTX$

(d) $\frac{1}{x}\overline{A}|_{\partial X}$ *for* $A \in \Gamma({}^\phi N\partial X)$ *is horizontal w.r.t.* $g_{M,b}$.

Proof. Part (a) is straightforward. To prove (b), we first show that $g_\phi(\overline{V},T)$ is well defined by showing that it is well defined for the different terms in the decomposition of g_ϕ. To do this note first that

$$N\frac{dx}{x^2}(\overline{V}) = \nabla^d_N \frac{dx}{x}(\frac{1}{x}\overline{V}) + \frac{dx}{x}(\nabla^d_N \frac{1}{x}\overline{V}) = O(x),$$

i.e. the expression $\frac{dx}{x^2}(\overline{V})$ is of order $O(x^2)$. Thus

$$\frac{dx}{x^2} \otimes \frac{dx}{x^2}(\overline{V},T) = O(x^0)\, dx(T) = O(x).$$

To see that the expression $h(\overline{V},T)$ only depends on $V, T|_{\partial X}$, note that the extension $\overline{V}$ pairs with any element $\varphi \in \Gamma([{}^{\mathrm{e}}N^*\partial X])$ to $O(x)$ and there is a natural map $x[{}^{\mathrm{e}}N^*\partial X] \to T^*X$. But the pairing of T with any element in $\Gamma(\partial X, T^*X)$ clearly only depends on $T|_{\partial X}$. The corresponding statement for g_Z is clear.

To prove the equality in (b) note that it is trivially true for vertical T. On the other hand

$$\langle \overline{V}, \mathsf{h}[T]\rangle_\phi = \langle \overline{V}, \frac{1}{x}\overline{A}\rangle_\phi = \frac{1}{Nx} N\langle \frac{1}{x}\overline{V}, \frac{1}{x}\overline{A}\rangle_d = \frac{1}{Nx}\left[\langle \nabla^d_N \frac{1}{x}\overline{A}, \frac{1}{x}\overline{V}\rangle_d + \langle \frac{1}{x}\overline{A}, \nabla^d_N \frac{1}{x}\overline{V}\rangle_d\right]$$

is 0 by definition of the extension. This also proves (c) and (d). ∎

As an aside, note that part (c) of the Lemma implies that the metric $g_{M,b}$ (or g_M) is really of the type (114) required in Appendix A.3, since

$$g_{M,b}(\mathsf{h}T_1, \mathsf{h}T_2) = g_d(\mathsf{h}T_1, \mathsf{h}T_2) = \phi^* g_{B,b}(T_1, T_2).$$

The following Lemma lists some ways to calculate the horizontal and vertical projections

Lemma 1.11 *Let* $\alpha \in \Gamma({}^bT^*X)$*,* $N \in \Gamma(TX)$ *and* $T \in \Gamma({}^bTX) = \Gamma([{}^dN^*\partial X])$*. Then* $x\alpha$ *is a section in* $[{}^dV^*\partial X]$ *and*

(a) $\nabla^d_N(x\alpha)(T)|_{\partial X} = \nabla^\phi_N\alpha(xT)|_{\partial X} = \, dx(N)\alpha(\mathsf{h}T)|_{\partial X}$.

(b) $x\nabla^\phi_N T|_{\partial X, {}^bTX} = - \, dx(N)\mathsf{h}T$

(c) $x\nabla^d_N T|_{\partial X, {}^bTX} = \, dx(N)\mathsf{v}T$

Proof. We use the method developed in Section 1.2. For simplicity, assume that $\alpha \in \Gamma(T^*X)$. Also write $A := (g_\phi)^{-1}(\alpha)$. Then

$$\begin{aligned} \nabla^d_N(x\alpha)(T) &= \nabla^\phi_N\alpha(xT) + O(x) \\ &= (L_Ag_\phi)(N, xT) + \, d\alpha(N, xT) + O(x) = (L_Ag_\phi)(N, xT) + O(x). \end{aligned}$$

The different parts of the Lie derivative of the metric can be calculated as in Lemma 1.5:

$$\begin{aligned} L_A(\frac{dx}{x^2} \otimes \frac{dx}{x^2})(N, xT) &= O(x^{-1})\, dx \otimes \, dx(N, xT) + O(x) = O(x) \\ L_Ag_Z(N, xT) &= O(x) \\ L_Ah(xN, T) &= (Nx)h(A, T) = (Nx)(\alpha(T) - g_Z(A, T)) = (Nx)\alpha(\mathsf{h}T) \end{aligned}$$

This proves the claim for $\alpha \in \Gamma(T^*X)$. It is easy to see that it also holds for forms of the type $\alpha = f\frac{dx}{x}$. Parts (b) and (c) are just reformulations of (a). ■

We now want to describe $\nabla^{M,b}$ and its associated tensors in terms of g_d and ∇^d. We start with the following

Definition 1.12 (a) *The curvature* $\Omega_\phi \in C^\infty(\partial X, \Lambda^2\, {}^\phi T^*X \otimes V\partial X)$ *is defined by*

$$\Omega_\phi(A, B) := \mathsf{v}[\frac{1}{x}\overline{\mathsf{n}A}, \frac{1}{x}\overline{\mathsf{n}B}]|_{\partial X} = \Omega_M(\frac{1}{x}\overline{\mathsf{n}A}, \frac{1}{x}\overline{\mathsf{n}B}), \quad \textit{for } A, B \in {}^\phi TX$$

(b) *The tensor* $S_\phi \in C^\infty(\partial X, V^*\partial X \otimes {}^\phi N^*\partial X \otimes V\partial X)$ *is defined by*

$$S_\phi(V)A := \mathsf{v}\frac{1}{x}\nabla^\phi_{\overline{V}}\overline{A}|_{\partial X}, \quad \textit{for } V \in V\partial X,\ A \in {}^\phi N\partial X.$$

It could be called the ϕ*-second fundamental form.*

(c) *A variant of (b) is* $S_d \in C^\infty(\partial X, V^*\partial X \otimes {}^dN^*\partial X \otimes V\partial X)$*, defined by*

$$S_d(V)\frac{1}{x}A := \mathsf{v}\nabla^d_{\overline{V}}\frac{1}{x}\overline{A}|_{\partial X} \overset{(13)}{=} S_\phi(V)A + \frac{dx}{x^2}(A)V, \quad \textit{for } V \in V\partial X,\ A \in {}^\phi N\partial X.$$

(d) $B_\phi \in C^\infty(\partial X, \Lambda^2V^*\partial X \otimes {}^\phi N\partial X)$ *is defined by*

$$B_\phi(V, W) := \mathsf{n}\frac{1}{x}\left([\overline{V}, \overline{W}] - \overline{[V, W]}\right), \quad \textit{for } V, W \in V\partial X.$$

The reader should verify that B_ϕ is indeed a well defined tensor: Since $[\overline{V},\overline{W}]$ and $\overline{[V,W]}$ have the same boundary value as elements in ${}^{\phi}TX$, their difference

$$[\overline{V},\overline{W}]-\overline{[V,W]} \quad \text{lies in} \quad x^{\phi}TX. \tag{19}$$

The nonvanishing of the tensor B_ϕ is linked the fact that g_ϕ might not be a product ϕ-metric near the boundary! In the product case (19) vanishes near the boundary and therefore $B_\phi=0$.

For a general exact ϕ-metric we have the following analogue of Lemma A.3:

Lemma 1.13 *Let $T_0\in\Gamma(TM), T_1,T_2\in\Gamma(M,{}^{b}TX|_M)$. Then*

$$\begin{aligned}\langle\nabla^{M,b}_{T_0}T_1,T_2\rangle_{M,b}=&\langle\nabla^{\phi}_{T_0}\overline{\mathsf{v}T_1},\overline{\mathsf{v}T_2}\rangle_\phi+\langle\nabla^{d}_{T_0}\overline{\mathsf{h}T_1},\overline{\mathsf{h}T_2}\rangle_d\\ &+\frac{1}{2}\left[\langle\Omega_\phi(\overline{\mathsf{h}T_0},\overline{\mathsf{h}T_1}),\overline{\mathsf{v}T_2}\rangle_\phi-\langle\Omega_\phi(\overline{\mathsf{h}T_1},\overline{\mathsf{h}T_2}),\overline{\mathsf{v}T_0}\rangle_\phi-\langle\Omega_\phi(\overline{\mathsf{h}T_2},\overline{\mathsf{h}T_0}),\overline{\mathsf{v}T_1}\rangle_\phi\right]\\ &+\langle S_\phi(\overline{\mathsf{v}T_0})\overline{\mathsf{h}T_1},\overline{\mathsf{v}T_2}\rangle_\phi-\langle S_\phi(\overline{\mathsf{v}T_0})\overline{\mathsf{h}T_2},\overline{\mathsf{v}T_1}\rangle_\phi\\ &+\frac{1}{2}\left[\langle B_\phi(\overline{\mathsf{v}T_1},\overline{\mathsf{v}T_2}),x\overline{\mathsf{h}T_0}\rangle_\phi-\langle B_\phi(\overline{\mathsf{v}T_2},\overline{\mathsf{v}T_0}),x\overline{\mathsf{h}T_1}\rangle_\phi-\langle B_\phi(\overline{\mathsf{v}T_0},\overline{\mathsf{v}T_1}),x\overline{\mathsf{h}T_2}\rangle_\phi\right]\end{aligned}$$

Proof. This is an application of the Koszul formula using Lemma 1.10(c). We only give an example of how the calculation differs from the one in Lemma A.3:

$$\begin{aligned}2\langle S_\phi(V)A,W\rangle_\phi &= -\overline{W}\langle\overline{V},\frac{1}{x}\overline{A}\rangle_\phi+\overline{V}\langle\frac{1}{x}\overline{A},\overline{W}\rangle_\phi+\frac{1}{x}\overline{A}\langle\overline{W},\overline{V}\rangle_\phi\\ &\qquad -\langle\overline{V},[\frac{1}{x}\overline{A},\overline{W}]\rangle_\phi+\langle\frac{1}{x}\overline{A},[\overline{W},\overline{V}]\rangle_\phi+\langle\overline{W},[\overline{V},\frac{1}{x}\overline{A}]\rangle_\phi\\ &= -\overline{W}\langle\overline{V},\frac{1}{x}\overline{A}\rangle_{M,b}+\overline{V}\langle\frac{1}{x}\overline{A},\overline{W}\rangle_{M,b}+\frac{1}{x}\overline{A}\langle\overline{W},\overline{V}\rangle_{M,b}\\ &\quad -\langle\overline{V},[\frac{1}{x}\overline{A},\overline{W}]\rangle_{M,b}+\langle\frac{1}{x}\overline{A},[\overline{W},\overline{V}]\rangle_{M,b}+\langle\overline{W},[\overline{V},\frac{1}{x}\overline{A}]\rangle_{M,b}+\langle\frac{1}{x}\overline{A},[\overline{W},\overline{V}]\rangle_\phi\\ &= 2\langle S_M(V)\frac{1}{x}\overline{A},W\rangle_{M,b}+\langle\frac{1}{x}\overline{A},[\overline{W},\overline{V}]\rangle_\phi\\ &= 2\langle S_M(V)\frac{1}{x}\overline{A},W\rangle_{M,b}-\langle B_\phi(V,W),A\rangle_\phi,\end{aligned}$$

and so on. ■

As a special case of this Lemma, note that

$$\langle\nabla^{M,b}_{T_0}\mathsf{v}T_1,\mathsf{v}T_2\rangle_{M,b}=\langle\nabla^{\phi}_{T_0}\overline{\mathsf{v}T_1},\overline{\mathsf{v}T_2}\rangle_\phi+\frac{1}{2}\langle B_\phi(\overline{\mathsf{v}T_1},\overline{\mathsf{v}T_2}),x\overline{\mathsf{h}T_0}\rangle_\phi. \tag{20}$$

This will play a minor role in the calculation of (102).

We can now calculate parts of the curvature R^d at the boundary more precisely.

Proposition 1.14 *Let $N\in\Gamma(TX)$ and $W,A,B\in\Gamma({}^{\phi}TX)$. Then at the boundary*

(a) $\langle R^d(W,N)A,B\rangle_\phi=\langle R^d(W,N)\tfrac{1}{x}A,\tfrac{1}{x}B\rangle_d$

$$=\langle R^d(W,N)\mathsf{v}A,\mathsf{v}B\rangle_\phi+\ dx(N)[\langle S_d(W)\tfrac{1}{x}\mathsf{n}B,\mathsf{v}A\rangle_\phi-\langle S_d(W)\tfrac{1}{x}\mathsf{n}A,\mathsf{v}B\rangle_\phi]$$

(b) $\langle R^d(W,N)\mathsf{n}A, \mathsf{n}B\rangle_\phi$ *is* $O(x)$ *and the corresponding coefficient depends on the choice of extension of* $\mathsf{n}A$, $\mathsf{n}B$ *into the interior. Still, one has*

$$\langle (\nabla^d_N R^d)(W,N)\mathsf{n}A, \mathsf{n}B\rangle_\phi = (\,dx(N))^2 \langle \Omega_\phi(A,B), W\rangle_\phi.$$

Proof. First note that the LHS in (a) is $O(x)$, whenever $W \in \Gamma([{}^\phi N\partial X])$. We thus can assume right away that $W \in \Gamma([{}^\phi V\partial X])$. Then, using the special extension, w.r.t. the normal vector field N, we get

$$\begin{aligned}
\langle R^d(W,N)\frac{1}{x}\mathsf{n}A, \frac{1}{x}\mathsf{v}B\rangle_d &= \langle R^d(\overline{W},N)\frac{1}{x}\overline{\mathsf{n}A}, \frac{1}{x}\overline{\mathsf{v}B}\rangle_d = -N\langle \nabla^d_{\overline{W}}\frac{1}{x}\overline{\mathsf{n}A}, \frac{1}{x}\overline{\mathsf{v}B}\rangle_d \\
&= -\,dx(N)\langle \nabla^d_{\overline{W}}\frac{1}{x}\overline{\mathsf{n}A}, \overline{\mathsf{v}B}\rangle_\phi = -\,dx(N)\langle S_d(\overline{W})\frac{1}{x}\overline{\mathsf{n}A}, \overline{\mathsf{v}B}\rangle_\phi.
\end{aligned}$$

Furthermore, the LHS,

$$\langle R^d(W,N)\frac{1}{x}A, \frac{1}{x}B\rangle_d = \langle R^d(A,N)\frac{1}{x}B, \frac{1}{x}W\rangle_d - \langle R^d(B,N)\frac{1}{x}A, \frac{1}{x}W\rangle_d, \tag{21}$$

is of order $O(x)$, whenever A, B are in $\Gamma([{}^\phi N\partial X])$. This finishes the proof of (a). To prove (b), assume that A, B are in $\Gamma([{}^\phi N\partial X])$. Then

$$\begin{aligned}
\langle (\nabla^d_N R^d)(W,N)\frac{1}{x}A, \frac{1}{x}B\rangle_d &= N\langle R^d(\overline{W},N)\frac{1}{x}\overline{A}, \frac{1}{x}\overline{B}\rangle_d \\
&= (Nx)\left[\langle R^d(\frac{1}{x}\overline{A},N)\frac{1}{x}\overline{B}, \frac{1}{x}\overline{W}\rangle_d - \langle R^d(\frac{1}{x}\overline{B},N)\frac{1}{x}\overline{A}, \frac{1}{x}\overline{W}\rangle_d\right] \\
&= (Nx)N\left[\langle \nabla^d_{\frac{1}{x}\overline{A}}\frac{1}{x}\overline{B}, \frac{1}{x}\overline{W}\rangle_d - \langle \nabla^d_{\frac{1}{x}\overline{B}}\frac{1}{x}\overline{A}, \frac{1}{x}\overline{W}\rangle_d\right] \\
&\qquad + (Nx)\langle \nabla^d_{[\frac{1}{x}\overline{A},N]}\frac{1}{x}\overline{B} - \nabla^d_{[\frac{1}{x}\overline{B},N]}\frac{1}{x}\overline{A}, \frac{1}{x}\overline{W}\rangle_d \\
&= (Nx)^2\langle [\frac{1}{x}\overline{A}, \frac{1}{x}\overline{B}], \overline{W}\rangle_\phi = (Nx)^2\langle \Omega_\phi(A,B), W\rangle_\phi.
\end{aligned}$$

This finishes the proof. ■

Using Definition 1.12 and (the proof of) Lemma 1.13, the RHS in Proposition 1.14(a) can be further decomposed into contributions from the intrinsic and the extrinsic geometry of the boundary fibration. For instance, the term

$$\langle R^d(W,\nu)\mathsf{v}A, \mathsf{v}B\rangle_\phi|_{\partial X} = \langle R^\phi(W,\nu)\mathsf{v}A, \mathsf{v}B\rangle_\phi|_{\partial X},$$

where $\nu = \frac{1}{x}\mathsf{X}_d$, should be considered as a tensor describing some aspect of the extrinsic geometry, since it vanishes whenever g_ϕ is a ϕ-product metric.

2 The ϕ-Calculus

As before, X denotes a manifold with fibred boundary. By the Schwartz kernel theorem, continuous operators from $\dot{C}^\infty(X)$ to $C^{-\infty}(X)$ are represented by elements of $C^{-\infty}(X^2)$. In this Chapter we will give an overview of the ideas developed in [MaMe3] and [Me0], [Me3] for the analysis of such operators whose kernels are distributions conormal to the diagonal and to the boundaries in a certain sense. The composition formula for these ϕ-pseudodifferential operators proved in Section 2.3 is an extension of the corresponding result in [MaMe3].

2.1 The b- and the ϕ-Blow up

Denote by β_L, β_R the left and right projections from X^2 to X. Given two sets of local coordinates on X near the boundary $x, \mathbf{y}, \mathbf{z}$ and $x', \mathbf{y}', \mathbf{z}'$, as in Section 1.1 we get coordinates on X^2 by lifting the first set from the left and the second set from the right to X^2:

$$x \triangleq \beta_L^* x, \quad \mathbf{y} \triangleq \beta_L^* \mathbf{y}, \quad \mathbf{z} \triangleq \beta_L^* \mathbf{z}, \qquad \text{and} \qquad x' \triangleq \beta_R^* x', \quad \mathbf{y}' \triangleq \beta_R^* \mathbf{y}', \quad \mathbf{z}' \triangleq \beta_R^* \mathbf{z}'.$$

A priori the two coordinate sets on X can be completely independent. It will usually be assumed though, that $x = x'$ is the fixed boundary defining function on X. Sometimes, depending of the geometrical situation – see the proof of Lemma 2.1 below, we also assume that $\mathbf{y} = \mathbf{y}'$ and $\mathbf{z} = \mathbf{z}'$ on X. In that case, we will say that the coordinates $\mathbf{y}, \mathbf{y}'$ or $\mathbf{z}, \mathbf{z}'$ on X^2 are *paired*.

For an easier analysis of the different classes of pseudodifferential operators on X we introduce: The b-kernel space is defined as the blow up (compare [Me3])

$$X_b^2 := [X^2, \partial X \times \partial X].$$

The corresponding blow down map is denoted $\beta_b : X_b^2 \to X^2$, and the induced left and right projections are $\beta_{b,L}$ and $\beta_{b,R}$. We write

$$\triangle_b := \overline{\beta_b^{-1}(\triangle - \partial\triangle)}, \quad \mathrm{bf} := \beta_b^{-1}(\partial X \times \partial X).$$



Figure 1

The face bf is diffeomorphic to $\partial X \times \partial X \times [-1,1]_\sigma$. Here, the coordinate σ is given by $\sigma = \frac{x-x'}{x+x'}$. The *fibre diagonal* $\partial X \times_Y \partial X$ is the preimage of the diagonal $\triangle_Y$ under the map $(\phi,\phi) : \partial X \times \partial X \to Y \times Y$. Its lift in bf is denoted by

$$F_\phi := \beta_b^{-1}(\partial X \times_Y \partial X) \times \{\sigma = 0\} \cong \partial X \times_Y \partial X.$$

Note that this definition depends on our (fixed) choice for the boundary defining function x. Blowing up F_ϕ in X_b^2 yields the kernel space for the ϕ-pseudodifferential operators (compare [MaMe3])

$$X_\phi^2 := [X_b^2, F_\phi].$$

Again, we write $\beta_\phi : X_\phi^2 \to X^2$ for the blow down map and $\beta_{\phi,L}$ and $\beta_{\phi,R}$ for the induced left and right projections onto X. Also, the lifted diagonal and the blown up faces are denoted by

$$\triangle_\phi := \overline{\beta_\phi^{-1}(\triangle - \partial\triangle)}, \quad \mathrm{ff} := \beta_{\phi,b}^{-1} F_\phi, \quad \phi\mathrm{bf} := \overline{\beta_{\phi,b}^{-1}(\mathrm{bf} - F_\phi)}.$$



Figure 2

As a general rule, given a boundary face F, we denote by ρ_F a boundary defining function for F. The following Lemma describes the fundamental property of the blow up X_ϕ^2:

Lemma 2.1 (a) *Restricted to $\triangle_\phi$ the spaces $\beta_{\phi,R}^{*}\,{}^{\phi}\mathcal{V}$ and $\beta_{\phi,L}^{*}\,{}^{\phi}\mathcal{V}$ each span the normal bundle $N\triangle_\phi$. Especially, recalling that $\triangle_\phi$ is canonically diffeomorphic to X, we have $TX_\phi^2|_{\triangle_\phi} \equiv TX \oplus {}^{\phi}TX$.*

(b) *$\beta_{\phi,R}^{*}W$, for $W \in \Gamma({}^{\phi}TX)$, is a ϕ-vector field at* lf *and* ϕbf *and is a b-vector field at* ff *and* rf.

(c) $\overset{\circ}{\mathrm{ff}}$ *is canonically diffeomorphic to ${}^{\phi}N\partial X \times_Y \partial X$ and* ff *to* $\mathrm{RC}_{\mathrm{fib}}({}^{\phi}N\partial X \times_Y \partial X)$.

Proof. (a) and (b) can be proved by calculating the lift of vector fields in local coordinates. First, taking local coordinates **near the corner in X^2**, we can introduce local coordinates on X_b^2, projective w.r.t. x':

$$s = \frac{x}{x'}, \quad x', \quad \mathbf{y}, \quad \mathbf{y}', \quad \mathbf{z}, \quad \mathbf{z}' \tag{22}$$

These coordinates are valid near bf but away from rf. There we calculate

$$ds = \frac{dx}{x'} - \frac{1}{sx'}\,dx' \quad \text{and} \quad {\beta_{b,R}}^* x^2 \frac{\partial}{\partial x} = x' s^2 \frac{\partial}{\partial s}, \quad {\beta_{b,R}}^* x \frac{\partial}{\partial \mathbf{y}} = x' s \frac{\partial}{\partial \mathbf{y}}.$$

Near the fibre diagonal $\boldsymbol{F_\phi}$ we can assume that the coordinates $\mathbf{y}$ and $\mathbf{y}'$ are paired, thus F_ϕ is described by $\{x' = 0, s = 1, \mathbf{y} = \mathbf{y}'\}$. Near ff, but away from bf, we can introduce coordinates on X^2_ϕ, projective w.r.t. x':

$$x', \quad S = \frac{s-1}{x'}, \quad \mathbf{Y} = \frac{\mathbf{y} - \mathbf{y}'}{x'}, \quad \mathbf{y}', \quad \mathbf{z}, \quad \mathbf{z}' \quad \text{and} \tag{23}$$

$$dS = \frac{ds}{x'} - \frac{s-1}{(x')^2}\,dx', \quad d\mathbf{Y} = \frac{d\mathbf{y} - \,d\mathbf{y}'}{x'} - \frac{\mathbf{y} - \mathbf{y}'}{(x')^2}\,dx'.$$

From this we get:

$${\beta_{\phi,R}}^* x^2 \frac{\partial}{\partial x} = (1 + x'S)^2 \frac{\partial}{\partial S}, \quad {\beta_{\phi,R}}^* x \frac{\partial}{\partial \mathbf{y}} = (1 + x'S) \frac{\partial}{\partial \mathbf{Y}}. \tag{24}$$

From these calculations (b) is obvious. Finally, near the lifted diagonal $\triangle_\phi$ we can add the condition that the coordinates $\mathbf{z}$ and $\mathbf{z}'$ are also paired. Then $\triangle_\phi$ is just $\{x' = 0, S = 0, \mathbf{Y} = 0\}, \mathbf{z} = \mathbf{z}'$ and the assertion of (a) also follows from (24).

For (c), we just mention that the map from ${}^\phi N\partial X \times_Y \partial X$ to the interior of ff is given as follows. A nonzero element $W \in \Gamma({}^\phi N\partial X)$ lifts via $\beta_{\phi,L}$ to a nonvanishing vector field $\beta^*_{\phi,L}W$ on ff, which generates a flow $\Phi_W(t)$ on that face. This flow is identified with the linear flow generated by W in the fibres of ${}^\phi N\partial X$. ■

The *small* calculus of b-pseudodifferential operators on X is defined as the space

$$\Psi^m_{b,\mathrm{cl}}(X) := \dot{C}^\infty_{\mathrm{bf}} I^m_{\mathrm{cl}}(X^2_b, \triangle_b; \beta^*_{b,R}\,{}^b\Omega(X)),$$

of distributions conormal to $\triangle_b$, but C^∞ everywhere else, and vanishing to infinite order at all the faces in X^2_b, except bf (compare [Me0] and Remark 3.1 for an explanation of these notions). The *small* calculus of ϕ-pseudodifferential operators on X is defined as

$$\Psi^m_{\phi,\mathrm{cl}}(X) := \dot{C}^\infty_{\mathrm{ff}} I^m_{\mathrm{cl}}(X^2_\phi, \triangle_\phi; \beta^*_{\phi,R}\,{}^\phi\Omega(X)).$$

Note that we *always* assume our conormal distributions to be classical at the diagonal. The "cl"-notation in the above definitions refers to the fact that the distributions are also assumed to be C^∞ at the faces bf or ff. More generally, given a C^∞-index set $\mathcal{E} = (E_{\mathrm{ff}}, E_{\phi\mathrm{bf}}, E_{\mathrm{rf}}, E_{\mathrm{lf}})$, see Appendix A.6, we can define the corresponding space in the big calculus of ϕ-pseudodifferential operators by

$$\Psi^{m,\mathcal{E}}_\phi(X) := \mathcal{A}^{\mathcal{E}} I^m_{\mathrm{cl}}(X^2_\phi, \triangle_\phi; \beta^*_{\phi,R}\,{}^\phi\Omega(X)),$$

and ditto for b-operators. These definitions also make sense for finite index sets, where the index of lowest order is interpreted as a conormal bound. Note that with this notation

$$\Psi^m_{\phi,\mathrm{cl}}(X) = \Psi^{m,\mathcal{E}}_\phi(X), \quad E_{\mathrm{ff}} = m + \mathbf{N}, \quad E_{\mathrm{lf}} = E_{\mathrm{rf}} = E_{\phi\mathrm{bf}} = \infty,$$

and the similar statement for b-operators is left to the reader.

An advantage of choosing the density bundle to be lifted from the right factor is that for any $\nu \in \Gamma({}^{\phi}\Omega(X))$ and $W \in \Gamma(X, {}^{\phi}TX)$ we have

$$L_{\beta_L^* W} \beta_R^* \nu = 0 \quad \text{on} \quad X^2, \qquad L_{\beta_{\phi,L}^* W} \beta_{\phi,R}^* \nu = 0 \quad \text{on} \quad X_\phi^2.$$

Thus, the lifts to X_ϕ^2 of vector fields in ${}^{\phi}\mathcal{V}$ from the left factor act on elements in $\Psi_\phi^{m,\mathcal{E}}(X)$ (as Lie derivatives) in a particularly simple manner.

The definitions above can be extended to operators between sections of vector bundles in the usual manner. In the case of the Dirac operator on a (ϕ-)Clifford bundle E we introduce the "coefficient bundle"

$$\mathrm{CB}_\phi = \beta_{\phi,L}^* E \otimes \beta_{\phi,R}^* (E^* \otimes {}^{\phi}\Omega(X))$$

on X_ϕ^2 and the analogous definition on X_b^2 to simplify the notation.

2.2 Mapping Properties

In order to describe elementary mapping properties of our pseudodifferential operators we use the pullback and pushforward results for conormal distributions on blown up spaces as described in [Me0] (see also [Lo] or Appendix A.6). First, we need a little

Lemma 2.2 (Densities and Blow up) *Let N be a manifold with corners, and $F \subset \partial Y$ a p-submanifold. Let $S_q \subset N^*F$ be a subspace of q-parabolic directions. Then the blow down map $\gamma : N_F = [N; F, S_q] \longrightarrow N$ has the property*

$$\gamma^* \Omega(N) = \rho_F^{n(F)+(q-1)\dim S_q} \Omega(N_F),$$

where $n(F) := \dim N - 1 - \dim F$ is the codimension of the blow up. ■

Note that the blow ups in this Chapter are all *simple*, i.e. $q = 1$.
Using this, we can describe the action of an element $P \in \Psi_\phi^{m,\mathcal{E}}(X)$ on a function $f \in \mathcal{A}^{\mathcal{F}}(X)$ as follows. Choose a nonvanishing section $u \in C^\infty(X, \Omega(X))$. Lemma 2.2 and the pullback Theorem A.18 then tell us that we have a conormal distribution on X_ϕ^2 of the type

$$\begin{aligned} \beta_{\phi,L}^* u \cdot P \cdot \beta_{\phi,R}^* f \in \mathcal{A}^{\mathcal{G}} I^m(X_\phi^2, \triangle_\phi, \Omega(X_\phi^2)) \qquad & G_{\mathrm{ff}} = E_{\mathrm{ff}} + F, \quad G_{\phi\mathrm{bf}} = E_{\phi\mathrm{bf}} + F - h - 1, \\ & G_{\mathrm{rf}} = E_{\mathrm{rf}} + F - h - 2, \quad G_{\mathrm{lf}} = E_{\mathrm{lf}} \end{aligned}$$

If $G_{\mathrm{rf}} > -1$, we can push forward this expression via $\beta_{\phi,L}$ to the left factor X (i.e. integrate the expression over the right factor) using Theorem A.20 and set

$$Pf = \beta_{\phi,L\,*}(\beta_{\phi,L}^* u \cdot P \cdot \beta_{\phi,R}^* f)/u, \quad \text{i.e} \quad \langle Pf, u \rangle_X = \langle P, \beta_\phi^*(f \boxtimes u) \rangle_{X_\phi^2}, \tag{25}$$

where u is only used for bookkeeping purposes w.r.t. the densities. By the pushforward Theorem A.20 this is a conormal distribution

$$Pf \in \mathcal{A}^G(X), \quad G = E_{\mathrm{lf}} \overline{\cup} (E_{\mathrm{ff}} + F) \overline{\cup} (E_{\phi\mathrm{bf}} + F - h - 1). \tag{26}$$

Analogously, if Q is a b-operator in $\Psi_b^{m,\mathcal{E}}(X)$, with $E_{\mathrm{rf}} + F - 1 > -1$ we can set

$$Qf = \beta_{b,L\,*}(\beta^*_{b,L}u \cdot Q \cdot \beta^*_{b,R}f)/u \in \mathcal{A}^G(X), \quad G = E_{\mathrm{lf}}\overline{\cup}(E_{\mathrm{bf}} + F). \tag{27}$$

As a special case note that elements in the small (b- or ϕ-) calculus always map $\mathcal{A}^F(X)$ to itself.

Having defined the action of ϕ- and b-pseudodifferential operators we can now describe their relationship with the corresponding classes of differential operators. First, note that the identity is an operator in the small calculus of order 0:

$$[\,\mathrm{Id}] \in \Psi^0_{b,\mathrm{cl}}(X), \quad \text{and} \quad [\,\mathrm{Id}] \in \Psi^0_{\phi,\mathrm{cl}}(X). \tag{28}$$

To describe the composition of a ϕ-differential operator, or ϕ-vector field W for simplicity, with a ϕ-pseudodifferential operator P, recall that for forms α_1, α_2 and a b-vector field W on X^2_ϕ the partial integration rule reads

$$\int L_W\alpha_1 \wedge \alpha_2 = -\int \alpha_1 \wedge L_W\alpha_2.$$

Hence, from the definition (25) we get

$$\langle W \circ Pf, u\rangle_X = -\langle Pf, L_W u\rangle = -\langle P, \beta^*_\phi(f \boxtimes L_W u)\rangle_{X^2_\phi} = \langle L_{\beta^*_{\phi,L}W}P, \beta^*_\phi(f \boxtimes u)\rangle_{X^2_\phi},$$

from which we can deduce that $W \circ P = L_{\beta^*_{\phi,L}W}P$. Here we have used the fact (Lemma 2.1) that the vector field $\beta^*_{\phi,L}W$ is tangent to the faces of X^2_ϕ. This argument shows that composition of a ϕ-differential operator and a ϕ-pseudodifferential operator is well defined:

$$\mathrm{Diff}^k_\phi(X) \circ \Psi^{m,\mathcal{E}}_\phi(X) \subset \Psi^{m+k,\mathcal{E}}_\phi(X), \tag{29}$$

for any index set $\mathcal{E}$. Also, from (28) we immediately infer that $\mathrm{Diff}^k_\phi(X) \subset \Psi^k_{\phi,\mathrm{cl}}(X)$.

Sobolev Spaces, Compactness and Trace Properties

We now want to analyze the mapping properties of our operators w.r.t. L^2- and Sobolev spaces. Using the volume form $dvol_\phi$ which is induced by the exact ϕ-metric g^ϕ and $dvol_b = x^{h+1}\, dvol_\phi$, we can introduce the spaces

$$L^2_\phi(X,E) = L^2(X,E;dvol_\phi), \qquad L^2_b(X,E) = L^2(X,E;dvol_b),$$

with norms $\|\cdot\|_\phi$ and $\|\cdot\|_b$. The associated scales of Sobolev spaces are

$$\begin{aligned} H^s_\phi(X,E) &= \{\xi \in C^{-\infty}(X,E) \;\mid\; \Psi^s_\phi(X,E)\xi \subset L^2_\phi(X,E)\} \\ H^s_b(X,E) &= \{\xi \in C^{-\infty}(X,E) \;\mid\; \Psi^s_b(X,E)\xi \subset L^2_b(X,E)\}, \end{aligned}$$

and the Sobolev norms $\|\cdot\|_{\phi,s}$, $\|\cdot\|_{b,s}$ can be defined as usual using fixed families of elliptic operators $P_s \in \Psi^s_{\phi,\mathrm{cl}}(X,E)$, $Q_s \in \Psi^s_{b,\mathrm{cl}}(X,E)$ (invertible, or – in the b-case – invertible modulo an element in $\Psi^{-\infty}_b(X,E)$). The main mapping properties of b- and ϕ-operators w.r.t. these spaces are described in the following

Proposition 2.3 (a) *$Q \in \Psi_b^{m,\mathcal{F}}(X,E)$ with $m \leq 0$ and index sets $F_{\mathrm{bf}} \geq 0$ and $F_{\mathrm{lf}}, F_{\mathrm{rf}} > 0$ is a bounded operator on $L_b^2(X,E)$.*

(b) *$P \in \Psi_\phi^{m,\mathcal{E}}(X,E)$ with $m \leq 0$ and index sets $E_{\mathrm{ff}} \geq 0$, $E_{\phi\mathrm{bf}} \geq h+1$, $E_{\mathrm{lf}} > \frac{h+1}{2}$ and $E_{\mathrm{rf}} > \frac{h+1}{2}$ is a bounded operator on $L_\phi^2(X,E)$.*

(c) *$P \in \Psi_\phi^{m,\mathcal{E}}(X,E)$ with $m \leq 0$ and index sets $E_{\mathrm{ff}} \geq 0$, $E_{\phi\mathrm{bf}} \geq h+1$, $E_{\mathrm{lf}} > 0$ and $E_{\mathrm{rf}} > h+1$ is a bounded operator on $L_b^2(X,E)$.*

Proof. Reduction to the case $m = -\infty$ is done as usual using Hörmander's argument (see [Me0] or Proposition 7 in [MaMe3]). In (a) it then suffices to verify the mapping property for an operator Q supported in a coordinate patch. There, write $Q = a(\mathbf{w},\mathbf{w}')\, dvol_b(\mathbf{w}')$. We can estimate

$$\begin{aligned}
\|Qf\|_b^2 &= \int_X |\int_X a(\mathbf{w},\mathbf{w}')f(\mathbf{w}')\, dvol_b(\mathbf{w}')|^2\, dvol_b(\mathbf{w}) \\
&\leq \int_X \int_X |a(\mathbf{w},\mathbf{w}')|\, dvol_b(\mathbf{w}') \int_X |a(\mathbf{w},\mathbf{w}')||f(\mathbf{w}')|^2\, dvol_b(\mathbf{w}')\, dvol_b(\mathbf{w}).
\end{aligned}$$

Now, it is easy to see that there is an operator $B = b(\mathbf{w},\mathbf{w}')\, dvol_b(\mathbf{w}')$ such that $b(\mathbf{w},\mathbf{w}') \geq |a(\mathbf{w},\mathbf{w}')|$. Then, the above is certainly smaller than

$$\sup_{\mathbf{w}}(\int_X b(\mathbf{w},\mathbf{w}')\, dvol_b(\mathbf{w}')) \cdot \sup_{\mathbf{w}'}(\int_X b(\mathbf{w},\mathbf{w}')\, dvol_b(\mathbf{w})) \cdot \int_X |f(\mathbf{w}')|^2\, dvol_b(\mathbf{w}').$$

The sup's are finite, since B maps the function 1 to $L^\infty(X)$ and the adjoint kernel satisfies the same estimates; (b) and (c) are then consequences of (a) using the fact that $\beta_{\phi,R}^*\Omega_\phi = (x')^{-h-1}\beta_{\phi,R}^*\Omega_b$ and $L_\phi^2(X) = x^{(h+1)/2}L_b^2(X)$. ■

Corollary 2.4 *For $P \in \Psi_{\phi,\mathrm{cl}}^m(X,E)$, $Q \in \Psi_{b,\mathrm{cl}}^m(X,E)$ the maps*

$$P : H_\phi^{s+m}(X,E) \longrightarrow H_\phi^s(X,E), \qquad Q : H_b^{s+m}(X,E) \longrightarrow H_b^s(X,E).$$

are continuous. ■

The standard results on compactness- and trace class-properties of pseudodifferential operators on a compact manifold generalize to our context as follows:

Proposition 2.5 (Sobolev, Rellich and Lidskii)

(a) *$H_b^{n/2+l+\epsilon}(X,E) \hookrightarrow C^l(\overset{\circ}{X},E)$, continuously.*

(b) *$x^\delta H_b^{s+\epsilon}(X,E) \hookrightarrow H_b^s(X,E)$ is compact.*

(c) *Let P be a bounded operator on $L_b^2(X,E)$ and assume $P : H_b^s(X,E) \to x^\delta H_b^{s+n+\epsilon}(X,E)$ continuously. Then P is trace class and $\mathrm{tr}(P) = \int_\triangle [P]$.* ■

The above Sobolev spaces and their variants will only play a minor role in subsequent Chapters of this work, since we will explicitly construct the integral kernels of the resolvent and the heat operator of D^d and not rely on abstract existence results. The main result that we are going to use is the following Corollary, which gives us a minimum requirement for the smallness of the error term in the construction.

Corollary 2.6 (Trace Class Operators)
Operators $Q \in \Psi_b^{-n-\epsilon,(\epsilon,\epsilon,\epsilon)}(X,E)$ *are trace class on* $L_b^2(X,E)$. ■

Finally, let us note how to calculate adjoints w.r.t. $L_b^2(X,E)$: Let Q be an operator in $\Psi_b(X,E)$ or $\Psi_\phi(X,E)$ and write Q^+ for the formal adjoint of Q obtained by flipping sides in X^2. Then

Lemma 2.7 *The* L_b^2*-adjoint of* Q *is given by* $Q^* = Q^+ \cdot dvol_{b,L}^{-1} \otimes dvol_{b,R}$. ■

In order to avoid any confusion due to the variety of different L^2-spaces, and to be able to use the standard constructions of the b-calculus (see [Me3]) without difficulty, we decide to **let all operators act on $\boldsymbol{L_b^2(X)}$**. However, the goal of this work is the description of the spectral properties of the Dirac operator D^d on $L^2(X,E;dvol_d)$, where $dvol_d = x^n\, dvol_\phi = x^v\, dvol_b$ is the volume form corresponding to the metric $g^d = x^2 g^\phi$. Thus, we have to consider the diagram (truly valid only on the domains of definition)

$$\begin{tikzcd}
L^2(X,E;x^n\,dvol_\phi) \arrow[r, "\mathsf{D}^d"] \arrow[d, "x^{v/2}"] & L^2(X,E;x^n\,dvol_\phi) \arrow[d, "x^{v/2}"] \\
L_b^2(X,E) \arrow[r, "\mathsf{D}^d-\frac{v}{2}\mathbf{c}_d(\frac{dx}{x})"] & L_b^2(X,E).
\end{tikzcd}$$

Thus, requiring all our operators to act on $L_b^2(X,E)$ means that we are ultimately looking at the operator

$$\boldsymbol{\mathsf{D} = \mathsf{D}^d - \frac{v}{2}\mathbf{c}_d(\frac{dx}{x}) = \frac{1}{x}\mathsf{D}^\phi + \frac{h}{2}\mathbf{c}_\phi(\frac{dx}{x^2}) \quad \text{on} \quad L_b^2(X,E).}$$

This choice will turn out to be particularly convenient in Sections 3.5 and 3.6.

2.3 Composition Formula

In this Section, we will prove that the composition of two operators in $\Psi_\phi(X)$ is again in $\Psi_\phi(X)$. To explain the main idea of the proof, consider first the "triple space" X^3 together with its projections onto X^2:

$$\begin{tikzcd}
X^2 & X^3 \arrow[l, "\pi_L"'] \arrow[r, "\pi_R"] \arrow[d, "\pi_M"] & X^2 \\
& X^2 &
\end{tikzcd} \tag{30}$$

The composition of two operators $A, B \in \dot{C}^\infty(X^2, \beta_R^*\Omega(X))$ is then simply given by

$$\pi_{M*}(\pi_L^* A \cdot \pi_R^* B).$$



Figure 3: X_b^3

In order to describe the composition properties of ϕ-pseudodifferential operators we need to replace the image spaces in (30) by X_ϕ^2 and modify X^3 in such a way that (30) becomes a diagram of b-fibrations. This will allow us to use the pullback- and pushforward results in [Me0] (see also Appendix A.6) to describe composition of ϕ-pseudodifferential operators.

The first step is to blow up the corner of X^2 in (30), thus replacing it by X_b^2. Blowing up the preimages of the corner under π_L, π_M and π_R at their intersections

$$T := (\partial X)^3, \quad F := X \times \partial X \times \partial X, \quad C := \partial X \times X \times \partial X, \quad S := \partial X \times \partial X \times X,$$

leads to the b-triple space X_b^3 defined as the blow up

$$X_b^3 := \left[X^3; T, F, C, S\right].$$

The projections in (30) induce *b-fibrations* $\pi_{b,o} : X_b^3 \to X_b^2$ for $o = L, M, R$ such that the diagram

$$\begin{tikzcd} X_b^3 \arrow[r, "\pi_{b,o}"] \arrow[d] & X_b^2 \arrow[d] \\ X^3 \arrow[r, "\pi_o"] & X^2 \end{tikzcd} \qquad (31)$$

is commutative. The next step is to replace X_b^2 in (31) by X_ϕ^2 by blowing up the fibre diagonal F_ϕ. To define the new LHS in (31), X_ϕ^3, we have to blow up the lifts of F_ϕ under π_L, π_M and π_R. We denote these faces (as afterwards the result of their blow up) by ϕ_F, ϕ_C, ϕ_S, ϕ_{FT}, ϕ_{CT}, ϕ_{ST}, ϕ_{TT}, and define

$$X_\phi^3 := \left[X_b^3; \phi_{TT}; \phi_{FT}; \phi_{CT}; \phi_{ST}; \phi_F; \phi_C; \phi_S\right].$$



Figure 4: X^3_ϕ

By R, M, C, F, S, C, T we denote the lifts to X^3_ϕ of the non-blown up parts of these faces in X^3_b. We will also have to consider the behavior of our operators at the lifted diagonals $\triangle_{\phi,o} := \pi^{-1}_{\phi,o}\triangle_\phi$, with $o = L, M, R$. The following is proved in [MaMe3]:

Proposition 2.8 *The projections in (30,31) induce b-fibrations $\pi_{\phi,o} : X^3_\phi \to X^2_\phi$ for $o = L, M, R$ such that the diagram*

$$\begin{tikzcd} X^3_\phi \arrow[r, "\pi_{\phi,o}"] \arrow[d] & X^2_\phi \arrow[d] \\ X^3 \arrow[r, "\pi_o"] & X^2 \end{tikzcd}$$

is commutative. ■

The coefficient matrices can be read off from Figure 4:

Lemma 2.9 (a) *π_L maps the face L to the interior of X^2_ϕ. Also*

$$\begin{array}{rclrcl} \pi_L^*\mathrm{lf} & = & \{R, C, \phi_C\} & \pi_L^*\mathrm{rf} & = & \{M, F, \phi_F\} \\ \pi_L^*\phi\mathrm{bf} & = & \{S, T, \phi_{FT}, \phi_{CT}\} & \pi_L^*\mathrm{ff} & = & \{\phi_S, \phi_{ST}, \phi_{TT}\} \end{array}$$

(b) *π_R maps the face R to the interior of X^2_ϕ. Also*

$$\begin{array}{rclrcl} \pi_L^*\mathrm{lf} & = & \{M, S, \phi_S\} & \pi_L^*\mathrm{rf} & = & \{L, C, \phi_C\} \\ \pi_L^*\phi\mathrm{bf} & = & \{F, T, \phi_{ST}, \phi_{CT}\} & \pi_L^*\mathrm{ff} & = & \{\phi_F, \phi_{FT}, \phi_{TT}\} \end{array}$$

(c) *π_M maps the face M to the interior of X^2_ϕ. Also*

$$\begin{array}{rclrcl} \pi_M^*\mathrm{lf} & = & \{R, S, \phi_S\} & \pi_M^*\mathrm{rf} & = & \{L, F, \phi_F\} \\ \pi_M^*\phi\mathrm{bf} & = & \{C, T, \phi_{ST}, \phi_{FT}\} & \pi_M^*\mathrm{ff} & = & \{\phi_C, \phi_{CT}, \phi_{TT}\} \end{array}$$

■

Starting with two operators $A \in \Psi_\phi^{l,\mathcal{L}}(X)$, $B \in \Psi_\phi^{r,\mathcal{R}}(X)$, and a placeholder function $u \in C^\infty(X, \Omega(X))$, their lift to X_ϕ^3 is

$$\gamma_{\phi,L}^* u \cdot \pi_{\phi,L}^* A \cdot \pi_{\phi,R}^* B \in \mathcal{A}^{\mathcal{G}} I^{(l-n/4, r-n/4)}(X_\phi^3, \triangle_{\phi,L} \cup \triangle_{\phi,R}; \Omega(X_\phi^3)). \tag{32}$$

The space on the RHS consists of distributions with conormal singularity at the union $\triangle_{\phi,L} \cup \triangle_{\phi,R}$ of interior manifolds in X_ϕ^3. This is described in [Me0]. At the boundary faces its elements have expansions described by the index set $\mathcal{G}$ as in Appendix A.6. To calculate the index set $\mathcal{G}$ we need to know the lifting properties of the density bundle ${}^\phi\Omega(X)$. The following is proved using Lemma 2.2:

Lemma 2.10 (a) $\gamma_b^* \Omega(X^3) = \rho_{b,F}\, \rho_{b,C}\, \rho_{b,S}\, \rho_{b,T}^2\, \Omega(X_b^3)$

(b) $\gamma_{\phi,b}^* \Omega(X_b^3) = \rho_{\phi T}^{2h+2}\, (\rho_{\phi FT}\, \rho_{\phi CT}\, \rho_{\phi ST}\, \rho_{\phi F}\, \rho_{\phi C}\, \rho_{\phi S})^{h+1} \Omega(X_\phi^3)$

(c) $\gamma_\phi^* \Omega(X^3) = \rho_F\, \rho_C\, \rho_S\, \rho_T^2\, \rho_{\phi F}^{h+2}\, \rho_{\phi C}^{h+2}\, \rho_{\phi S}^{h+2}\, \rho_{\phi FT}^{h+3}\, \rho_{\phi CT}^{h+3}\, \rho_{\phi ST}^{h+3}\, \rho_{\phi TT}^{2h+4} \Omega(X_\phi^3)$

(d) $\gamma_{\phi,F}^* \Omega(X) \cdot \pi_{\phi,L}^* {}^\phi\Omega(X) \cdot \pi_{\phi,R}^* {}^\phi\Omega(X) = \rho_L^0\, \rho_M^{-h-2}\, \rho_R^{-h-2}$

$\cdot \rho_F^{-h-1}\, \rho_C^{-h-1}\, \rho_S^{-2h-3}\, \rho_T^{-2h-2}\, \rho_{\phi F}^0\, \rho_{\phi C}^0\, \rho_{\phi S}^{-h-2}\, \rho_{\phi FT}^{-h-1}\, \rho_{\phi CT}^{-h-1}\, \rho_{\phi ST}^{-h-1}\, \rho_{\phi TT}^0 \Omega(X_\phi^3)$ ■

From this and the pullback Theorem A.18 (or really its extension to distributions conormal at the boundary faces *and* to some interior submanifold, see [Me0]) it follows directly that the index $\mathcal{G}$ is given by

$$\begin{array}{rcl@{\qquad}rcl}
G_L & = & R_{\mathrm{rf}} & G_F & = & L_{\mathrm{rf}} + R_{\mathrm{bf}} - h - 1 \\
G_M & = & L_{\mathrm{rf}} + R_{\mathrm{lf}} - h - 2 & G_C & = & L_{\mathrm{lf}} + R_{\mathrm{rf}} - h - 1 \\
G_R & = & L_{\mathrm{lf}} - h - 2 & G_S & = & L_{\phi\mathrm{bf}} + R_{\mathrm{lf}} - 2h - 3 \\
 & & & G_T & = & L_{\phi\mathrm{bf}} + R_{\phi\mathrm{bf}} - 2h - 2 \\
G_{\phi_F} & = & L_{\mathrm{rf}} + R_{\mathrm{ff}} & G_{\phi_{FT}} & = & L_{\phi\mathrm{bf}} + R_{\mathrm{ff}} - h - 1 \\
G_{\phi_C} & = & L_{\mathrm{lf}} + R_{\mathrm{rf}} & G_{\phi_{CT}} & = & L_{\phi\mathrm{bf}} + R_{\phi\mathrm{bf}} - h - 1 \\
G_{\phi_S} & = & L_{\mathrm{ff}} + R_{\mathrm{lf}} - h - 2 & G_{\phi_{ST}} & = & L_{\mathrm{ff}} + R_{\phi\mathrm{bf}} - h - 1 \\
 & & & G_{\phi_{TT}} & = & L_{\mathrm{ff}} + R_{\mathrm{ff}}.
\end{array}$$

Thus, by the pushforward Theorem A.20 (again, it is really the extension of this result to distributions conormal to $\triangle_{\phi,L} \cup \triangle_{\phi,R}$ in [Me0] that is needed), the conormal distribution in (32) can be pushed forward via $\pi_{\phi,M}$ whenever $G_M = L_{\mathrm{rf}} + R_{\mathrm{lf}} - h - 2 > -1$ giving

$$\pi_{M\,*}(\gamma_{\phi,L}^* u \cdot \pi_{\phi,L}^* A \cdot \pi_{\phi,R}^* B) \in \mathcal{A}^{\pi_{\phi,M\,\sharp}\mathcal{G}} I^{l+r}(X_\phi^2, \triangle_\phi; \Omega(X_\phi^2)).$$

Here the pushforward of the index set $\mathcal{G}$ is given by the rule

$$\pi_{\phi,M\,\sharp}\mathcal{G}(B) = \overline{\bigcup}_{F \in \mathcal{M}(X_\phi^3), \pi_{\phi,M}(F) \subset B} G_F,$$

which by Lemma 2.9 is

$$\begin{array}{rcl@{\qquad}rcl}
\pi_{\phi,M\,\sharp}\mathcal{G}(\mathrm{lf}) & = & G_R \overline{\cup} G_S \overline{\cup} G_{\phi_S} & \pi_{\phi,M\,\sharp}\mathcal{G}(\mathrm{rf}) & = & G_L \overline{\cup} G_F \overline{\cup} G_{\phi_F} \\
\pi_{\phi,M\,\sharp}\mathcal{G}(\phi\mathrm{bf}) & = & G_C \overline{\cup} G_T \overline{\cup} G_{\phi_{ST}} \overline{\cup} G_{\phi_{FT}} & \pi_{\phi,M\,\sharp}\mathcal{G}(\mathrm{ff}) & = & G_{\phi_C} \overline{\cup} G_{\phi_{CT}} \overline{\cup} G_{\phi_{TT}}
\end{array}$$

Rewriting everything in terms of the ϕ-density bundle lifted from the right

$$\beta^*_{\phi,R}{}^{\phi}\Omega(X) = \rho_{\mathrm{lf}}^{-h-2}\,\rho_{\phi\mathrm{bf}}^{-h-1}\Omega(X^2_\phi)/\beta^*_{\phi,L}\Omega(X),$$

we get the final composition formula:

Theorem 2.11 (Composition Formula) *Let* $A \in \Psi_\phi^{l,\mathcal{L}}(X)$ *and* $B \in \Psi_\phi^{r,\mathcal{R}}(X)$ *and assume* $L_{\mathrm{rf}} + R_{\mathrm{lf}} - h - 2 > -1$. *Then* $A \circ B \in \Psi_\phi^{r+l,\mathcal{M}}(X)$ *with index set given by*

$$\begin{array}{lcl}
M_{\mathrm{lf}} & = & L_{\mathrm{lf}}\overline{\cup}(L_{\phi\mathrm{bf}} + R_{\mathrm{lf}} - h - 1)\overline{\cup}(L_{\mathrm{ff}} + R_{\mathrm{lf}}) \\
M_{\mathrm{rf}} & = & R_{\mathrm{rf}}\overline{\cup}(L_{\mathrm{rf}} + R_{\phi\mathrm{bf}} - h - 1)\overline{\cup}(L_{\mathrm{rf}} + R_{\mathrm{ff}}) \\
M_{\phi\mathrm{bf}} & = & (L_{\mathrm{lf}} + R_{\mathrm{rf}})\overline{\cup}(L_{\phi\mathrm{bf}} + R_{\phi\mathrm{bf}} - h - 1)\overline{\cup}(L_{\phi\mathrm{bf}} + R_{\mathrm{ff}})\overline{\cup}(L_{\mathrm{ff}} + R_{\phi\mathrm{bf}}) \\
M_{\mathrm{ff}} & = & (L_{\mathrm{lf}} + R_{\mathrm{rf}})\overline{\cup}(L_{\phi\mathrm{bf}} + R_{\phi\mathrm{bf}} - h - 1)\overline{\cup}(L_{\mathrm{ff}} + R_{\mathrm{ff}})
\end{array}$$

■

All the results above also hold for ϕ-pseudodifferential operators with coefficients in vector bundles.

3 Normal Operators and the Resolvent Construction

In this Chapter we will present a construction of the resolvent $G(\lambda)$ of the Dirac operator D. We do this by first solving the equation

$$(\mathsf{D}-\lambda)Q(\lambda) = I + R(\lambda) \quad \text{on} \quad X^2_\phi \tag{33}$$

with successively smaller and smaller remainder terms $R(\lambda)$, then use the composition formula, Theorem 2.11, to (formally) invert $I + R(\lambda)$ and obtain a smoothing remainder term. In a final step this remainder term is then also removed.

The construction will proceed by the symbolic solution of (33) and the subsequent removal of error terms appearing at the faces ff, ϕbf, lf. The technical tools used are the so-called normal operators, which allow to give a representation of (33) at each of the above faces. These concepts will be introduced at the corresponding stages of the construction in Sections 3.1, 3.3 and 3.5. Prototypes of this kind of construction can be found in [MaMe1] and [Me3].

3.1 Symbol Map

The symbol map ${}^\phi\sigma$ in the small calculus of ϕ-pseudodifferential operators, describes the conormal singularities at $\triangle_\phi$. It is given by the composition

$$\begin{aligned} {}^\phi\sigma_m : \Psi^m_{\phi,\mathrm{cl}}(X) = \dot{C}^\infty_{\mathrm{ff}} I^m_{\mathrm{cl}}(X^2_\phi, \triangle_\phi; \beta^*_{\phi,R}{}^\phi\Omega(X)) &\longrightarrow C^\infty_c I^m_{\mathrm{cl}}(N\triangle_\phi, \triangle_\phi; \Omega(N\triangle_\phi/\triangle_\phi)) \\ &\xrightarrow{F} \mathcal{S}^m_{\mathrm{cl}}(N^*\triangle_\phi) \cong \mathcal{S}^m_{\mathrm{cl}}({}^\phi T^*X). \end{aligned} \tag{34}$$

Here, the first map comes from an identification of $N\triangle_\phi$ with a tubular neighborhood of $\triangle_\phi$, the second map is Fourier transform and the identification at the end is just Lemma 2.1. Also, using radial compactification (for instance by introducing $\rho := |\,|^{-1}$ as a defining function for the boundary at infinity) in the fibres of ${}^\phi T^*X$ we could have written

$$\mathcal{S}^m_{\mathrm{cl}}({}^\phi T^*X) = \rho^{-m} C^\infty(\mathrm{RC}({}^\phi T^*X)),$$

which by multiplication with ρ^m and restriction to the boundary maps to $C^\infty({}^\phi S^*X)$.

Remark 3.1 The existence of the map (34) can be taken to be the defining property of the space $I^m_{\mathrm{cl}}(X^2_\phi, \triangle_\phi; \beta^*_{\phi,R}{}^\phi\Omega(X))$ of distributions conormal at $\triangle_\phi$ of order m and C^∞ up to ff.

Putting everything together, the mth order symbol fits into a short exact sequence

$$0 \longrightarrow \Psi^{m-1}_{\phi,\mathrm{cl}}(X) \longrightarrow \Psi^m_{\phi,\mathrm{cl}}(X) \xrightarrow{{}^\phi\sigma_m} C^\infty({}^\phi S^*X) \longrightarrow 0 \tag{35}$$

of *filtered algebra* homomorphisms.

Note that we will use both versions (34, 35) of the symbol without distinguishing them notationally. Also, the generalization of these definitions to the calculus with coefficients in a vector bundle is left to the reader. Especially, using the identification of $N\triangle_\phi$ with ${}^\phi TX$:

Proposition 3.2 *Let* $W \in \Gamma({}^{\phi}TX)$, $Q \in \Psi_{\phi}^{m}(X)$, $Q_E \in \Psi_{\phi}^{m}(X, E)$.

(a) ${}^{\phi}\sigma_1(W) = iW$, *i.e.* ${}^{\phi}\sigma_{m+1}(W \circ Q) = iW \otimes {}^{\phi}\sigma_m(Q)$

(b) ${}^{\phi}\sigma_1(\mathsf{D}^{\phi}) = {}^{\phi}\sigma_1(x\mathsf{D}^{d}) = {}^{\phi}\sigma_1(x\mathsf{D}) = i\mathbf{c}_{\phi}$, *i.e.* ${}^{\phi}\sigma_{m+1}(\mathsf{D}^{\phi} \circ Q_E) = i\mathbf{c}_{\phi} \otimes {}^{\phi}\sigma_m(Q_E)$,

etc., where W *(and* $\mathbf{c}_{\phi}$*) are interpreted as linear functions on* ${}^{\phi}T^{*}X$. ■

As usual, an element P in $\Psi_{\phi}^{m>0}(X, E)$ will be called *elliptic*, when its symbol ${}^{\phi}\sigma_m(P)$ is invertible in $C^{\infty}({}^{\phi}S^{*}X)$. Thus, for instance, the Dirac operator D^{ϕ} is elliptic.

3.2 Symbolic Construction

To perform the symbolic part of the construction of the inverse of the Dirac operator $\mathsf{D} - \lambda$ write

$$(\mathsf{D} - \lambda)^{-1} = (x\mathsf{D} - x\lambda)^{-1} \cdot x.$$

Thus, we can as well invert the operator $P(\lambda) = x\mathsf{D} - x\lambda$, which is an elliptic operator in the ϕ-calculus:

Proposition 3.3 *There are holomorphic families,*

$$\lambda \mapsto Q_0(\lambda) \in \Psi_{\phi,\mathrm{cl}}^{-1}(X, E), \qquad \lambda \mapsto R_0(\lambda) \in \Psi_{\phi,\mathrm{cl}}^{-\infty}(X, E),$$

such that $P(\lambda)Q_0(\lambda) = 1 + R_0(\lambda)$.

Proof. This is the standard symbolic construction using the symbol sequence

$$0 \longrightarrow \Psi_{\phi}^{m-1}(X, E) \longrightarrow \Psi_{\phi}^{m}(X, E) \xrightarrow{{}^{\phi}\sigma_m} C^{\infty}({}^{\phi}S^{*}X, \mathrm{END}(E)) \longrightarrow 0.$$

The symbol ${}^{\phi}\sigma_1(P(\lambda))$ is independent of λ and invertible. We can therefore find a $g \in C^{\infty}({}^{\phi}S^{*}X, \mathrm{END}(E))$ and, because of the surjectivity of the symbol map ${}^{\phi}\sigma_{-1}$, also a $G \in \Psi_{\phi,\mathrm{cl}}^{-1}(X, E)$ with

$${}^{\phi}\sigma_{-1}(G) = g, \qquad P(\lambda)G = 1 - S(\lambda), \quad \lambda \mapsto S(\lambda) \in \Psi_{\phi,\mathrm{cl}}^{-1}(X, E) \quad \text{holomorphic.}$$

We now set as usual

$$Q_0(\lambda) = G \sum_j S(\lambda)^j, \qquad R_0(\lambda) = P(\lambda)Q_0(\lambda) - 1.$$

The construction can be made holomorphic in the parameter λ. ■

In general the remainder $R_0(\lambda)$ will not vanish to any positive order at the face ff and therefore (compare Section 2.2) cannot be expected to be compact. It will therefore be necessary to "solve away" (at least) the leading term of $R_0(\lambda)$ at ff, i.e. to modify the parametrix $Q_0(\lambda)$ in such a way that the new remainder also vanishes at ff. This will be done in the next two Sections.

3.3 Normal Operator at ff

The normal operator at ff of an operator $P \in \Psi^*_{\phi,\mathrm{cl}}(X)$ is defined as the restriction of the conormal distribution P to that face. To describe this in more detail, first recall that the interior of the front face ff is

$$\overset{\circ}{\mathrm{ff}} = {}^\phi N\partial X \times_Y \partial X \tag{36}$$

and ff fibres over ∂X via the left and right projections

$$\mathrm{ff} \xrightarrow{\beta_{\phi,L}} \partial X \qquad \mathrm{ff} \xrightarrow{\beta_{\phi,R}} \partial X.$$

The density bundle for the ϕ-calculus restricts to ff as the density bundle on the fibres of the left projection

$$\beta^*_{\phi,R}{}^\phi\Omega(X)|_{\mathrm{ff}} \cong \Omega_{\mathrm{fib}}({}^\phi N\partial X \times_Y \partial X \overset{\beta_{\phi,L}}{\to} \partial X) \cong \beta^*_{\phi,R}\Omega({}^\phi N\partial X/Y) \tag{37}$$

This allows us to define the normal map at ff as follows

Definition 3.4 (Suspended Calculus and the Normal Map)

(a) *The space $\Psi^m_{\mathrm{sus}}({}^\phi N\partial X)$ of* suspended *pseudodifferential operators on ${}^\phi N\partial X$ is defined as the space*

$$\dot{C}^\infty I^{m+1/4}_{\mathrm{cl}}(\mathrm{ff}, \triangle_\phi; \Omega_{\mathrm{fib}}(\beta_{\phi,L})) = \mathcal{S}_{\mathrm{fib}} I^{m+1/4}_{\mathrm{cl}}({}^\phi N\partial X \times_Y \partial X, F_\phi; \beta^*_{\phi,R}\Omega({}^\phi N\partial X/Y))$$

acting as ($C^\infty(Y) \times \mathbf{R}^{h+1}$-invariant) convolution operators on (e.g.) $\mathcal{S}_{\mathrm{fib}}({}^\phi N\partial X)$.

(b) *The normal map $N_{\mathrm{ff}} : \Psi^m_{\phi,\mathrm{cl}}(X) \to \Psi^m_{\mathrm{sus}}({}^\phi N\partial X)$ is given by $N_{\mathrm{ff}}(A) := A|_{\mathrm{ff}}$.* ∎

It is again straightforward to extend these definitions to the the calculus with coeffients in a vector bundle as well as to any extended calculus, which allows restriction to ff to be defined. Let us note the main properties of this construction:

Lemma 3.5 (a) *The suspended pseudodifferential operators form a calculus, i.e. composition gives a map*

$$\Psi^m_{\mathrm{sus}}({}^\phi N\partial X) \times \Psi^l_{\mathrm{sus}}({}^\phi N\partial X) \xrightarrow{\circ} \Psi^{l+m}_{\mathrm{sus}}({}^\phi N\partial X).$$

(b) *The normal map fits into a short exact sequence*

$$0 \longrightarrow \rho_{\mathrm{ff}}\Psi^m_{\phi,\mathrm{cl}}(X) \longrightarrow \Psi^m_{\phi,\mathrm{cl}}(X) \xrightarrow{N_{\mathrm{ff}}} \Psi^m_{\mathrm{sus}}({}^\phi N\partial X) \longrightarrow 0,$$

which is a sequence of filtered algebras.

Proof. First, the exactness of the sequence in (b) should be clear. The algebraic properties follow, since composition in $\Psi_{\mathrm{sus}}({}^\phi N\partial X)$ is induced by composition in $\Psi_\phi(X)$: Taking two operators P, Q in the *small* calculus $\Psi^*_{\phi,\mathrm{cl}}(X)$, it follows from the composition formula, Theorem 2.11, that $(P \circ Q)|_{\mathrm{ff}}$ is 0 whenever $P|_{\mathrm{ff}}$ or $Q|_{\mathrm{ff}}$ is 0. Thus, the composition of $P|_{\mathrm{ff}}$ and $Q|_{\mathrm{ff}}$ is independent of the choice of extension to X^2_ϕ. ■

Again, the description of the normal operators for ϕ-differential operators is particularly easy. Since, as we have seen, the composition of a ϕ-differential operator, or ϕ-vector field W, and a ϕ-pseudodifferential operator P is given by

$$W \circ P = L_{\beta^*_{\phi,L} W} P = \beta^*_{\phi,L} W \cdot P, \quad \text{and thus} \quad (W \circ P)|_{\mathrm{ff}} = \beta^*_{\phi,L} W|_{\mathrm{ff}} \cdot P|_{\mathrm{ff}},$$

the normal operator of W can be viewed as the restriction of $\beta^*_{\phi,L} W$ to the front face ff, acting on $\Psi^m_{\mathrm{sus}}({}^\phi N\partial X)$.

This action can be described geometrically using the identification (36). The vector field $\beta^*_{\phi,L} W$ is tangent to the fibres of $\beta_{\phi,R} : \mathrm{ff} \to Y$, i.e. it is tangent to (the left factor of) ${}^\phi N\partial X/Y$ in the interior of ff. Using the identification

$$T^\phi N\partial X/Y \cong \pi^{*\phi} N\partial X \oplus_{\partial X} V\partial X \tag{38}$$

we have

Proposition 3.6 *Define the map* $N_\phi : {}^\phi TX \to T^\phi N\partial X/Y$ *as the composition*

$$N_\phi : {}^\phi TX \longrightarrow {}^\phi TX|_{\partial X} \cong {}^\phi N\partial X \oplus V\partial X \overset{(\pi^*, \mathrm{Id})}{\longrightarrow} T^\phi N\partial X/Y.$$

Then for $Q \in \Psi^m_{\phi,\mathrm{cl}}(X)$, $Q_E \in \Psi^m_{\phi,\mathrm{cl}}(X,E)$ *and* $W \in \Gamma({}^\phi TX)$ *we have*

(a) $N_{\mathrm{ff}}(W \circ Q) = N_\phi(W) \circ N_{\mathrm{ff}}(Q)$.

(b) $N_{\mathrm{ff}}((x\mathsf{D} - x\lambda) \circ Q_E) = N_{\mathrm{ff}}(\mathsf{D}^\phi \circ Q_E) = (N_\phi(\mathbf{c}_\phi) \circ \pi^* \nabla^{E,\phi}) \circ N_{\mathrm{ff}}(Q_E)$. ■

Recall that for two vector fields $A, B \in \Gamma(\partial X, {}^\phi TX)$, the commutator $[\mathsf{n}A, \mathsf{v}B]$ lies in ${}^\phi N\partial X$, but is not necessarily 0. However, the normal operator

$$N_\phi(\mathsf{D}^\phi) = (N_\phi(\mathbf{c}_\phi) \circ \pi^* \nabla^{E,\phi}) = \mathsf{D}^{\phi,V} + \mathsf{n}\mathbf{c}_\phi \, d_{{}^\phi N\partial X}$$

of $P(\lambda)$ has the following special property

Lemma 3.7 $\mathsf{D}^{\phi,V}$ *and* $\mathsf{n}\mathbf{c}_\phi \circ d$ *anticommute over* ∂X.

Proof. First note that it follows from 1.5(c) that for T tangent to the boundary

$$[\nabla^{E,\phi}_T, \mathsf{n}] = 0 \quad \text{and} \quad [\nabla^{E,\phi}_T, \mathsf{v}] = 0.$$

Using this, we can write

$$\begin{aligned} [\mathsf{v}\mathbf{c}_\phi \pi^* \nabla^E, \mathsf{n}\mathbf{c}_\phi \pi^* \nabla^E] &= (\mathsf{v}\mathbf{c}_\phi, \mathsf{n}\mathbf{c}_\phi)[\pi^*\nabla^E, \pi^*\nabla^E] + \mathsf{v}\mathbf{c}_\phi[\pi^*\nabla^E, \mathsf{n}\mathbf{c}_\phi]\pi^*\nabla^E \\ &\qquad - \mathsf{v}\mathbf{c}_\phi[\pi^*\nabla^E, \mathsf{n}\mathbf{c}_\phi]\pi^*\nabla^E + [\mathsf{v}\mathbf{c}_\phi, \mathsf{n}\mathbf{c}_\phi]\pi^*\nabla^E\pi^*\nabla^E \\ &= (\mathsf{v}\mathbf{c}_\phi, \mathsf{n}\mathbf{c}_\phi)\pi^* F^E = 0, \end{aligned}$$

which proves the claim. ■

This Lemma will allow us to calculate solutions for $N_\phi(P(\lambda))$ by separating the horizontal and vertical variables.

3.4 Inversion at ff

In this Section we use the normal operator defined in the previous Section to improve the error term obtained in Proposition 3.3 at the front face ff. This means that we have to solve the equation

$$N_{\mathrm{ff}}(P(\lambda)Q_1(\lambda)) = N_\phi(\mathsf{D}^\phi) \circ N_{\mathrm{ff}}(Q_1(\lambda)) \overset{!}{=} -N_{\mathrm{ff}}(R_0(\lambda)) \in \Psi^\infty_{\mathrm{sus}}({}^\phi N\partial X, E),$$

i.e. we have to invert the operator $N_\phi(\mathsf{D}^\phi) = (\mathsf{D}^{\phi,V} + \mathsf{nc}_\phi\, d_{{}^\phi N\partial X})$ over ${}^\phi N\partial X/Y$.

Clearly, sections in the null space of $\mathsf{D}^{\phi,V}$ will be of special importance in this context. At each point $y \in Y$ the null space of the operator $\mathsf{D}^{\phi,V}|_{Z_y}$ is given by

$$\mathcal{K}_y = \mathrm{null}(\mathsf{D}^{\phi,V})|_{Z_y} \subset L^2(Z_y, dvol_Z).$$

From now on we *assume* that

$$\boldsymbol{\mathcal{K} = (\mathrm{null}(\mathsf{D}^{\phi,V})|_{Z_y})_{y\in Y} \to Y \quad \text{is a vector bundle over } Y}, \tag{39}$$

i.e. $\mathrm{null}(\mathsf{D}^{\phi,V})$ is the space of sections of the vector bundle $\mathcal{K} \to Y$. Denote by $\Pi_\circ, \Pi_\perp$ the projections onto this null space and its orthogonal complement. At the boundary this gives us the decomposition into “zero modes” and “nonzero modes”

$$C^\infty(\partial X, E) \cong \Pi_\circ C^\infty(\partial X, E) \oplus \Pi_\perp C^\infty(\partial X, E).$$

Such decompositons also exist at the faces lf, ϕbf, ff and it follows from Lemma 3.7 that $N_\phi(\mathsf{D}^\phi)$ preserves this decomposition over ff. As explained in Appendix A.6 it will be useful to consider spaces of polyhomogeneous, conormal sections whose top (in the sense defined there) coefficients lie in either part of the decomposition. A more refined index notation adapted to this situation is also described in Appendix A.6.

The Dirac Operator on $\mathbf{R}^n$

We start with the analysis of our model operator restricted to $\Pi_\circ$, i.e. the (constant coefficient) Dirac operator on $\mathbf{R}^n$: Denote by g_n the euclidean metric on $\mathbf{R}^n$ and by $dvol_n$ the euclidean volume form. Let $\mathcal{W}$ be a Clifford module over $\mathbf{R}^n$ with Clifford action $\mathbf{c}_n \in \Gamma(T\mathbf{R}^n, \mathrm{End}(\mathcal{W}))$ and trivial connection d. The Dirac operator is then $\mathsf{D}_n = \mathbf{c} \circ d$.

Writing R for the tautological (or radial) vector field in $T\mathbf{R}^n$ the fundamental solution for D_n is

$$E_n = \mathrm{C}_n |\cdot|^{-n}\mathbf{c}_n(\mathsf{R}), \quad \text{or} \quad E_n(\mathbf{y}) = \mathrm{C}_n \sum_{j=1}^n r^{-n} y^j \mathbf{c}(\,dy^j),$$

when one prefers the use of the standard euclidean coordinates $\mathbf{y} \overset{\wedge}{=} (y^1, \dots, y^n)$, $r = |\mathbf{y}|$, on $\mathbf{R}^n$. Then

$$\mathsf{D}_n E_n = \delta \quad \text{in} \quad C^{-\infty}(\mathbf{R}^n, \mathrm{End}(\mathcal{W})).$$

Our idea is to analyze the behavior of solutions for D_n, by first looking at solutions for the conformally transformed Dirac operator D_b on the cylinder $\mathbf{R}_+ \times S^{n-1}$. For this operator we can the use the known methods of the b-calculus from [Me3].

Thus identify $\mathbf{R}^n \setminus 0$ with $\mathbf{R}_+ \times S^{n-1}$ and introduce the b-metric $g_b = r^{-2}g_n$. The corresponding b-volume form is $dvol_b = r^{-n}\, dvol_n$. It then follows again as in Appendix A.2 that $\mathcal{W}$ is a Clifford module over $\mathbf{R}_+ \times S^{n-1}$, with Clifford action $\mathbf{c}_b = r\mathbf{c}_n$ and Dirac operator

$$\mathsf{D}_b = r^{(n-1)/2} r\mathsf{D}_n r^{-(n-1)/2} = r\mathsf{D}_n - \frac{n-1}{2}\mathbf{c}_n(\, dr). \tag{40}$$

This operator is b-elliptic and selfadjoint on $L^2_b(\mathbf{R}_+ \times S^{n-1})$ and $r\mathsf{D}_n$ is a b-elliptic, b-differential operator on $\mathbf{R}_+ \times S^{n-1}$. We have

$$L^2(\mathbf{R}^n, \mathcal{W}; dvol_n) \cong r^{-n/2} L^2(\mathbf{R}_+ \times S^{n-1}, \mathcal{W}; dvol_n) \equiv r^{-n/2} L^2_b(\mathbf{R}_+ \times S^{n-1}, \mathcal{W}).$$

The main result is now

Lemma 3.8 *The operator* $r\mathsf{D}_{n,\alpha} : r^\alpha H^{k+1}_b(\mathbf{R}_+ \times S^{n-1}, \mathcal{W}) \to r^\alpha H^k_b(\mathbf{R}_+ \times S^{n-1}, \mathcal{W})$ *is invertible if* $\alpha \notin \mathbf{Z}$. *More exactly the extended set of indicial roots is* $i\mathbf{N} \cup -i((n-1)-\mathbf{N})$, *especially the indicial roots are all simple.*

Proof. First, we know from the general b-calculus that the operator $\mathsf{D}_{b,\alpha}$ is invertible except if $i\alpha$ is an indicial root, where it has a null space in the "extended L^2-sense". Since $\mathsf{D}_b|_{r=0}$ anticommutes with $\mathbf{c}_b(\frac{dr}{r})$, one easily finds that the set of indicial roots $\operatorname{spec}_I(\mathsf{D}_b)$ is invariant under multiplication with -1. Also it follows immediately from (40) that $\operatorname{spec}_I(r\mathsf{D}_n) = -i\frac{n-1}{2} + \operatorname{spec}_I(\mathsf{D}_b)$. It therefore suffices to show that

$$\{z \in \operatorname{spec}_I(r\mathsf{D}_n) \mid \operatorname{Im}(z) \geq -(n-1)\} \overset{!}{=} -i(n-1) \cup i\mathbf{N}.$$

This is easy. Let $u_\alpha \in \operatorname{null}_-(r\mathsf{D}_{n,\alpha}) \subset r^\alpha L^\infty(\mathbf{R}_+ \times S^{n-1})$. If $\alpha > -n$ then the distribution u_α becomes integrable in 0 w.r.t. $dvol_n$ and extends to a distribution (also denoted u_α) on $\mathbf{R}^n$. This distribution fulfills

$$\mathsf{D}_n u_\alpha = \delta w, \quad \text{i.e.} \quad \mathsf{D}_n(u_\alpha - E_n(w)) = 0$$

in all of $\mathbf{R}^n$ for some $w \in \mathcal{W}$. By elliptic regularity, we can conclude that $u_\alpha - E_n(w)$ is in $C^\infty(\mathbf{R}^n, \mathcal{W})$, from which the claim follows. ■

This result can be used to solve $\mathsf{D}_n u = f$ for f with compact support in $\mathbf{R}^n$. First, if $f \in C^\infty_c(\mathbf{R}_+ \times S^{n-1}, \mathcal{W})$ we also have $rf \in C^\infty_c(\mathbf{R}_+ \times S^{n-1}, \mathcal{W})$. Choosing $\alpha \notin \mathbf{Z}$ set

$$u_\alpha = (r\mathsf{D}_{n,\alpha})^{-1} rf \in r^\alpha H^\infty_b(\mathbf{R}_+ \times S^{n-1}, \mathcal{W}).$$

Then the usual results for elliptic b-pseudodifferential operators imply that u_α has expansions at $r = 0$ and $1/r = 0$ of the form

$$\begin{aligned} u_\alpha &\sim \sum_{\mathbf{Z} \ni j > \alpha} a_j(\omega) r^j \quad \text{for} \quad r \to 0 \\ u_\alpha &\sim \sum_{\mathbf{Z} \ni j > -\alpha} b_j(\omega) \rho^j \quad \text{for} \quad \rho = r^{-1} \to 0. \end{aligned}$$

If we choose $\alpha > -n$, the distribution u_α again extends to a distribution on $\mathbf{R}^n$. Writing $\mathbf{B}^n := \mathrm{RC}(\mathbf{R}^n)$ for the radial compactification of $\mathbf{R}^n$, we have shown the more precise statement

$$u_\alpha - E_n(w) \in \rho^{n-1} C^\infty(\mathbf{B}^n, \mathcal{W}).$$

Of course, due to the translational invariance of the problem, this argument works for any $f \in C_c^\infty(\mathbf{R}^n, \mathcal{W})$ by first shifting the support of f away from 0. We now show

Lemma 3.9 (a) $\mathsf{D}_n : \rho^\alpha H_b^\infty(\mathbf{B}^n, \mathcal{W}) \to \rho^{\alpha+1} H_b^\infty(\mathbf{B}^n, \mathcal{W})$.

(b) *For any* $f \in \mathcal{A}^F(\mathbf{B}^n, \mathcal{W})$ *there exists a*

$$u \in \mathcal{A}^U(\mathbf{B}^n, \mathcal{W}), \qquad U = (n - 1 + \mathbf{N}) \cup ((\mathbf{Z}_{\geq F} \overline{\cup} F) - 1),$$

which solves $\mathsf{D}_n u = f$.

Proof. Choose a cutoff function $\varphi \in C_c^\infty(\mathbf{B}^n)$. Then, the compactly supported section φf, as treated above, is responsible for the term $n - 1 + \mathbf{N}$ in the index set U.

Choosing a noninteger $\alpha < F$ we know that $(1 - \varphi) f \in \rho^\alpha H_b^\infty(\mathbf{R}_+ \times S^{n-1}, \mathcal{W})$, and by Lemma 3.8 there is $v_\alpha \in \rho^{\alpha-1} H_b^\infty(\mathbf{R}_+ \times S^{n-1}, \mathcal{W})$ satisfying $\mathsf{D}_n v_\alpha = (1 - \varphi) f$. By the general theory of b-pseudodifferential operators (compare [Me3], Chapter 5)

$$v_\alpha \in \mathcal{A}^G(\mathbf{R}_+ \times S^{n-1}, \mathcal{W}), \quad G(r = 0) = \{1 - \alpha < j \in \mathbf{Z}\}, \quad G(\rho = 0) = (F \overline{\cup} \{\alpha < j \in \mathbf{Z}\}) - 1.$$

Now, choosing another cutoff function $\eta \in C_c^\infty(\mathbf{B}^n)$ with $(1 - \eta)(1 - \varphi) = 1 - \varphi$, we can write

$$\mathsf{D}_n (1 - \eta) v_\alpha = (1 - \eta) \mathsf{D}_n v_\alpha - \mathbf{c}(\, d\eta) v_\alpha = (1 - \varphi) f - \mathbf{c}(\, d\eta) v_\alpha.$$

But $\mathbf{c}(\, d\eta)$ is compactly supported and can therefore be treated as in the first part. ■

We have solved the model problem in the zero-modes at ff. The model problem in the nonzero-modes is easy. Noting that

$$\Psi_{\text{sus}}^{-\infty, F}({}^\phi N \partial X, E) = \mathcal{A}^F(\mathrm{RC}({}^\phi N \partial X \times_Y \partial X), \mathrm{CB}_\phi) = \mathcal{A}^F(\mathrm{ff}, \mathrm{CB}_\phi)$$

we get

Lemma 3.10 (Solution of the Model Problem at ff)

(a) *For every* $f_\circ \in \Pi_\circ \Psi_{\text{sus}}^{-\infty, F}({}^\phi N \partial X, E)$ *there exists*

$$u_\circ \in \Pi_\circ \Psi_{\text{sus}}^{-\infty, U}({}^\phi N \partial X, E), \qquad U = (h + \mathbf{N}) \cup ((F \overline{\cup} \mathbf{Z}_{\geq F}) - 1),$$

which solves $N_{\mathrm{ff}}(\mathsf{D}^\phi) u_\circ = f_\circ$.

(b) *For every* $f_\perp \in \Pi_\perp \Psi_{\text{sus}}^{-\infty, F}({}^\phi N \partial X, E)$ *there exists an* $u_\perp \in \Pi_\perp \Psi_{\text{sus}}^{-\infty, F-1}({}^\phi N \partial X, E)$ *which solves* $N_{\mathrm{ff}}(\mathsf{D}^\phi) u_\perp = f_\perp$.

Proof. First, note that $N_{\mathrm{ff}}(\mathsf{D}^\phi)$ restricted to $\Pi_\circ H_b^\infty(\mathrm{ff}, \mathrm{CB}_\phi)$ is just the family of (constant coefficient) fibre Dirac operators given by $\mathsf{nc}_\phi \circ d$ on ${}^\phi N \partial X \times_Y \partial X / F_\phi$. Hence (a) follows by applying Lemma 3.9 fibre by fibre. ■

Using Lemma 3.10 and the fact that $N_{\mathrm{ff}}(P(\lambda)) = N_{\mathrm{ff}}(\mathsf{D}^\phi)$ it is now straightforward to prove the following refinement of Proposition 3.3

Proposition 3.11 *There are holomorphic families*

$$\lambda \mapsto Q_1^N(\lambda) \in \Psi_\phi^{-1,\mathcal{E}}(X,E), \quad \textit{and} \quad \lambda \mapsto R_1^N(\lambda) \in \Psi_\phi^{-\infty,\mathcal{F}}(X,E),$$

such that such that

$$P(\lambda)Q_1^N(\lambda) = 1 + R_1^N(\lambda),$$

and with index sets $\mathcal{E}$, $\mathcal{F}$ given by

$$F_{\phi\mathrm{bf}} = (h+1+\mathbf{N})^{\overline{\cup}N}, \quad F_{\mathrm{ff}} = N + \mathbf{N}$$

$$E_{\phi\mathrm{bf}} = (((h,0),\dots,(h,N-1))^\circ, (h+1+\mathbf{N})^{\overline{\cup}N}) = [\circ](h+\mathbf{N})^{\overline{\cup}N}, \quad E_{\mathrm{ff}} = \mathbf{N},$$

(all other index sets are ∞).

Note that the special case $N = 1$ will suffice for the construction of the resolvent!

Proof. For purposes of notational simplicity we are going to ignore coefficients. From the symbolic construction, we have holomorphic families

$$\lambda \mapsto Q_0(\lambda) \in \Psi_{\phi,\mathrm{cl}}^{-1}(X), \qquad \lambda \mapsto R_0(\lambda) \in \Psi_{\phi,\mathrm{cl}}^{-\infty}(X), \quad \text{with} \quad P(\lambda)Q_0(\lambda) = 1 + R_0(\lambda).$$

Setting $S(\lambda) = -R_0(\lambda)$, we want to solve

$$P(\lambda)G(\lambda) = S(\lambda)$$

holomorphically in λ to order N at ff. For this, we write $G(\lambda)$ and $S(\lambda)$ in a (finite) Taylor series (sum):

$$G(\lambda) = G_0(\lambda) + x'G_1(\lambda) + (x')^2G_2(\lambda) + \ldots + (x')^{N-1}G_{N-1}(\lambda),$$

$$S(\lambda) = S_0(\lambda) + x'S_1(\lambda) + (x')^2S_2(\lambda) + \ldots + (x')^{N-1}S_{N-1}(\lambda) + (x')^NS_N(\lambda),$$

and the maps

$$\lambda \mapsto G_j(\lambda) \in \Psi_\phi^{-\infty,\mathcal{I}_j}(X) \quad \text{and} \quad \lambda \mapsto S_j(\lambda) \in \Psi_\phi^{-\infty,\mathcal{J}_j}(X)$$

are holomorphic. As part of the proof we will show that

$$I_{j,\mathrm{ff}} = 0, \quad I_{j,\phi\mathrm{bf}} = [\circ](h-j+\mathbf{N})^{\overline{\cup}j+1}, \quad J_{j,\mathrm{ff}} = 0, \quad J_{j,\phi\mathrm{bf}} = (h-j+1+\mathbf{N})^{\overline{\cup}j}.$$

In the first step, we have to solve

$$N_{\mathrm{ff}}(S_0(\lambda)) = N_{\mathrm{ff}}(P(\lambda)G_0(\lambda)) = N_{\mathrm{ff}}(\mathsf{D}^\phi)N_{\mathrm{ff}}(G_0(\lambda)).$$

Write $s_0(\lambda)$ for $N_{\mathrm{ff}}(S_0(\lambda))$. It is a holomorphic family $\lambda \mapsto s_0(\lambda) \in \dot{C}^\infty(\mathrm{ff})$. By Lemma 3.10, we can find a holomorphic family

$$\lambda \mapsto g_0(\lambda) = g_0(\lambda)^\perp + g_0(\lambda)^\circ \in \Pi_\perp \dot{C}^\infty(\mathrm{ff}) + \Pi_\circ\mathcal{A}^{h+\mathbf{N}}(\mathrm{ff}), \quad N_{\mathrm{ff}}(\mathsf{D}^\phi)g_0(\lambda) = s_0(\lambda),$$

which can be extended to a holomorphic family

$$\lambda \mapsto G_0(\lambda) \in \Psi^{-\infty,\mathcal{I}_0}, \quad I_{0,\mathrm{bf}} = ((h,0)^\circ, h+1+\mathbf{N}), \quad N_{\mathrm{ff}}(\mathsf{D}^\phi G_0(\lambda) - S_0(\lambda)) = 0.$$

Thus, we have shown the case $j = 0$.

Assuming inductively that the equation has been solved up to order j, with index sets as indicated, we set

$$S_{j+1}(\lambda) = (x')^{-1}(P(\lambda)G_j(\lambda) - S_j(\lambda)).$$

Since $P(\lambda)G_j(\lambda)$ has index set $(h+1-j+\mathbf{N})^{\overline{\cup}j+1}$ and $S_j(\lambda)$ has index set $(h-j+1+\mathbf{N})^{\overline{\cup}j}$ at bf, the index set for $S_{j+1}(\lambda)$ is just $\mathcal{J}_j$. Now, the equation

$$N_{\mathrm{ff}}(\mathsf{D}^{\phi}G_{j+1}(\lambda) - S_{j+1}(\lambda)) = 0$$

can be solved holomorphically in λ by Lemma 3.10 and G_{j+1} has index set $\mathcal{I}_{j+1}$. Setting $Q_1^N(\lambda) = Q_0(\lambda) + G(\lambda)$ and $R_1^N(\lambda) = (x')^N S_N(\lambda)$ yields the result. ■

For reference in the next Sections we note the case $N = 1$ as a Corollary. Set $Q_1(\lambda) = x'Q_1^1(\lambda)$ and $R_1(\lambda) = \frac{x'}{x}R_1^1(\lambda)$:

Corollary 3.12 *$D_\lambda Q_1(\lambda) = 1 + R_1(\lambda)$ with holomorphic families*

$$\lambda \mapsto R_1(\lambda) \in \Psi_{\phi}^{\infty,\mathcal{F}}(X,E), \qquad F_{\phi\mathrm{bf}} = (h+1+\mathbf{N}), \qquad F_{\mathrm{ff}} = 1 + \mathbf{N}$$

$$\lambda \mapsto Q_1(\lambda) \in \Psi_{\phi}^{-1,\mathcal{E}}(X,E), \qquad E_{\phi\mathrm{bf}} = [\circ](h+1+\mathbf{N}), \quad E_{\mathrm{ff}} = 1 + \mathbf{N},$$

the other terms in the index sets are ∞. ■

Thus, we have solved the resolvent equation (33) up to a remainder term, which is C^∞ in the interior and which vanishes to first order at the front face. Unfortunately, by Section 2.2, this new remainder is still not compact, since it does not vanish to positive order at ϕbf (w.r.t. to the coefficients CB_b, which are the relevant ones at that face). Thus the next two Sections will describe how to improve the remainder at ϕbf.

3.5 Reduced Normal Operators at bf and ϕbf

In this Section we define the normal operator at the face ϕbf and describe the model action of the Dirac operator D at this face. Recall the relation between the b- and the ϕ-coefficient bundles: $\mathrm{CB}_b = (x')^{h+1}\mathrm{CB}_\phi$. The normal operators at ϕbf and bf are defined by

Definition 3.13 *Let $\mathcal{E}$, $\mathcal{F}$ be index sets for X_ϕ^2 and X_b^2 respectively, with $E_{\phi\mathrm{bf}} = h+1+\mathbf{N}$ and $F_{\mathrm{bf}} = \mathbf{N}$.*

(a) $N_{\mathrm{bf}} : \Psi_b^{-\infty,\mathcal{F}}(X,E) \longrightarrow \mathcal{A}^{(F_{\mathrm{lf}},F_{\mathrm{rf}})}(\mathrm{bf},\mathrm{CB}_b) \qquad A \longmapsto A|_{\mathrm{bf},\mathrm{CB}_b}$

(b) $N_{\phi\mathrm{bf}} : \Psi_\phi^{-\infty,\mathcal{E}}(X,E) \longrightarrow \mathcal{A}^{(E_{\mathrm{lf}},E_{\mathrm{rf}},E_{\mathrm{ff}})}(\phi\mathrm{bf},\mathrm{CB}_b) \qquad A \longmapsto A|_{\phi\mathrm{bf},\mathrm{CB}_b}$ ■

The corresponding exact sequences are

$$0 \longrightarrow \rho_{\mathrm{bf}}\Psi_b^{-\infty,\mathcal{F}}(X,E) \longrightarrow \Psi_b^{-\infty,\mathcal{F}}(X,E) \xrightarrow{N_{\mathrm{bf}}} \mathcal{A}^{(F_{\mathrm{lf}},F_{\mathrm{rf}})}(\mathrm{bf},\mathrm{CB}_b) \longrightarrow 0$$

$$0 \longrightarrow \rho_{\phi\mathrm{bf}}\Psi_\phi^{-\infty,\mathcal{E}}(X,E) \longrightarrow \Psi_\phi^{-\infty,\mathcal{E}}(X,E) \xrightarrow{N_{\phi\mathrm{bf}}} \mathcal{A}^{(E_{\mathrm{lf}},E_{\mathrm{rf}},E_{\mathrm{ff}})}(\phi\mathrm{bf},\mathrm{CB}_b) \longrightarrow 0.$$

Note also, that whenever $E_{\mathrm{ff}} = \infty$ we have

$$\Psi_\phi^{-\infty,\mathcal{E}}(X,E) \subset \Psi_b^{-\infty,(E_{\mathrm{rf}}-h-1,\mathbf{N},E_{\mathrm{lf}})}(X,E),$$

and the normal map $N_{\phi\mathrm{bf}}$, restricted to the LHS, then equals N_{bf}. We will not try to describe the general behavior of the above sequences under composition of operators. Instead we will restrict ourselves to the analysis of the normal action of the Dirac operator D, which will be sufficient for our purposes.

Recall that the Dirac operator D does *not* map calculi of the above type to themselves, since its lift to X_b^2 or X_ϕ^2 is not tangent to bf or ϕbf. From Appendix A.6 we know that

$$\mathsf{D}_\lambda : \Psi_b^{-\infty,(E_{\mathrm{rf}},[\circ]E_{\mathrm{lf}},[\circ]\mathbf{N}_{\mathrm{bf}})}(X,E) \longrightarrow \Psi_b^{-\infty,(E_{\mathrm{rf}},E_{\mathrm{lf}},\mathbf{N}_{\mathrm{bf}})}(X,E)$$

$$\mathsf{D}_\lambda : \Psi_\phi^{-\infty,(E_{\mathrm{rf}},[\circ]E_{\mathrm{lf}},[\circ]\mathbf{N}_{\phi\mathrm{bf}},E_{\mathrm{ff}})}(X,E) \longrightarrow \Psi_\phi^{-\infty,(E_{\mathrm{rf}},E_{\mathrm{lf}},\mathbf{N}_{\phi\mathrm{bf}},E_{\mathrm{ff}}-1)}(X,E),$$

i.e. D_λ maps nicely, when we restrict to conormal distributions with top cofficients in the zero modes at lf and ϕbf. This allows to define the normal action $N_{\mathrm{bf}}(\mathsf{D}_\lambda)$ on the sections of $\mathcal{K}$ over bf (resp. $N_{\phi\mathrm{bf}}(\mathsf{D}_\lambda)$ on ϕbf), defined as usual by

$$N_{\mathrm{bf}}(\mathsf{D}_\lambda)N_{\mathrm{bf}}(A) = \Pi_\circ N_{\mathrm{bf}}(\mathsf{D}_\lambda A), \quad \text{for} \quad A \in \Psi_b^{-\infty}(X,E), \quad N_{\mathrm{bf}}(A) \in \mathcal{K}.$$

The Model Operator at bf

Let us describe the action of $N_{\mathrm{bf}}(\mathsf{D}_\lambda)$ at bf. First, identify bf $-$ rf with the fibre bundle

$$\beta_{b,L} : \partial X \times \partial X \times [0,\infty[_s \to \partial X, \quad s = x/x', \qquad \beta_{b,R}^* {}^b\Omega(X) = {}^b\Omega_{\mathrm{fib}}(\beta_{b,L}).$$

Thus, writing $B = Y \times [0,\infty[$, the space of sections in the null space of $\mathsf{D}^{\phi,V}$ over bf is

$$\Pi_\circ C^\infty(\mathrm{bf},\mathrm{CB}_b) = C^\infty(\partial X, C^\infty(B, \mathcal{K} \otimes {}^b\Omega(B))),$$

and $N_{\mathrm{bf}}(\mathsf{D}_\lambda)$ acts as the operator on

$$C^\infty(B,\mathcal{K} \otimes {}^b\Omega(B)) = \Pi_\circ C^\infty(\partial X \times \mathbf{R}_+, E_{\partial X}), \tag{41}$$

constant in the parameter space ∂X, given by fixing coefficients at the boundary

$$I_b(\mathsf{D}_\lambda) := \Pi_\circ \mathsf{D}_\lambda|_{\partial X,{}^bTX} = [\Pi_\circ \frac{1}{x}\mathsf{D}^\phi + \frac{h}{2}\mathbf{c}_\phi(\frac{dx}{x^2}) - \lambda]|_{\partial X,{}^bTX}. \tag{42}$$

This definition is *much less innocent* than it looks at first sight. The expression actually may involve hidden endomorphisms originating from vertical differentiations compressed to $\mathcal{K}$. We therefore prefer another way to define this "indicial operator":

Lemma 3.14 (a) *Let $\xi \in C^\infty(X,E)^\circ$, i.e. $\mathsf{D}_\lambda\xi \in C^\infty(X,E)$. Then*

$$I_b(\mathsf{D}_\lambda)(\xi|_{\partial X}) := \Pi_\circ(\mathsf{D}_\lambda\xi)|_{\partial X}$$

is well defined and only depends on $\xi|_{\partial X} \in \Pi_\circ C^\infty(\partial X,E)$.

(b) *The family*

$$z \mapsto I_b(\mathsf{D}_\lambda)(z)(\xi|_{\partial X}) := \Pi_\circ(x^{-iz}\mathsf{D}_\lambda x^{iz}\xi|_{\partial X}) \in \operatorname{Diff}^1(Y, \mathcal{K}),$$

is well defined (i.e. independent of the choice of extension ξ of $\xi|_{\partial X}$) and holomorphic in z.

(c) $I_b(\mathsf{D}_\lambda)(z) = M_{x\to z} I_b(\mathsf{D}_\lambda) M_{z\to x}$, *where M denotes the Mellin transform (see (48)).*

(d) *Let $\xi \in C^\infty(X, E)$ such that $\mathsf{D}_\lambda^l \xi \in C^\infty(X, E)$. Then*

$$\Pi_\circ(\mathsf{D}_\lambda^l\xi)|_{\partial X} = [I_b(\mathsf{D}_\lambda)]^l(\xi|_{\partial X}).$$

Moreover any $\xi|_{\partial X}$ can be extended to $\widetilde{\xi}$ such that $(\mathsf{D}_\lambda^l\widetilde{\xi})|_{\partial X} = [I_b(\mathsf{D}_\lambda)]^l(\xi|_{\partial X})$.

Proof. In (a) and (b), only the independence of the extension of $\xi|_{\partial X}$ might be worth a little thought. It follows from the fact, that for $\eta \in C^\infty(X, E)$ we have

$$\Pi_\circ(\mathsf{D}_\lambda x\eta)|_{\partial X} = \Pi_\circ(x\mathsf{D}_\lambda\eta + x\mathbf{c}_d(\frac{dx}{x})\eta)|_{\partial X} = \Pi_\circ(x\mathsf{D}\eta)|_{\partial X} = 0.$$

Thus, the expression defining $I_b(\mathsf{D}_\lambda)(\xi|_{\partial X})$ vanishes for the difference of two extensions of $\xi|_{\partial X}$.

Part (d) is proved by induction. We have proved the case $l = 1$ in (a), now assume that

$$\Pi_\circ\mathsf{D}_\lambda^{l-1}\xi|_{\partial X} = [I_b(\mathsf{D}_\lambda)]^{l-1}(\xi|_{\partial X}).$$

Then starting with ξ such that $\mathsf{D}_\lambda^l\xi \in C^\infty(X, E)$ means that we have $\mathsf{D}_\lambda^{l-1}\xi \in C^\infty(X, E)^\circ$. Therefore,

$$\Pi_\circ\mathsf{D}_\lambda^l\xi|_{\partial X} = \Pi_\circ\mathsf{D}(\mathsf{D}_\lambda^{l-1}\xi)|_{\partial X} = I_b(\mathsf{D}_\lambda)[I_b(\mathsf{D}_\lambda)]^{l-1}(\xi|_{\partial X}),$$

by (a) and the hypothesis of the induction. ■

Geometrical Description of the Indicial Operator

Let us take the time to describe the geometry underlying our definition of the indicial family $I_b(\mathsf{D})(z)$. This will not be strictly necessary for the analysis of the model problem at ϕbf and bf, but gives some insight in the specialties of "ϕ-geometry". Recall that bwe write $\nu = \frac{1}{x}\mathsf{X}_d$. Then, starting with $\xi|_{\partial X} \in \Gamma(Y, \mathcal{K})$ and an extension$\xi \in C^\infty(X, E)^\circ$, we can use Lemma 3.14 to get at the boundary

$$\begin{aligned} I_b(\mathsf{D})(iz)(\xi|_{\partial X}) &= \Pi_\circ(x^z\mathsf{D}x^{-z}\xi|_{\partial X}) = -(z + v/2)\Pi_\circ\mathbf{c}_d(\frac{dx}{x})\xi + \Pi_\circ\mathsf{D}^d\xi \\ &= -(z + v/2)\Pi_\circ\mathbf{c}_d(\frac{dx}{x})\xi + \Pi_\circ\nabla_\nu^{E,d}x\mathsf{D}^d\xi \\ &= -(z + v/2)\Pi_\circ\mathbf{c}_d(\frac{dx}{x})\xi + \Pi_\circ\left([\nabla_\nu^{E,d}, \mathbf{c}_d \circ x\nabla_{\cdot}^{E,d}]\xi + x\mathsf{D}^d\nabla_\nu^{E,d}\xi\right) \\ &= -(z + v/2)\Pi_\circ\mathbf{c}_d(\frac{dx}{x})\xi + \Pi_\circ\mathbf{c}_d(xF^{E,d}(\nu, \cdot))\xi - \Pi_\circ\mathbf{c}_d\nabla_{x\nabla_{\cdot}^{\phi}\nu}^{E,d}\xi. \qquad (43) \end{aligned}$$

In the last step we have used that $\Pi_\circ x\mathsf{D}^d\nabla^{E,d}_\nu\xi|_{\partial X} = 0$, the definition of the curvature $F^{E,d}$ and the fact that $A - x\nabla^d_A\nu = -x\nabla^\phi_A\nu$. Note that we are only referring to the "d-structure" of the bundle E in this calculation. It is therefore valid for any Dirac operator D. It now follows from Lemma 1.11 that $\nabla^\phi_W\nu \in {}^bTX$ for $W \in \Gamma({}^\phi TX)$, and

$$\nabla^\phi_A\nu = -\mathsf{h}\frac{1}{x}A - \frac{dx}{x^2}(A)\mathsf{X}_d + O(x)^bTX, \quad \text{for} \quad A \in \Gamma([{}^\phi N\partial X]).$$

Also, using the special extension described in Lemma 1.8, we find that $W \mapsto \nabla^\phi_{\overline{W}}\nu$ is a well defined tensor in $C^\infty(\partial X, V^*\partial X \otimes {}^\phi TX)$, independent of the vector field used in the definition of the extension (which therefore could be different from ν). Writing

$$2\langle\nabla^\phi_{\overline{W}}\nu, \overline{V}\rangle_\phi = (L_\nu g_\phi)(\overline{W}, \overline{V}), \quad V, W \in \Gamma(V\partial X), \tag{44}$$

we see that this describes yet another aspect of the deviation of g_ϕ from a product metric near the boundary.

Since the Clifford action $\mathbf{c}_d$ is parallel w.r.t. $\nabla^{E,d}$ we can use the special extension introduced in Section 1.4 and decompose

$$\mathbf{c}_d = \overline{\mathsf{n}\mathbf{c}_d} + \overline{\mathsf{v}\mathbf{c}_d}, \quad \nabla^{E,d}_\nu\overline{\mathsf{n}\mathbf{c}_d} = \nabla^{E,d}_\nu\overline{\mathsf{v}\mathbf{c}_d} = 0.$$

Plugging this into (43) we get

$$I_b(\mathsf{D})(iz)\xi = \Pi_\circ\left(-(z+v/2)\mathbf{c}_d(\frac{dx}{x})\xi + (\overline{\mathsf{n}\mathbf{c}_d})\nabla^{E,d}_{\cdot}\xi + \mathbf{c}_d(xF^{E,d}(\nu,\cdot))\xi - \overline{\mathsf{v}\mathbf{c}_d}\nabla^{E,d}_{x\nabla^\phi_{\cdot}\nu}\xi\right).$$

Thus $I_b(\mathsf{D})$ is a b-differential operator on the vector bundle $\mathcal{K} \to B$ pulled back from Y to $B = Y \times \mathbf{R}_+$. As explained, the last summand is an additional *endomorphism* stemming from the fact that we are looking at a situation where the metric g_ϕ is not a product ϕ-metric. Let us now analyze the curvature term in the formula. Of course, this term depends on the choice of Clifford bundle E, but we can always look at the general decomposition

$$F^{E,d}(\nu, A) = \frac{1}{2}\mathbf{c}_d(R^d(\nu, A)) + F^{E/S,d}(\nu, A).$$

The contribution coming from the first summand can then be further analyzed using Proposition 1.14:

$$\begin{aligned}\frac{1}{2}\mathbf{c}_{d,1}\mathbf{c}_d(xR^d(\nu,\cdot_1)) &= \frac{1}{4}(x\mathbf{c}_d, \mathbf{c}_d, \mathbf{c}_d)\langle R^d(\nu,\cdot)\cdot,\cdot\rangle_d \\ &= \frac{1}{4}(\mathsf{v}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d)\langle xR^d(\nu,\cdot)\cdot,\cdot\rangle_d + \frac{1}{2}(\mathsf{v}\mathbf{c}_d, \mathsf{n}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d)\langle S_d(\cdot)\cdot,\cdot\rangle_d(45)\end{aligned}$$

Using the symmetries of the curvature tensor and the description of S_ϕ given in Lemma 1.13, the reader can check that (45)can be written as

$$\frac{1}{2}\mathsf{v}\mathbf{c}_d(\mathrm{Ric}^{d,V}(\mathsf{X}_d,\cdot)) + \frac{v}{2}\mathbf{c}_d(\frac{dx}{x}) + \frac{1}{2}\mathsf{n}\mathbf{c}_d(\,\mathrm{tr}(S_M)) + \frac{1}{4}(\mathsf{n}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d)\langle xB_\phi(\cdot, cdot),\cdot\rangle_d.$$

For $A \in {}^dTX$ we have used the notation

$$\mathrm{Ric}^{d,V}(\mathsf{X}_d, A) = (\mathsf{v}g_d)^{-1}_{aa'}\langle R^d(\mathsf{X}_d,\cdot_a)\cdot_{a'}, A\rangle_d$$

for the "vertical" Ricci curvature and $S_M(\cdot)\mathsf{n}A \in \mathrm{End}(V\partial X)$ is the second fundamental form of the boundary fibration. Definition 1.12 can be used to check that all these terms are in $C^\infty(\partial X, \mathrm{End}(E))$.

To be able to interpret this formula, we need to note some more of the **geometrical data** of this construction:

- $g_{B,b}$ is the metric on ${}^bTB$ given by $(\frac{dx}{x})^2 + g_B|_{x=0}$.
- The Levi Civita connection $\nabla^{B,b} : \Gamma({}^bTB) \to \Gamma(T^*B \otimes {}^bTB)$ is the connection induced by the connection $\nabla^d : \Gamma(\phi^{*b}TB) \to \Gamma(\phi^{*b}TB \otimes T^*Y)$ over the boundary (see also Lemma 1.13). Here, elements of TY act on $\Gamma(\phi^{*b}TB)$ via their *horizontal lift* to TX
- $\mathbf{c}_{B,b} \in \Gamma({}^bTB \otimes \mathrm{End}(\mathcal{K}))$ is the ($\mathbf{R}_+$-invariant) Clifford action induced by $\Pi_\circ \mathsf{h}\mathbf{c}_d|_{\partial X}$ on $\mathcal{K}$ (Here we use $[\mathsf{D}^{\phi,V}, \mathsf{n}\mathbf{c}_d] = 0$ at the boundary).
- The grading operators $\boldsymbol{\epsilon} = \boldsymbol{\epsilon}_{\phi T^*X}$ and $\boldsymbol{\epsilon}_B = \boldsymbol{\epsilon}_{{}^bT^*B}$ induce gradings on $\mathcal{K}$.
- The hermitian metric $\langle \cdot, \cdot \rangle_{\mathcal{K}}$ is the L^2-metric: $\langle \xi, \eta \rangle_{\mathcal{K}} = \phi_*(\langle \xi, \eta \rangle_E \, dvol_Z)$.
- For $A \in C^\infty(Y, \mathrm{End}(\mathcal{K}))$:
$$\mathrm{tr}_{\mathcal{K}}(A) = \int_{\partial X/Y} \mathrm{tr}_E(A) \, dvol_Z = \phi_*[\, \mathrm{tr}_E(A) \, dvol_Z] \in C^\infty(Y)$$
- $\widetilde{\nabla}^{\mathcal{K}}$ is the pullback to B of the compressed connection $\Pi_\circ \nabla^{E,d}$ over Y. Here again, elements of TY act via their *horizontal lift* to TX

In this context the usual problem about the non-unitarity of the compressed connection $\widetilde{\nabla}^{\mathcal{K}}$ arises: For $T \in \Gamma(TB)$ denote by $\phi^*T \in \Gamma(H(\partial X \times \mathbf{R}_+))$ the horizontal lift of T. Then

$$\begin{aligned} T \cdot \langle \xi, \eta \rangle_{\mathcal{K}} &= T \cdot \phi_*[\langle \xi, \eta \rangle_E \, dvol_Z] \\ &= \phi_*[\langle \nabla^{E,d}_{\phi^*T} \xi, \eta \rangle_E \, dvol_Z] + \phi_*[\langle \xi, \nabla^{E,d}_{\phi^*T} \eta \rangle_E \, dvol_Z] + \phi_*[\langle \xi, \eta \rangle_E \, \boldsymbol{\iota}(\phi^*T) \, d \, dvol_Z]. \end{aligned}$$

The last summand in this expression (compare Appendix A.3 or [BGV] Chapter 9) is just

$$-\phi_*[\, \mathrm{tr}(\cdot S_M(\cdot)T) \langle \xi, \eta \rangle_E \, dvol_Z]$$

and the unitary (Clifford-) connection on $\mathcal{K}$ is given by

$$\nabla^{\mathcal{K}}_T \xi := \widetilde{\nabla}^{\mathcal{K}}_T \xi + \frac{1}{2} \Pi_\circ \; \mathrm{tr}(\cdot S_M(\cdot)T)\xi.$$

We can summarize our results as follows:

Proposition 3.15 *In terms of the connection $\nabla^{\mathcal{K}}$ the indicial operator of D has the form*

$$\begin{aligned} I_b(\mathsf{D}) \;&=\; \mathbf{c}_{B,b} \circ \nabla^{\mathcal{K}} \\ &\quad + \Pi_\circ \left[\mathsf{v}\mathbf{c}_d(F^{E/S,d}(\mathsf{X}_d, \cdot)) + \frac{1}{2}\mathsf{v}\mathbf{c}_d(\mathrm{Ric}^{d,V}(\mathsf{X}_d,\cdot)) + \frac{1}{4}(\mathsf{n}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d, \mathsf{v}\mathbf{c}_d)\langle x B_\phi(\cdot, cdot), \cdot\rangle_d - \overline{\mathsf{v}\mathbf{c}}_d \nabla^{E,d}_{x\nabla^\phi_\cdot \nu} \right]. \end{aligned}$$

This is an odd operator w.r.t. $\boldsymbol{\epsilon}$. It is also odd w.r.t. $\boldsymbol{\epsilon}_B$ whenever $h+1$ is even. Recall that the last summand is an endomorphism *on $\mathcal{K}$.* ■

Especially, this Proposition shows that in general $I_b(\mathsf{D})$ is *not* the induced Dirac operator on $\mathcal{K}$! However, the last three summands on the RHS vanish when the metric g_ϕ is a product ϕ-metric. Also, if X is spin and $E = S(X) \otimes E'$, we have $F^{E/S,d}(\nu,\cdot) = F^{E',d}(\nu,\cdot)$, which vanishes whenever the twisting bundle E' has product structure near ∂X.

Inverting the Indicial Family

We now want to analyze the structure of the indicial family $I_b(\mathsf{D}_\lambda)(z)$ in more detail. We emphasize that we make no assumption on the parity of the dimension of the base space Y at this stage. The operator $I_b(\mathsf{D})$ can be written as

$$I_b(\mathsf{D}) := \mathsf{D}_Y + \mathbf{c}_d(\frac{dx}{x})x\frac{\partial}{\partial x} \quad \text{with} \quad \mathsf{D}_Y(\xi|_{\partial X}) = I_b(\mathsf{D})(0)(\xi|_{\partial X}) = \Pi_\circ(\mathsf{D}\xi)|_{\partial X}.$$

Proposition 3.15, or the calculation

$$\begin{aligned} [\mathsf{D}^d, \mathbf{c}_d(\frac{dx}{x})]|_{\partial X} \;&=\; \left(\mathbf{c}_d[\nabla^{E,d}, \mathbf{c}_d(\frac{dx}{x})] + [\mathbf{c}_d, \mathbf{c}_d(\frac{dx}{x})]\nabla^{E,d} \right)|_{\partial X} \\ &=\; (\mathbf{c}_d, \mathbf{c}_d)\mathsf{v}g_d(\cdot,\cdot)|_{\partial X} - 2\nabla^{E,d}_{\mathsf{X}_d}|_{\partial X} = -v, \end{aligned}$$

show that D_Y anticommutes with $\boldsymbol{\sigma} = \mathbf{c}_d(\frac{dx}{x})$. Also, D_Y is selfadjoint with purely discrete spectrum. This implies that for any eigenvalue a of D_Y the negative $-a$ is also an eigenvalue and the corresponding eigenspaces are intertwined by $\boldsymbol{\sigma}$. Denote by $P(\pm a)$ the projector onto the eigenspaces for $\pm a$. These are invariant subspaces for the indicial family $I_b(\mathsf{D}_\lambda)(z)$ and for $a \neq 0$ we get the matrix representation

$$P(\pm a) I_b(\mathsf{D}_\lambda)(z) \mathrel{\hat{=}} \begin{pmatrix} a - \lambda & -iz \\ iz & -a-\lambda \end{pmatrix},$$

w.r.t. a basis of eigenvectors of D_Y of the form $(\boldsymbol{\psi}, \boldsymbol{\sigma}\boldsymbol{\psi})$. The inverse, whenever it exists, has the form

$$P(\pm a) I_b(\mathsf{D}_\lambda)(z)^{-1} \mathrel{\hat{=}} \frac{1}{\lambda^2 - a^2 - z^2} \begin{pmatrix} -a-\lambda & iz \\ -iz & a - \lambda \end{pmatrix}. \tag{46}$$

In the case $a = 0$ we have

$$P(0) I_b(\mathsf{D}_\lambda)(z) = iz\boldsymbol{\sigma} - \lambda, \qquad P(0) I_b(\mathsf{D}_\lambda(z))^{-1} = \frac{1}{z^2 - \lambda^2}(iz\boldsymbol{\sigma} + \lambda). \tag{47}$$

Thus, writing spec(D_Y) for the (necessarily symmetric) set of eigenvalues of D_Y, we find the following description of the set of indicial roots, i.e. those points z, where $I_b(\mathsf{D}_\lambda)$ is not invertible:

$$\operatorname{spec}_I(I_b(\mathsf{D}_\lambda)(z)) = \big\{ \pm\sqrt{\lambda^2 - c_j^2} \,|\quad c_j \in \operatorname{spec}(\mathsf{D}_Y)\big\}.$$

Formula (46,47) shows that these indicial roots are all simple (i.e. the corresponding poles are), as long as λ is not an eigenvalue of D_Y. Also, for $\lambda = 0$ the indicial root in 0 (which exists if 0 is an eigenvalue of D_Y) is always simple. If λ is a nonzero eigenvalue of D_Y, then 0 is an indicial root of order two.

The Model Operator at ϕbf

Having analyzed the behavior of the model operator on the fibres of the face bf, let us now briefly describe the situation at the face $\phi\mathrm{bf} = \mathrm{bf} - F_\phi$. As usual, this face is a fibre bundle

$$\phi\mathrm{bf} \cong \partial X \times \partial X \times [-1,1]_\sigma - F_\phi \xrightarrow{\beta_{\phi,R}} \partial X, \qquad \sigma = \frac{x - x'}{x + x'}.$$

In this compactified picture, the fibre over a point $p \in \partial X$ is the manifold $\partial X \times [-1,1]_\sigma - (Z_{\phi(p)} \times \{0\})$ with boundaries

$$\mathrm{rf} \cap \phi\mathrm{bf}|_p \cong \partial X \times \{1\}, \qquad \mathrm{lf} \cap \phi\mathrm{bf}|_p \cong \partial X \times \{-1\}, \qquad \mathrm{ff} \cap \phi\mathrm{bf}|_p \cong S^\phi N\partial X|_{Z_{\phi(p)}},$$

and the model operator $N_{\phi\mathrm{bf}}(\mathsf{D}_\lambda)$ on the fibres is just the restriction of $N_{\mathrm{bf}}(\mathsf{D}_\lambda)$ to this space. More concretely

Lemma 3.16 *Fix a $p \in \partial X$ and set*

$$B_p = [Y \times [0,\infty[_s;\ \{\phi(p)\} \times \{s = 1\}], \qquad s = x/x'.$$

By restriction, the function ρ_{ff} gives a defining function for the face ff *in B_p obtained by the blow up of $\{\phi(p)\} \times \{s = 1\}$ in B_p. Then*

(a) *$N_{\phi\mathrm{bf}}(\mathsf{D}_\lambda)$ is the operator $I_b(\mathsf{D})$ restricted to $\mathcal{K}$ over B_p*

(b) *$\rho_{\mathrm{ff}} I_b(\mathsf{D})$ is a b-elliptic operator on B_p.*

(c) *The extended set of indicial roots at* ff *of the operator $\rho_{\mathrm{ff}}\, I_b(\mathsf{D}_\lambda)$ is given by*

$$\operatorname{spec}_I(I_{\mathrm{ff}}(\rho_{\mathrm{ff}}\, I_b(\mathsf{D}_\lambda))(z)) = i\mathbf{N} \cup -i((n-1) - \mathbf{N}) \subset \mathbf{Z}.$$

Proof. This Lemma is a special case of a similar statement for elliptic operators on a compact manifold minus a point. An instance of this was presented in Lemma 3.8 and we use this result for the proof of (c). Write $x = \rho_{\mathrm{ff}} \cdot \rho_{\phi\mathrm{bf}}$ around ff and consider the little calculation

$$0 = \rho_{\mathrm{ff}}\, \mathbf{c}_\phi(\frac{dx}{x^2})|_{\mathrm{ff}} = \mathbf{c}_\phi(\frac{d\rho_{\phi\mathrm{bf}}}{\rho_{\phi\mathrm{bf}}^2}) + \frac{\rho_{\mathrm{ff}}}{\rho_{\phi\mathrm{bf}}}\mathbf{c}_\phi(\frac{d\rho_{\mathrm{ff}}}{\rho_{\mathrm{ff}}^2})|_{\mathrm{ff}}, \quad \text{i.e.} \quad \frac{1}{x}\mathbf{c}_\phi(\,d\rho_{\mathrm{ff}})|_{\mathrm{ff}} = -\mathbf{c}_\phi(\frac{d\rho_{\phi\mathrm{bf}}}{\rho_{\phi\mathrm{bf}}^2})|_{\mathrm{ff}}.$$

This allows us to calculate the indicial family at $\mathrm{ff} \cap \phi\mathrm{bf}$ as

$$\begin{aligned} I_{\mathrm{ff}}(\rho_{\mathrm{ff}}\, I_b(\mathsf{D}_\lambda))(-iz) &= \rho_{\mathrm{ff}}^{-z+1} I_b(\mathsf{D}_\lambda)\rho_{\mathrm{ff}}^{z}|_{(\mathrm{ff}\cap\phi\mathrm{bf})_{\phi(p)}} = \rho_{\mathrm{ff}}\, I_b(\mathsf{D}_\lambda)|_{(\mathrm{ff}\cap\phi\mathrm{bf})_{\phi(p)}} + \frac{z}{x}\mathbf{c}_\phi(\, d\rho_{\mathrm{ff}})|_{(\mathrm{ff}\cap\phi\mathrm{bf})_{\phi(p)}} \\ &= \rho_{\mathrm{ff}}\, I_{\phi\mathrm{bf}}(\mathsf{D}_\lambda)|_{(\phi\mathrm{bf}\cap\mathrm{ff})_{\phi(p)}} - z\cdot\mathbf{c}_\phi(\frac{d\rho_{\phi\mathrm{bf}}}{\rho_{\phi\mathrm{bf}}^2})|_{(\mathrm{ff}\cap\phi\mathrm{bf})_{\phi(p)}} = \rho_{\phi\mathrm{bf}}^{z}\frac{1}{\rho_{\phi\mathrm{bf}}}\mathsf{D}^\phi\rho_{\phi\mathrm{bf}}^{-z}|_{(\mathrm{ff}\cap\phi\mathrm{bf})_{\phi(p)}}, \end{aligned}$$

which is just the indicial family whose roots were calculated to be $i\mathbf{N} \cup -i((n-1)-\mathbf{N})$ in Lemma 3.8. ■

3.6 Inversion at ϕbf

In Corollary 3.12 we have constructed the resolvent up to an error term R_1, which does not exhibit decay at the face ϕbf. In this Section we want to improve this error at the face ϕbf. To do this, we first remove the Taylor series of $R_1|_{\phi\mathrm{bf}}$ at the face ff obtaining a remainder lying in the "zero modes" over the blown down face bf. This term can then be removed using the inversion of $I_b(\mathsf{D}-\lambda)$ described in the last Section.

Proposition 3.17 (Reduction to bf)
The equality $(\mathsf{D}-\lambda)Q_2^\phi(\lambda) = 1 + R_2^\phi(\lambda) + R_2^b(\lambda)$ *holds for holomorphic families*

$$\begin{aligned} \lambda &\mapsto R_2^\phi(\lambda) \in \Psi_\phi^{-\infty,\mathcal{F}(\lambda)}(X,E), \quad F(\lambda)_{\mathrm{ff}} = (1+\mathbf{N})\overline{\cup}(\widehat{1+\mathbf{N}}), \quad F(\lambda)_{\phi\mathrm{bf}} = h+2+\mathbf{N} \\ \lambda &\mapsto R_2^b(\lambda) \in \Psi_b^{-\infty,\mathcal{G}(\lambda)}(X,E), \quad G(\lambda)_{\mathrm{bf}} = [\circ]\mathbf{N} \\ \lambda &\mapsto Q_2^\phi(\lambda) \in \Psi_\phi^{-1,\mathcal{E}(\lambda)}(X,E), \quad E(\lambda)_{\phi\mathrm{bf}} = [\circ](h+1+\mathbf{N}), \quad E(\lambda)_{\mathrm{ff}} = (1+\mathbf{N})\overline{\cup}(\widehat{2+\mathbf{N}}), \end{aligned}$$

all other index sets equal ∞.

Proof. In this proof, we will uniformly assume that the coefficient bundle is CB_b and drop it from our notation. Then

$$s(\lambda) = R_1(\lambda)|_{\phi\mathrm{bf}} \in \mathcal{A}^{\mathcal{H}}(\phi\mathrm{bf}), \quad H_{\mathrm{ff}} = -h+\mathbf{N}, \quad H_{\mathrm{lf}} = H_{\mathrm{rf}} = \infty.$$

In order to reduce $s^\circ(\lambda)$ to an error term on the face bf, we have to solve away its expansion at ff. Since by Lemma 3.16 the indicial roots of $\rho_{\mathrm{ff}} I_{\phi\mathrm{bf}}(\mathsf{D}-\lambda)$ are just $i\mathbf{Z}$, the theory of elliptic b-differential operators (see Lemma 5.44 in [Me3], or Lemma 3.20 below for a similar statement) tells us that the equation

$$I_{\phi\mathrm{bf}}(\mathsf{D}-\lambda)g^\circ(\lambda) - s^\circ(\lambda) = u^\circ(\lambda)$$

for the zero-mode part can be solved with

$$g^\circ(\lambda) \in \Pi_\circ\mathcal{A}^{\mathcal{I}}(\phi\mathrm{bf}), \quad I_{\mathrm{ff}} = (-h+1+\mathbf{N})\overline{\cup}(\widehat{-h+1+\mathbf{N}}), \quad I_{\mathrm{lf}} = I_{\mathrm{rf}} = \infty,$$

$$\text{and} \quad u^\circ(\lambda) \in \Pi_\circ\, \dot{C}^\infty(\phi\mathrm{bf}) \subset \Pi_\circ\, \dot{C}^\infty(\mathrm{bf}).$$

Extend g° and u° to kernels $G^\circ(\lambda)$ and $U^\circ(\lambda)$ on X^2_ϕ. All this can be done holomorphically in λ. Noting that

$$G^\circ(\lambda) \in \mathcal{A}^{\mathcal{I}}(X^2_\phi), \quad I_{\mathrm{ff}} = (1-h+\mathbf{N})\overline{\cup}(1\widehat{-h+}\mathbf{N}), \quad I_{\phi\mathrm{bf}} = [\circ]\mathbf{N}, \quad I_{\mathrm{rf}} = I_{\mathrm{lf}} = \infty,$$

we get, using Lemma A.17,

$$T(\lambda) := (\mathsf{D}-\lambda)G^\circ(\lambda)-S(\lambda)-U^\circ(\lambda) \in \mathcal{A}^{\mathcal{J}}(X), \quad J_{\mathrm{ff}} = (1+\mathbf{N})\overline{\cup}(\widehat{1+\mathbf{N}}), J_{\phi\mathrm{bf}} = [\perp](h+1+\mathbf{N}).$$

It remains to solve away $t(\lambda)$, the restriction of $T(\lambda)$ to ϕbf. To do this, just define $G^\perp(\lambda)$ to be x' times the extension of $(\mathsf{D}^{\phi,V})^{-1}t(\lambda)$ to X^2_ϕ. Then, setting

$$Q^\phi_2(\lambda) := Q_1(\lambda) + G^\circ(\lambda) + G^\perp(\lambda), \quad R^b_2(\lambda) = U^\circ(\lambda), \quad R^\phi_2(\lambda) = (\mathsf{D}-\lambda)G^\perp(\lambda) - T(\lambda)$$

proves the claim. ■

Meromorphic Solution at bf

For the formulation of the next Proposition, recall that the indicial roots of the operator $I_b(\mathsf{D}-\lambda)$ were given by $\pm\sqrt{\lambda^2 - c_j^2}$, where the c_j denote the eigenvalues of D_Y. We have agreed implicitly to read this as $\pm\lambda$ if $c_j = 0$. For $\mathrm{Im}(\lambda) > 0, a \in \mathbf{R} - \{0\}$ we have

$$\sqrt{\lambda^2 - a^2} = \sqrt{\lambda - a}\sqrt{\lambda + a}.$$

This can be continued to a meromorphic function on the Riemann surface $\Sigma_a \to \mathbf{C}$, defined as the minimal branched covering of $\mathbf{C}$ with that property, with poles only at the branching points $\pm a$.

Denote by $\Sigma \to \mathbf{C}$ the (infinite) branched covering over $\mathbf{C}$, onto which all the functions $\sqrt{\lambda^2 - c_j^2}$ can be continued to be meromorphic, with poles only at the branching points. This means that Σ has branching points over $\pm c_j \neq 0$ around which it is uniformized by the functions $\sqrt{\lambda \mp c_j}$. We emphasize that 0 is not a branching point of this covering. We will denote points over $\lambda \in \mathbf{C}$ by $\Lambda \in \Sigma$ and write $\theta_j(\Lambda)$ for the function on Σ, obtained by continuation of $\theta_j(\lambda) = -i\sqrt{\lambda^2 - c_j^2}$ from $\mathrm{Im}(\lambda) > 0$. Note that on $\mathrm{Im}(\lambda) > 0$ the function $\theta_j(\lambda)$ has positive real part.

More concretely, introduce the *physical domain* PD as that part of the preimage of the upper half plane $\{\lambda | \ \mathrm{Im}(\lambda) > 0\}$ in Σ in which $\theta_j(\Lambda) = \theta_j(\lambda)$ or, equivalently, where all $\theta_j(\Lambda)$ have positive real part. Define the set $\Theta(\Lambda)$ as the smallest C^∞-index set containing $(\theta_j(\Lambda) | j \in \mathbf{N})$. This is a generating set for $\Theta(\Lambda)$ and a minimal such *away* from accidental multiplicities, i.e. in the sense of Appendix A.6:

$$\mathrm{lead}(\Theta(\Lambda)) = (\theta_j(\Lambda) | j \in \mathbf{N}), \qquad \text{if} \quad \theta_j(\Lambda) \notin \theta_k(\Lambda) + \mathbf{N} \quad \text{for} \quad j \neq k.$$

Let us note some important properties of these index sets as a

Lemma 3.18 (a) $0 < \mathrm{Re}(\theta_j(\Lambda)) < \mathrm{Re}(\theta_k(\Lambda)) \quad$ *for* $\Lambda \in \mathrm{PD}, j < k$

(b) $\theta_j(-r) = \overline{\theta_j(r)} \quad (\ = -\theta_j(r) \quad$ *if* $\quad |r| \geq c_j)$ *for* $r \in \mathbf{R} = \partial\mathrm{PD}$

(c) *The set of accidental multiplicities* $\mathcal{M} \subset \Sigma$ *is discrete and* $\mathcal{M} \cap \mathrm{PD} \subset i]0, \infty[$.

Proof. An accidental multiplicity in PD is a point λ with $\operatorname{Im}(\lambda) > 0$ such that

$$-i\sqrt{\lambda^2 - a^2} = -i\sqrt{\lambda^2 - b^2} + m,$$

with a, b positive eigenvalues of D_Y and $m \in \mathbf{Z}$. This can be rewritten as

$$(2m\lambda)^2 = [(b+m)^2 - a^2][a^2 - (b-m)^2],$$

from which the assertion in (c) immediately follows. ■

We can now formulate the next step in the construction of the resolvent

Proposition 3.19 *The equation* $(\mathsf{D} - \lambda)Q_3^b(\Lambda) = -R_2^b(\Lambda) + R_3^b(\Lambda)$ *can be solved with*

$$\begin{array}{lcll} Q_3^b(\Lambda) & \in & \Psi_b^{-\infty,\mathcal{E}}(X,E), & E(\Lambda)_{\mathrm{lf}} = [\circ]\Theta(\Lambda), \quad E(\Lambda)_{\mathrm{rf}} = \Theta(\Lambda), \quad E(\Lambda)_{\mathrm{bf}} = [\circ]\mathbf{N}, \\ R_3^b(\Lambda) & \in & \Psi_b^{-\infty,\mathcal{F}}(X,E), & F(\Lambda)_{\mathrm{lf}} = 1 + \Theta(\Lambda), \; F(\Lambda)_{\mathrm{rf}} = \Theta(\Lambda), \; F(\Lambda)_{\mathrm{bf}} = 1 + \mathbf{N}. \end{array}$$

The maps

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto Q_3^b(\Lambda) \in \Psi_b^{-\infty,(E_{\mathrm{bf}},-R,-R)}(X,E),$$

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto R_3^b(\Lambda) \in \Psi_b^{-\infty,(F_{\mathrm{bf}},-R,-R)}(X,E)$$

into the calculus with bounds are meromorphic with poles only at the branching points of Σ.

Proof. We remind the reader that we will be working with the coefficient bundle CB_b. Assume for the moment that $\operatorname{Im}(\lambda) > 0$, i.e. $I_b(\mathsf{D} - \lambda)$ does not have any real indicial roots. Also assume for simplicity that λ is not an accidental multiplicity i.e. $\theta_j(\lambda) \notin \theta_k(\lambda)$ for $j \neq k$. Since

$$v^\circ(\lambda) := -R_2^b(\lambda)|_{\mathrm{bf}} \in \Pi_\circ \dot{C}^\infty(\mathrm{bf}),$$

we can solve $I_{\mathrm{bf}}(\mathsf{D} - \lambda)g^\circ(\lambda) = -v^\circ(\lambda)$ with

$$g^\circ(\lambda) \in \Pi_\circ \mathcal{A}^{\mathcal{G}}(\mathrm{bf}), \quad G_{\mathrm{lf}} = \Theta(\lambda), \quad G_{\mathrm{rf}} = \Theta(\lambda).$$

We can extend $g^\circ(\lambda)$ to X_b^2 in such a way that all the leading coefficients of $\operatorname{lead}(\Theta(\lambda))$ at lf are in the zero modes and independent of x' close to bf. This gives an element

$$G^\circ(\lambda) \in \mathcal{A}^{\mathcal{H}}(X_b^2), \quad H_{\mathrm{lf}} = [\circ]\Theta(\lambda), \quad H_{\mathrm{rf}} = \Theta(\lambda), \quad H_{\mathrm{bf}} = \mathbf{N}.$$

This operator can be used to solve away the error term $R_2^b(\lambda)$. By Lemma A.17 we have

$$-T(\lambda) := (\mathsf{D} - \lambda)G^\circ(\lambda) - R_2^b(\lambda) \in \mathcal{A}^{\mathcal{I}}(X_b^2), \quad I_{\mathrm{bf}} = [\perp]\mathbf{N}, \; I_{\mathrm{lf}} = [\perp]\Theta(\lambda), \; I_{\mathrm{rf}} = \Theta(\lambda),$$

where the form of I_{lf} is due to our special choice of extension. We are left with the problem of solving away an error perpendicular to the zero-modes.

This is done as usual by extending $(\mathsf{D}^{\phi,V})^{-1}T(\lambda)|_{\mathrm{bf}}$, to X_b^2 such that the coefficients associated to $[\perp]\mathrm{lead}(\Theta(\lambda))$at lf are just $(\mathsf{D}^{\phi,V})^{-1}$ applied to the corresponding coefficients of $T(\lambda)$ at lf. Define $G^{\perp}(\lambda)$ to be x times this extension, thus

$$G^{\perp}(\lambda) \in \mathcal{A}^{\mathcal{J}}(X_b^2), \qquad J_{\mathrm{lf}} = [\perp]\Theta(\lambda) + 1, \quad J_{\mathrm{rf}} = \Theta(\lambda), \quad J_{\mathrm{bf}} = [\perp](1 + \mathbf{N}).$$

With Lemma A.17 it is now easy to see that, setting $Q_3^b(\lambda) = G^{\circ}(\lambda) + G^{\perp}(\lambda)$, we have

$$R_3^b(\lambda) = (\mathsf{D} - \lambda)Q_2^b(\lambda) - R_2^b(\lambda) = (\mathsf{D} - \lambda)G^{\perp}(\lambda) - T(\lambda),$$

with index sets as indicated in the Proposition.

To prove the claimed meromorphy, a more careful analysis of the term $g^{\circ}(\lambda)$ above is needed. Using $s = x/x'$, recall that since $v^{\circ} \in \Pi_{\circ}\, \dot{C}^{\infty}(\mathrm{bf})$, the Mellin transform

$$M(v^{\circ})(z) = \int_0^{\infty} s^{iz} v^{\circ}(s) \frac{ds}{s} \tag{48}$$

is holomorphic and (still assuming $\mathrm{Im}(\lambda) > 0$) we have

$$-g^{\circ}(\lambda)(s) = I_{\mathrm{bf}}(\mathsf{D} - \lambda)^{-1} v^{\circ}(\lambda)(s) = \frac{1}{2\pi} \int_{\mathrm{Im}(z)=0} s^{-iz} I_{\mathrm{bf}}(\mathsf{D} - \lambda)(z)^{-1} M(v^{\circ})(z)\, dz \tag{49}$$

The inverse of the indicial family has the form (46, 47). The integral thus is well defined, as long as λ stays away from the real axis. When $|\lambda| < R/\sqrt{2}$ approaches the real axis from $\mathrm{Im}(\lambda) > 0$, only the eigenvalues smaller than $R/\sqrt{2}$ contribute a singularity. Denote by $c_1, \ldots, c_N$ the eigenvalues of D_Y which are smaller than a given $L > 2R$. Writing $P(c_j)$ and $P(\geq L)$ for the projections onto the corresponding eigenspaces of D_Y, we get the decomposition

$$\begin{aligned} I_b(\mathsf{D} - \lambda)(z)^{-1} \;&=\; P(\geq L) I_b(\mathsf{D} - \lambda)(z)^{-1} + P(0)\frac{1}{z^2 - \lambda^2}(iz\boldsymbol{\sigma} + \lambda) \\ &\quad + \sum_{j=1}^{N} P(\pm c_j) \frac{1}{\lambda^2 - c_j^2 - z^2} \begin{pmatrix} -c_j - \lambda & iz \\ -iz & c_j - \lambda \end{pmatrix}. \end{aligned} \tag{50}$$

The first term in this decomposition is uniformly bounded in z and has no poles in the strip $-L/2 \leq \mathrm{Im}(z) \leq L/2$. Since the term $M(v^{\circ})(z)$ is rapidly decreasing at real infinity, performing the integral in (49) for this part yields a section in $\mathcal{A}^{(L/2,L/2)}(\mathrm{bf})$, which depends holomorphically on $|\lambda| < R/\sqrt{2}$.

Plugged into (49), the summands of the third term yield integrals of the form

$$\frac{1}{2\pi} \int_{\mathrm{Im}(z)=0} s^{-iz} P(\pm c_j) \frac{1}{\lambda^2 - c_j^2 - z^2} \begin{pmatrix} -c_j - \lambda & iz \\ -iz & c_j - \lambda \end{pmatrix} M(v^{\circ})(z)\, dz.$$

For small s this can be evaluated by shifting contours to $\mathrm{Im}(z) \gg 0$. From the residue theorem (using the same method for the term stemming from the 0-eigenvalue) we get an expansion for $g^{\circ}(\lambda)(s)$ of the form

$$A_0(\lambda) s^{-i\lambda} + \sum_{j=1}^{N} A_j(\Lambda) \frac{s^{\theta_j(\Lambda)}}{\theta_j(\Lambda)} + O_{\Lambda}(s^{L/2}). \tag{51}$$

Here, the coefficients A_j are holomorphic with values in the $\pm c_j$-eigenspaces of D_Y. The expansion for $1/s \to 0$ can be similarly obtained by shifting contours to $\operatorname{Im}(z) \gg 0$.

The expression (51) is meromorphic in the sense of being an expansion with meomorphic coefficients. Its extension to X_b^2 is still meromorphic in this strong sense, i.e. as an element in $\Psi_b^{-\infty,\mathcal{E}(\Lambda)}(X)$, as long as we stay away from accidental multiplicities. Away from these points, the extensions of the leading coefficients A_j of the expansion can (be clearly distinguished and therefore) still be chosen to lie in the $\pm c_j$-eigenspaces of D_Y.

Around points of accidental multiplicity, the extension of (51) is still meromorphic in the weaker sense claimed in the Proposition. ■

Removal of the Remainder at lf

The next step in the parametrix construction is to solve away the error terms at the left face lf. The approach to this problem is based on the following Lemma:

Lemma 3.20 *For any family of sections $f(\Lambda) \in \mathcal{A}^{1+\Theta(\Lambda)}(X, E)$ there is a family $u(\Lambda) \in \mathcal{A}^{1+\Theta(\Lambda)}(X, E)$, such that $(\mathsf{D} - \lambda)u(\Lambda) - f(\Lambda) \in \dot{C}^{\infty}(X, E)$. The construction gives a meromorphic map*

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto u(\Lambda) \in \mathcal{A}^{-R}(X, E),$$

whenever $f(\Lambda)$ is meromorphic in that sense.

Proof. This Lemma is a variation of Lemma 5.44 in [Me3], with an easier proof. Again, we first assume that $\operatorname{Im}(\lambda) > 0$. We have to show how to solve for a section f of the form $f = x^{\theta_j(\lambda)+m} g(\lambda)$, with $g(\lambda)$ meromorphic in the C^∞-sections over X and $m \in \mathbf{N}_+$. As usual, decompose $g = g^\circ + g^\perp$ with $g^\circ \in C^\infty(X, E)^\circ$ and $g^\perp \in C^\infty(X, E)^\perp$. Starting with g°, we can just set

$$u_0^\circ(\lambda) = x^{\theta_j(\lambda)+m} I_b(\mathsf{D} - \lambda)(i(\theta_j(\lambda) + m))^{-1} g^\circ(\lambda), \tag{52}$$

which, according to (46, 47), continues to be meromorphic with extra poles arising only at points of accidental multiplicity.

Ansatz (52) solves our problem to first order in the zero modes:

$$(\mathsf{D} - \lambda)u_0^\circ(\lambda) - x^{\theta_j(\lambda)+m} g(\lambda) = x^{\theta_j(\lambda)+m} v_0(\lambda)^\perp.$$

Again, solving for $x^{\theta_j(\lambda)+m} v_0(\lambda)^\perp$ is easy. Just set

$$u_0^\perp = x^{\theta_j(\lambda)+m+1} (D^{\phi,V})^{-1} v_0^\perp(\lambda)$$

This solves the problem to first order. Iteration of this procedure finishes the proof. ■

The obvious parametrized modification of this Lemma can now be used to get rid of the expansion of the error $R_3^b(\lambda)$ at the left boundary of X_b^2.

Proposition 3.21 *The equation $D_\lambda Q_4^b(\Lambda) = -R_3^b(\Lambda) + R_4^b(\Lambda)$ can be solved with*

$$\begin{aligned} R_4^b(\Lambda) &\in \Psi_b^{-\infty,\mathcal{F}}(X, E), \quad F(\Lambda)_{\mathrm{rf}} = \Theta(\Lambda), \quad F(\Lambda)_{\mathrm{lf}} = \infty, \quad F(\Lambda)_{\mathrm{bf}} = 1 + \mathbf{N}, \\ Q_4^b(\Lambda) &\in \Psi_b^{-\infty,\mathcal{E}}(X, E), \quad E(\Lambda)_{\mathrm{lf}} = 1 + \Theta(\Lambda),\ E(\Lambda)_{\mathrm{rf}} = \infty,\ E(\Lambda)_{\mathrm{bf}} = 1 + \mathbf{N}. \end{aligned}$$

The maps

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto Q_4^b(\Lambda) \in \Psi_b^{-\infty,(E_{\mathrm{bf}},-R,-R)}(X,E),$$

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto R_4^b(\Lambda) \in \Psi_b^{-\infty,(F_{\mathrm{bf}},\infty,-R)}(X,E)$$

are meromorphic. ■

Remark 3.22 Note that, eventhough $\lambda = 0$ might be a point of accidental multiplicity, the coefficient of the highest power $s^{-i\lambda}$ in the expansions of $Q_3^b(\Lambda)$ and $Q_4^b(\Lambda)$, will be meromorphic near 0. We will use this fact in Section 3.8, but we abstain from introducing even more specialized notation for this.

3.7 End of the Construction

So far, we have found a parametrix of the type

$$(\mathsf{D} - \lambda)Q(\Lambda) = I + R(\Lambda), \qquad \text{where}$$

$$Q(\Lambda) := Q_2^\phi(\lambda) + Q_3^b(\Lambda) + Q_4^b(\Lambda), \quad R(\Lambda) := R_2^\phi(\lambda) + R_4^b(\Lambda)$$

have the properties described in the above series of Propositions – in a loose sense, we have obtained a remainder term R, which vanishes to first order at the faces ff, ϕbf and to infinite order at lf. From the composition formula in Theorem 2.11 its powers are of the form

$$R^j \in \Psi_\phi^{-\infty,\mathcal{E}_j}(X,E), \quad E_{j,\mathrm{ff}} \geq j - \varepsilon, \quad E_{j,\phi\mathrm{bf}} \geq h + 1 + j - \varepsilon, \quad E_{j,\mathrm{lf}} = \infty.$$

The ε is used to take care of the "log's" which appear at the face ff. The expansion of R^j at the right face does not improve with increasing j, but becomes eventually constant in any fixed compact range of powers. Thus, the Neumann series for $(I + R)^{-1}$ makes sense as the asymptotic sum

$$(1 + R)^{-1} = \sum_{k=0}^{\infty} (-R)^k = 1 + \widetilde{R}$$

at the faces lf, ϕbf, and ff, and setting

$$Q_{\mathrm{sm}}(\Lambda) = Q(\Lambda)(1 + R)^{-1} = Q(\Lambda) + Q(\Lambda)\widetilde{R}(\Lambda) =: Q(\Lambda) + \widetilde{Q}(\Lambda)$$

we get

$$\mathsf{D}_\lambda Q_{\mathrm{sm}}(\Lambda) = 1 + R_{\mathrm{sm}}(\Lambda), \qquad R_{\mathrm{sm}}(\Lambda) = \text{"}R^\infty(\Lambda)\text{"}$$

To easily describe the index sets for these operators, we allow for a little sloppiness in the notation: Given an index set $\mathcal{F}$, denote by

$$T\mathcal{F} := \mathcal{F}|_{[\inf(\mathcal{F}),\inf(\mathcal{F})+1[} \tag{53}$$

the finite index set which coincides with $\mathcal{F}$ to first order. Also, we will write down the index set at ff w.r.t. the ϕ-density, and those at the faces ϕbf, lf and rf w.r.t. the b-density, in order to avoid the notorious term "$h+1$". Then

$$\begin{aligned}
\widetilde{R} \in \Psi_\phi^{-\infty,\mathcal{G}}(X,E), &\quad G_{\mathrm{lf},b} = \infty,\ TG_{\phi\mathrm{bf},b} = \{(1,0)\},\ TG_{\mathrm{rf},b} = T\Theta(\Lambda),\\
&\quad TG_{\mathrm{ff},\phi} = \{(1,0),(1,1)\}\\
R_{\mathrm{sm}} \in \Psi^{-\infty,\mathcal{H}}(X,E), &\quad TH_{\mathrm{rf},b} = T\Theta(\Lambda),\ TH_{\mathrm{lf},b} = \infty\\
\widetilde{Q} \in \Psi_\phi^{-1,\mathcal{I}}(X,E), &\quad I_{\mathrm{lf},b} = [\circ]\Theta(\Lambda),\ T_{<1}I_{\phi\mathrm{bf},b} = [\circ]T_{<1}(\Theta(\Lambda)+\Theta(\Lambda)),\\
&\quad TI_{\mathrm{rf},b} = T\Theta(\Lambda),\ T_{<1}I_{\mathrm{ff},\phi} = [\circ]T_{<1}(\Theta(\Lambda)+\Theta(\Lambda))\\
Q_{\mathrm{sm}} \in \Psi_\phi^{-1,\mathcal{J}}(X,E), &\quad J_{\mathrm{lf},b} = [\circ]\Theta(\Lambda),\ T_{<1}J_{\phi\mathrm{bf},b} = [\circ]T_{<1}(\{0\}\cup\Theta(\Lambda)+\Theta(\Lambda)),\\
&\quad TJ_{\mathrm{rf},b} = T\Theta(\Lambda),\ T_{<1}J_{\mathrm{ff},\phi} = [\circ]T_{<1}(\Theta(\Lambda)+\Theta(\Lambda)) \qquad (54)
\end{aligned}$$

Also, to first order, the expansion of $R_{\mathrm{sm}}(\Lambda)$ at the right face is of type (51) with s replaced by s^{-1}.

The removal of this smoothing remainder is now standard. Since, for instance,

$$R_{\mathrm{sm}}(i) : L_b^2(X,E) \longrightarrow \dot{C}^\infty(X,E),$$

the kernel of the projection M onto the null space of $1+R_{\mathrm{sm}}(i)$ lies in $\dot{C}^\infty(X^2,\mathrm{CB}_b)$. Setting $Q_{\mathrm{inv}}(\Lambda) = Q_{\mathrm{sm}}(\Lambda) + M$ yields

$$(\mathsf{D}-\lambda)Q_{\mathrm{inv}}(\Lambda) = 1 + R_{\mathrm{sm}}(\Lambda) + (\mathsf{D}-\lambda)M = 1 + R_{\mathrm{inv}}(\Lambda),$$

where the new remainder R_{inv} has the same expansions as R_{sm}. Therefore the remainder is a compact family (compare Corollary 2.6) of the form:

$$R_{\mathrm{inv}} : \Sigma \subset B_{R/\sqrt{2}}(0) \longrightarrow \mathcal{K}(x^R L_b^2(X,E)).$$

Since $1+R_{\mathrm{inv}}(i)$ is invertible by construction we infer from analytic Fredholm theory that the inverse of $1+R_{\mathrm{inv}}(\Lambda)$ is a meromorphic family of the same type:

Proposition 3.23 *The inverse* $(1+R_{\mathrm{inv}}(\Lambda))^{-1}$ *is of the form* $1+S(\Lambda)$*, where*

$$S(\Lambda) \in \Psi^{-\infty,(\infty,H_{\mathrm{rf}})}(X,E), \quad TH_{\mathrm{rf}} = T\Theta(\Lambda),$$

is meromorphic as a map

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto \Psi^{-\infty,(\infty,-R)}(X,E).$$

Proof. The proof closely follows the proof of the analytic Fredholm theorem. For a Λ_0 in $B_{R/\sqrt{2}}(0)$ denote by $M_0 = M(\Lambda_0) \in \dot{C}^\infty(X^2)$ the projection (of finite rank) onto the null space of $1+R_{\mathrm{inv}}(\Lambda_0)$. Then w.r.t. $1-M_0$ and M_0 the operator $1+R_{\mathrm{inv}}(\Lambda)$ can be written as the matrix

$$1+R_{\mathrm{inv}}(\Lambda) = \begin{pmatrix} A(\Lambda) & B(\Lambda) \\ C(\Lambda) & D(\Lambda) \end{pmatrix}.$$

The operator $A(\Lambda)$ is of the form $A(\Lambda) = 1 + T(\Lambda)$, with

$$T(\Lambda) = R_{\mathrm{inv}}(\Lambda) - M_0 R_{\mathrm{inv}}(\Lambda) - R_{\mathrm{inv}}(\Lambda)M_0 + M_0 R_{\mathrm{inv}}(\Lambda) M_0 \in \Psi^{-\infty,(\infty,H_{\mathrm{rf}})}(X,E).$$

Also $A(\Lambda)$ is invertible for Λ close to Λ_0 with an inverse of the same type. To see this, we write

$$(1+T)^{-1} = 1 - T + T(1+T)^{-1}T,$$

and use the following "semi-ideal" property of the space $\Psi_b^{-\infty,(\infty,a,b)}(X,E)$, which is described in [Me3], and which shows that the last term in the above sum is of the same type as the operator T:

Lemma 3.24 *For* $Q_1, Q_2 \in \Psi_b^{-\infty,(\infty,a,b)}(X,E)$ *and* $A \in \mathcal{L}(x^{-b}L_b^2(X,E))$ *we have* $Q_1 A Q_2 \in \Psi_b^{-\infty,(\infty,a,b)}(X,E)$. ■

The rest of the proof now follows as usual: Assuming the invertibility of $A(\Lambda)$, the operator $1 + R_{\mathrm{inv}}(\Lambda)$ is invertible exactly if the endomorphism $D - CA^{-1}B$ is invertible, since

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}^{-1} = \begin{pmatrix} A^{-1} + A^{-1}B(D-CA^{-1}B)^{-1}CA^{-1} & -A^{-1}B(D-CA^{-1}B)^{-1} \\ -(D-CA^{-1}B)^{-1}CA^{-1} & (D-CA^{-1}B)^{-1} \end{pmatrix}.$$

Thus the inverse exists, and is of the claimed form, whenever $\det(D - CA^{-1}B) \neq 0$. The determinant is not constantly 0, since $1 + R_{\mathrm{inv}}(i)$ is invertible by construction. ■

Putting everything together, we obtain one of the central results in this work

Theorem 3.25 (The Resolvent of D) *The resolvent of* D *continued meromorphically to* Σ *from* $\mathrm{Im}(\lambda) > 0$ *is given by*

$$G(\Lambda) \equiv G^-(\lambda) := (Q_{\mathrm{sm}}(\Lambda) + M)(1 + R_{\mathrm{inv}}(\Lambda))^{-1} \in \Psi_\phi^{-1,\mathcal{J}}(X,E),$$

with index set $\mathcal{J}$ *as in (54). It is meromorphic as a map*

$$\Sigma \supset B_{R/\sqrt{2}}(0) \ni \Lambda \mapsto G^-(\Lambda) \in \Psi_\phi^{-1,(-R,-2R,-2R,-R)}(X,E),$$

■

Remark 3.26 Let us end this Section with a remark about some variants of this construction: First, it is possible to construct the meromorphic continuation of $(\mathsf{D} - \lambda)^{-1}$ from $\mathrm{Im}(\lambda) < 0$. This will be needed in view of Stone's formula (62). Equivalently, we can use the continuation $G^+(\Lambda)$ of $(\mathsf{D} + \lambda)^{-1}$ from $\mathrm{Im}(\lambda) > 0$. The results in this Chapter hold *verbatim* for this continuation (with λ replaced by $-\lambda$, *but the terms* Λ, $\theta_j(\Lambda)$ *etc. remain unchanged!*).

Also, a generalized inverse for $\mathsf{D} - \lambda_0$ can be constructed for any fixed $\lambda_0 \in \mathbf{C}$. For $\alpha \in \mathbf{R}$ not equal to the real part of any $\pm\theta_j(\lambda_0)$ we define the C^∞-index set

$$\Theta_\alpha(\lambda_0) := \{s = \pm\theta_j(\lambda_0) \mid \quad \mathrm{Re}(s) > \alpha\} + \mathbf{N}.$$

Then the construction of the parametrix as an operator on $x^\alpha L_b^2(X,E)$ up to a smoothing term can be followed through as before. The integral in (49) is now over $\mathrm{Im}(z) = \alpha$ and the index set $\Theta(\lambda_0)$ has to be replaced by $\Theta_\alpha(\lambda_0)$ at the left face and by $\Theta_{-\alpha}(\lambda_0)$ at the right face. The constructions in this Section and Section 3.6 then yield

Proposition 3.27 $\mathsf{D}_{\lambda_0} Q_{\mathrm{sm},\alpha}(\lambda_0) = 1 + R_{\mathrm{sm},\alpha}(\lambda_0)$, *holds with* $R_{\mathrm{sm},\alpha} \in \Psi^{-\infty,\mathcal{H}}(X,E)$, $Q_{\mathrm{sm},\alpha} \in \Psi_\phi^{-1,\mathcal{J}}(X,E)$, *and index sets*

$$\begin{aligned} &TH_{\mathrm{rf},b} = T\Theta_{-\alpha}(\lambda_0), \quad TH_{\mathrm{lf},b} = \infty \\ &J_{\mathrm{lf},b} = [\circ]\Theta_\alpha(\Lambda), \quad T_{<1}J_{\phi\mathrm{bf},b} = [\circ]T_{<1}(\{0\} \cup \Theta_\alpha(\lambda_0) + \Theta_{-\alpha}(\lambda_0)), \\ &TJ_{\mathrm{rf},b} = T\Theta_{-\alpha}(\lambda_0), \quad T_{<1}J_{\mathrm{ff},\phi} = [\circ]T_{<1}(\Theta_\alpha(\lambda_0) + \Theta_{-\alpha}(\lambda_0)). \end{aligned}$$

Especially, the operator $(\mathsf{D} - \lambda_0) : x^\alpha L^2_b(X,E) \to x^\alpha L^2_b(X,E)$ *is Fredholm, if* $\alpha \neq \mathrm{Re}(\pm\theta_j(\lambda_0))$. ■

3.8 Consequences

In this Section we present some standard applications of our construction of the resolvent. Our presentation here is an adaptation of the developments in Chapter 6 of [Me3] to the case of our Dirac operator D. As a special feature, this approach allows a description of the continuous part of the spectral measure for D, based solely on Stone's formula, without using any of the machinery of Hilbert space scattering theory.

The arguments in this Section do not really differ from those in the *b*-case. They have been included here, since a detailed treatment of the Dirac operator along these lines does not seem to exist in the literature. A treatment of the *b*-Laplace operator can be found in [Ch].

Structure of the Null Spaces

As an immediate application of the resolvent construction, and its extension described at the end of the last Section, we can describe the structure of sections in the weighted null spaces of $\mathsf{D} - \lambda_0$.

Proposition 3.28 *Recall the notation introduced in (53). For* $\lambda_0 \in \mathbf{C}$, $r_0 \in \mathbf{R}$ *and* $\alpha \in \mathbf{R}$ *such that* $\alpha \neq \mathrm{Re}(\pm\theta_j(\lambda_0))$ *we have*

(a) $\mathrm{null}_{x^\alpha L^2_b}(\mathsf{D} - \lambda_0) \subset \mathcal{A}^{[\circ]I_\alpha(\lambda_0)}(X,E), \qquad TI_\alpha(\lambda_0) = T\Theta_\alpha(\lambda_0),$

(b) $\mathrm{null}_{L^2_b}(\mathsf{D} - r_0) \subset x^\varepsilon H^\infty_b(X,E), \quad \mathrm{null}_{L^2_b}(\mathsf{D}^2 - r_0) \subset x^\varepsilon H^\infty_b(X,E)$

(c) $\mathrm{null}_-(\mathsf{D} - r_0) := \bigcap_{\varepsilon>0} \mathrm{null}_{x^{-\varepsilon}L^2_b}(\mathsf{D} - r_0) \subset \mathcal{A}^{[\circ]I(r_0)}, \qquad TI(r_0) = T\Theta(r_0).$

(d) $\mathrm{null}_-(\mathsf{D}^2 - r_0^2) := \bigcap_{\varepsilon>0} \mathrm{null}_{x^{-\varepsilon}L^2_b}(\mathsf{D}^2 - r_0^2) \subset \mathcal{A}^{[\circ]J(r_0)}$, *with*

$$TJ(r_0) = T[\Theta_0(-r_0)\overline{\cup}(\Theta(r_0) \cup (\Theta(r_0) + \Theta_0(-r_0))\overline{\cup}(\Theta(r_0) + \Theta_0(-r_0) + \Theta(-r_0))].$$

All of these spaces are of finite dimension.

Proof. Let ξ be a section in $x^\alpha L^2_b(X,E)$ such that $(\mathsf{D} - \lambda_0)\xi = 0$. Then, starting with the parametrix described in Proposition 3.27, we get the equation

$$(\mathsf{D} - \overline{\lambda_0})Q_{\mathrm{sm},-\alpha}(\overline{\lambda_0}) = 1 + R_{\mathrm{sm},-\alpha}(\overline{\lambda_0})$$

on $x^{-\alpha}L_b^2(X,E)$. The adjoint of this equation can then be applied to ξ:

$$0 = Q_{\mathrm{sm},-\alpha}(\overline{\lambda_0})^*(\mathsf{D}-\lambda_0)\xi = (1 + R_{\mathrm{sm},-\alpha}(\overline{\lambda_0})^*)\xi.$$

Since by Lemma 2.7 the remainder $R_{\mathrm{sm},-\alpha}(\overline{\lambda_0})^*$ has an expansion with index set of type $T\Theta_\alpha(\lambda_0)$ at the left face, we find $\xi \in \mathcal{A}^{[\circ]I_\alpha(\lambda_0)}(X,E)$, with $I_\alpha(\lambda_0)$ as above. Part (b) is an immediate consequence of this, since all indeces in $\Theta_0(r_0)$ have positive real part.

The proof of (c) then follows from this, once one recalls that $\Theta(r_0)$ is obtained from $\Theta(\lambda)$ by continuation from $\mathrm{Im}(\lambda) > 0$. Alternatively

$$\Theta(r_0) = \bigcap_{\alpha>0} \Theta_{-\alpha}(r_0).$$

All of these null spaces are of finite dimension, since the operator $(\mathsf{D}-\lambda_0)$ is Fredholm on $x^\alpha L_b^2(X,E)$.

To analyze the structure of the null space of $\mathsf{D}^2 - r_0^2$, with $r_0 \in \mathbf{R}$, we temporarily write $G(s) := Q_{\mathrm{sm},\alpha}(s)$ and $R(s) := R_{\mathrm{sm},\alpha}(s)$. Then $G(r_0)G(-r_0)$ is our first guess at a parametrix:

$$\begin{aligned}(\mathsf{D}^2 - r_0^2)G(r_0)G(-r_0) &= (\mathsf{D}+r_0)(\mathsf{D}-r_0)G(r_0)G(-r_0) = (\mathsf{D}+r_0)(1+R(r_0))G(-r_0)\\ &= 1 + R(-r_0) + (\mathsf{D}+r_0)R(r_0)G(-r_0)\end{aligned}$$

with $R(-r_0) \in \Psi^{-\infty,\mathcal{E}}(X,E)$, $(\mathsf{D}+r_0)R(r_0)G(-r_0) \in \Psi^{-\infty,\mathcal{F}}(X,E)$ and index sets given by

$$E_{\mathrm{lf}} = \infty, \quad TE_{\mathrm{rf}} = T\Theta_{-\alpha}(-r_0) \quad \text{and} \quad F_{\mathrm{lf}} = \infty,$$

$$TF_{\mathrm{rf}} = T[\Theta_\alpha(-r_0)\overline{\cup}(\Theta_{-\alpha}(r_0) \cup (\Theta_{-\alpha}(r_0) + \Theta_\alpha(-r_0)))\overline{\cup}(\Theta_{-\alpha}(r_0) + \Theta_\alpha(-r_0) + \Theta_{-\alpha}(-r_0))].$$

Taking the limit $\alpha \searrow 0$ as before proves the claim. ■

Note that this Proposition only makes a statement about the *form* of null sections, not about their existence. For instance, since D is selfadjoint on L_b^2, the operator $\mathsf{D}-\lambda_0$ does not have an L_b^2-null space when λ_0 is not purely real. Also, part (d) should only be regarded as an “a priori”-result. We will show below that the top coefficients in $\mathrm{null}_-(\mathsf{D}^2 - r_0^2)$ are much simpler.

Proposition 3.28 implies that the pairing

$$x^{-\varepsilon}L_b^2(X,E) \times x^\varepsilon L_b^2(X,E) \longrightarrow \mathbf{C} \tag{55}$$

induces a pairing between the L_b^2-nullspaces and the extended L_b^2-nullspaces. The relationship between the extended nullspaces $\mathrm{null}_-(\mathsf{D} \pm s)$ and $\mathrm{null}_-(\mathsf{D}^2 - r^2)$ can therefore be described as follows

Lemma 3.29 *The sequence*

$$0 \longrightarrow \mathrm{null}_-(\mathsf{D}-r) \longrightarrow \mathrm{null}_-(\mathsf{D}^2 - r^2) \xrightarrow{\mathsf{D}-r} [\mathrm{null}_{L_b^2}(\mathsf{D}-r)^\perp \cap \mathrm{null}_-(\mathsf{D}+r)] \longrightarrow 0$$

is exact.

Proof. Injectivity of the left map is clear, so is exactness in the middle. We only have to show that the map on the right is (well defined and) surjective. Also, it should be clear that for $\xi \in \mathrm{null}_-(\mathsf{D}^2 - r^2)$, the expression $(\mathsf{D} - r)\xi$ lies in $\mathrm{null}_-(\mathsf{D} + r)$ and pairs to 0 with $\mathrm{null}_{L^2_b}(\mathsf{D} - r)$ under (55). Surjectivity then follows from the fact that

$$\mathrm{null}_{L^2_b}(\mathsf{D} - r)^\perp = \mathrm{null}_{x^\varepsilon L^2_b}(\mathsf{D} - r)^\perp = \overline{\mathrm{im}_{x^{-\varepsilon} L^2_b}(\mathsf{D} - r)} = \mathrm{im}_{x^{-\varepsilon} L^2_b}(\mathsf{D} - r),$$

where we have chosen ε such that $(\mathsf{D} - r)$ is Fredholm on $x^{-\varepsilon} L^2_b(X, E)$. Since $\varepsilon > 0$ with this property can be chosen arbitrarily small, the claim follows. ■

For a more detailed description of the null spaces and the generalized eigenspaces we need to analyze the generalized eigenspaces of the model operator $I_b(\mathsf{D})$, first. Let $P \in \mathrm{Diff}_b(B, \mathcal{K})$ be an $\mathbf{R}_+$-invariant operator (think of $P = I_b(\mathsf{D})$ or $P = I_b(\mathsf{D})^2$). The *formal null space* of P at $z \in \mathrm{spec}_I(P)$ is defined as the extended $x^{-iz} L^2_b(B, \mathcal{K})$-null space of P, i.e.

$$F(P, z) = \{u = \sum_{l=0}^{\mathrm{ord}(z)-1} a_l x^{-iz} \log(x)^l \quad | \quad a_l \in \Gamma(Y, \mathcal{K}), \quad Pu = 0\}.$$

For $z \notin \mathrm{spec}_b(P)$ we have $F(P, z) = \{0\}$. This can be seen, noting that any u with an expansion as above fulfills

$$0 \;=\; P x^{iz} \sum_{l \le p} a_l \log(x)^l = x^{iz} I_b(P)(z) \sum_{l \le p} a_l \log(x)^l + x^{iz} \sum_{l \le p-1} b_l \log(x)^l.$$

Since $I_b(P)(z)$ is invertible, it follows that the coefficient of the highest log-power has to be 0. Thus, the whole expansion must be 0.

In the case of the Dirac operator $P = I_b(\mathsf{D} \mp \lambda)$ the structure of the formal eigenspaces can be inferred from (46) and (47). Let us start with a

Definition 3.30 *Let* $\mathrm{Im}(\lambda) > 0$. *For* ξ_j *in the* c_j*-eigenspace of* D_Y *set*

$$\begin{aligned} \pi^\pm(\xi_j, \lambda) \;&:=\; \sqrt{\lambda + c_j}\xi_j \mp i\sqrt{\lambda - c_j}\boldsymbol{\sigma}\xi_j \\ \pi^\pm(\xi_0, \lambda) \;&:=\; \frac{1}{2}(\xi_0 \pm i\boldsymbol{\sigma}\xi_0) \quad \textit{(the projection onto the } \mp i\textit{-eigenspace of } \boldsymbol{\sigma}\textit{).} \end{aligned}$$

The sections $\pi^\pm$ can be continued to Σ. Recall however that $\theta_j(-r) = \overline{\theta_j(r)}$ for $\lambda = r \in \partial\mathrm{PD} = \mathbf{R}$ and

$$\pi^\pm(\xi_j, -r) = \begin{cases} \pm\,\mathrm{sgn}(r)\boldsymbol{\sigma}\pi^\pm(\xi_j, r) & |r| > c_j \neq 0 \\ \pm\boldsymbol{\sigma}\pi^\mp(\xi_j, r) & |r| \le c_j \neq 0 \end{cases} \tag{56}$$

Now the formal eigenspaces of $\mathsf{D} \pm \lambda$ have the following form

Proposition 3.31 *Let* $\mathrm{Im}(\lambda) \geq 0$.

(a) *For any* c_j *and* $\lambda \neq \pm c_j$*:*

$$\begin{aligned} F(I_b(\mathsf{D} - \lambda), \mp i\theta_j(\lambda)) \;&=\; \left\langle x^{\mp\theta_j(\lambda)} \pi^\mp(\xi_j, \lambda) \,\middle|\, \mathsf{D}_Y \xi_j = c_j \xi_j \right\rangle, \\ F(I_b(\mathsf{D} + \lambda), \mp i\theta_j(\lambda)) \;&=\; \left\langle x^{\mp\theta_j(\lambda)} \boldsymbol{\sigma}\pi^\pm(\xi_j, \lambda) \,\middle|\, \mathsf{D}_Y \xi_j = c_j \xi_j \right\rangle, \end{aligned}$$

Especially, recalling that $\theta_0(\lambda) = -i\lambda$*:*

$$\begin{aligned} F(I_b(\mathsf{D} - \lambda), \pm\lambda \neq 0) &= \langle \pi^{\pm}(\xi_0) x^{\mp i\lambda} \,|\, \mathsf{D}_Y \xi_0 = 0 \rangle, \\ F(I_b(\mathsf{D} + \lambda), \pm\lambda \neq 0) &= \langle \pi^{\pm}(\xi_0) x^{\pm i\lambda} \,|\, \mathsf{D}_Y \xi_0 = 0 \rangle. \end{aligned}$$

(b) *For* $\lambda \in \{\pm c_j\}$ *we get:*

$$\begin{aligned} F(I_b(\mathsf{D} - c_j), 0) &= \langle \xi_j x^0 \log(x) + c_j/2\, \boldsymbol{\sigma} \xi_j x^0, \xi_j x^0 \,|\, \mathsf{D}_Y \xi_j = c_j \xi_j \rangle, \quad j \neq 0 \\ F(I_b(\mathsf{D} + c_j), 0) &= \langle \boldsymbol{\sigma} \xi_j x^0 \log(x) + c_j/2\, \xi_j x^0, \boldsymbol{\sigma} \xi_j x^0 \,|\, \mathsf{D}_Y \xi_j = c_j \xi_j \rangle, \quad j \neq 0 \\ F(I_b(\mathsf{D}), 0) &= \langle \xi_0 x^0 \,|\, \mathsf{D}_Y \xi_0 = 0 \rangle. \end{aligned}$$

(c) *For any* c_j^2 *and* $\lambda^2 \neq c_j^2$*:*

$$F(I_b(\mathsf{D})^2 - \lambda^2), \mp i\theta_j(\lambda)) = \left\langle x^{\mp\theta_j(\lambda)} \eta_j \;\middle|\; \mathsf{D}_Y^2 \eta_j = c_j^2 \eta_j \right\rangle,$$

(d) *For* $\lambda^2 = c_j^2$

$$F(I_b(\mathsf{D})^2 - c_j^2, 0) = \left\langle x^0 \log(x) \eta_j^1 + x^0 \eta_j^0 \,|\, \mathsf{D}_Y^2 \eta_j^i = c_j^2 \eta_j^i \right\rangle$$

■

These spaces encode the “top order behavior” of generalized eigensections of D or D^2. To describe this, introduce the notation $G(P, r) := \sum_{\mathrm{Im}(z) = -r} F(P, z)$. By Lemma 3.18 we know that for $\lambda \in \mathbf{C}$

$$G(\mathsf{D} - \lambda, 0) = \bigoplus_{|c_j| \leq |\lambda|} F(\mathsf{D} - \lambda, \pm\theta_j(\lambda)).$$

It then follows from Lemma 3.14 that the maps

$$\mathrm{null}_-(\mathsf{D} - r_0) \to G(I_b(\mathsf{D}) - r_0, 0), \quad \mathrm{null}_-(\mathsf{D}^2 - r_0^2) \to G(I_b(\mathsf{D})^2 - r_0^2, 0) \tag{57}$$

given by restriction to the top coefficients, are well defined and have kernel contained in $x^{\varepsilon} H_b^{\infty}$. The images of these maps are denoted by $G'(I_b(\mathsf{D}) - r_0, 0)$ and $G'(I_b(\mathsf{D})^2 - r_0^2, 0)$. They can be identified with the quotient between the corresponding extended L_b^2-nullspace and the true L_b^2-nullspace

$$\begin{aligned} G'(I_b(\mathsf{D}) - r_0, 0) &\cong [\mathrm{null}_{L_b^2}(\mathsf{D} - r_0)^{\perp} \subset \mathrm{null}_-(\mathsf{D} - r_0)] \\ G'(I_b(\mathsf{D})^2 - r_0^2, 0) &\cong [\mathrm{null}_{L_b^2}(\mathsf{D}^2 - r_0^2)^{\perp} \subset \mathrm{null}_-(\mathsf{D}^2 - r_0^2)], \end{aligned}$$

where $\perp$ again refers to the pairing (55) above.

Using part (a) of Proposition 3.31, we can now abstractly “construct” generalized eigensections for D, representing elements in $G'(I_b(\mathsf{D}) - r_0, 0)$, as follows: Fixing a cutoff function χ near the boundary we have seen that for $\mathrm{Im}(\lambda) > 0$:

$$\begin{gathered} (\mathsf{D} - \lambda) x^{-\theta_j(\lambda)} \pi^-(\xi_j, \lambda) \chi = O(x^{1 - \theta_j(\lambda)}) \\ (\mathsf{D} + \lambda) x^{-\theta_j(\lambda)} \boldsymbol{\sigma} \pi^+(\xi_j, \lambda) \chi = O(x^{1 - \theta_j(\lambda)}) \end{gathered}$$

For small $\operatorname{Im}(\lambda) > 0$ this is certainly in L^2_b and we can apply the resolvent to it. Thus define

$$\begin{aligned} U^-(\xi_j,\lambda) &:= x^{-\theta_j(\lambda)}\pi^-(\xi_j,\lambda)\chi - (\mathsf{D}-\lambda)^{-1}(\mathsf{D}-\lambda)x^{-\theta_j(\lambda)}\pi^-(\xi_j,\lambda)\chi \qquad (58)\\ U^+(\xi_j,\lambda) &:= x^{-\theta_j(\lambda)}\boldsymbol{\sigma}\pi^+(\xi_j,\lambda)\chi - (\mathsf{D}-\lambda)^{-1}(\mathsf{D}-\lambda)x^{-\theta_j(\lambda)}\boldsymbol{\sigma}\pi^+(\xi_j,\lambda)\chi. \qquad (59)\end{aligned}$$

These sections can be continued meromorphically over Σ. Before noting more of their properties we have to take another look at the resolvent of D.

Let us give a rough description of the real poles of the resolvent:

Proposition 3.32 (Poles of the Resolvent) (a) *Let $\Lambda_0 \in \Sigma$ be located above λ_0 on the real axis, away from any $\pm c_j$ but possibly near 0. Then $G^\mp(\Lambda)$ has a pole of order at most 1 in Λ_0:*

$$G^\mp(\Lambda) = \frac{A^\mp_{\Lambda_0}}{\lambda-\lambda_0} + B^\mp_{\Lambda_0} + (\lambda-\lambda_0)C^\mp_{\Lambda_0}(\Lambda),$$

and $A^\mp_{\Lambda_0}$ is the projection onto the L^2_b-null space of $\mathsf{D}\mp\lambda_0$.

(b) *If λ_0 equals some $\pm c_j \neq 0$ then $G^\mp(\Lambda)$ has a pole of order at most two. Use θ_j as a uniformizing function of Σ around this branching point by writing $\Lambda = \Lambda_0 + \theta_j(\Lambda)^2$. Then*

$$G^\mp(\Lambda) = \frac{A^\mp_{\Lambda_0}}{\theta^2} + \frac{B^\mp_{\Lambda_0}}{\theta} + C^\mp_{\Lambda_0}(\theta).$$

Again, $A^\mp_{\Lambda_0}$ is the projection onto the L^2_b-null space of $\mathsf{D}\mp\lambda_0$.

Proof. To prove (a), note that λ_0 is not a branching point and $G^-(\lambda_0+i\varepsilon) = (\mathsf{D}-(\lambda_0+i\varepsilon))^{-1}$. It follows from the selfadjointness of D and the spectral radius formula that $\|G^-(\lambda_0+i\varepsilon)\| \leq \varepsilon^{-1}$. Thus $G^-(\lambda)$ can have a pole of order at most 1 with a Laurent expansion as in (a). The operators $A^-_{\Lambda_0}, B^-_{\Lambda_0} : x^\varepsilon L^2_b(X,E) \to x^{-\varepsilon}L^2_b(X,E)$, fulfill the equalities

$$(\mathsf{D}-\lambda_0)A^-_{\Lambda_0} = 0, \quad (\mathsf{D}-\lambda_0)B^-_{\Lambda_0} = 1 - A^-_{\Lambda_0}, \quad (\mathsf{D}-\lambda_0)C^-_{\Lambda_0} = -B^-_{\Lambda_0}$$

from the first two of which the claim follows.

The proof of (b) is analogous, when one uses the fact that around a branching point $\pm c_j$ the surface Σ is uniformized by $\theta_j(\Lambda)$. ∎

The main properties of our generalized eigensetions (58), (59) are now listed in the following

Proposition 3.33 (Generalized Eigenfunctions)

(a) $(\mathsf{D}\mp\lambda)U^\mp(\xi_j,\Lambda) = 0$

(b) $U^\mp(\xi_j,\Lambda)$ *is regular near* $\Lambda = 0$

(c) $U^\mp(\xi_j,0) \perp \operatorname{null}_{L^2_b}(\mathsf{D})$.

(d) $U^-(\xi_j,\Lambda) = x^{-\theta_j(\Lambda)}\pi^-(\xi_j,\Lambda) + \sum_{c_k\le|\lambda|} x^{\theta_k(\Lambda)} A_{kj}(\Lambda)\pi^-(\xi_j,\Lambda) + O(x^\varepsilon)$

(e) $U^+(\xi_j,\Lambda) = x^{-\theta_j(\lambda)}\boldsymbol{\sigma}\pi^+(\xi_j,\lambda) + \sum_{c_k\le|\lambda|} x^{\theta_k(\Lambda)} B_{kj}(\Lambda)\boldsymbol{\sigma}\pi^+(\xi_j,\Lambda) + O(x^\varepsilon)$

close to $\partial\mathrm{PD}$ *and where the families of endomorphisms*

$$A_{kj}(\Lambda) : \langle \pi^-(\xi_j,\Lambda)\rangle \to \langle \pi^+(\xi_k,\Lambda)\rangle, \quad B_{kj}(\Lambda) : \langle \boldsymbol{\sigma}\pi^+(\xi_j,\Lambda)\rangle \to \langle \boldsymbol{\sigma}\pi^-(\xi_k,\Lambda)\rangle$$

are meromorphic in that area.

Proof. (a) should be clear, (b) follows from Proposition 3.32, since $(\mathsf{D}-\lambda)x^{-\theta_j(\lambda)}\pi^-(\xi_j,\lambda)\chi$ is orthogonal to the L^2_b-null space of $\mathsf{D}-\lambda$.

Also (d) and (e) follow from Proposition 3.31, since $U^\pm$ certainly belong to the formal eigenspaces to first order and the sums on the RHS are just the most general linear combination of formal eigensections, which are L^2_b for Λ in the physical domain. ■

Restricting to $r \in \partial\mathrm{PD}$ we can rewrite (d) and (e) in the above Proposition in the following way. First,

$$\Pi_r := E_{]-|r|,|r|[}(\mathsf{D}_Y) = \Pi_r^+ \oplus \Pi_r^-, \quad \text{with} \quad \Pi_r^\pm := \langle \pi^\pm(\xi_l,r) \mid c_l < |r|\rangle.$$

Then we can write the endomorphisms in (d) and (e) as:

$$A(r) := \bigoplus_{c_j,c_k<|r|} A_{kj}(r) : \Pi_r^- \longrightarrow \Pi_r^+, \qquad B(-r) := \bigoplus_{c_j,c_k<|r|} B_{kj}(-r) : \Pi_r^+ \longrightarrow \Pi_r^-,$$

and, introducing the definition $\mathbb{X}(r)(\pi^\pm(\xi_j,r)) := \pi^\pm(\xi_j,r)x^{\pm\theta_j(r)}$, and using (56), the generalized eigensections of D in (d) and (e) can now be written

$$U^-(\xi_j,r) = \mathbb{X}(r)(1+A(r))\pi^-(\xi_j,r), \quad U^+(\xi_j,-r) = \mathbb{X}(r)(1+B(-r))\pi^+(\xi_j,r), \tag{60}$$

modulo $O(x^\varepsilon)$. More properties of the “scattering matrices” A and B will be described in Proposition 3.38 below.

Choosing an orthonormal basis of eigenvectors ξ_j^a of the c_j-eigenspace of D_Y (with the additional condition that $\boldsymbol{\sigma}\xi_0^a = i\xi_0^a$) we get an independent set of sections

$$U^-(\xi_0^1,\Lambda),\dots,U^-(\xi_0^{a_0},\Lambda),\dots,U^-(\xi_j^1,\Lambda),\dots,U^-(\xi_j^{a_j},\Lambda) \quad c_j \le |\lambda| \tag{61}$$

in $\mathrm{null}_-(\mathsf{D}-\lambda)/\mathrm{null}_{L^2_b}(\mathsf{D}-\lambda)$. The set of generalized eigensections (61) spans a subspace of dimension $\dim(\Pi_{|\lambda|})/2$. To see that this span is the whole space, we need a second way to obtain information about the dimension.

What is needed here is a scalar product on the spaces of formal eigensections of D. To define this, let P be in $\mathrm{Diff}_b(B,\mathcal{K})$ as before. The *boundary pairing* for P is defined as a the sesquilinear pairing

$$b_P : F(P,z)\times F(P^*,\overline{w}) \longrightarrow \mathbf{C}, \quad b_P(\xi,\eta) := i\int_B \langle P(\chi\xi),\chi\eta\rangle_E - \langle \chi\xi, P^*(\chi\eta)\rangle_E \, dvol_b,$$

where χ is a cutoff function supported near $x=0$ and $\langle\cdot,\cdot\rangle_E$ is antilinear in the *second* variable; b_P is independent of the choice of χ. It is nondegenerate iff $z=w$ and 0 otherwise (see [Me3], Chapter 6). The boundary pairing for the Dirac operator has the following fundamental properties:

Lemma 3.34 *Let* $u \in F(I_b(\mathsf{D}-\lambda), a)$, $v \in F(I_b(\mathsf{D}-\lambda), b)$ *be elements of the general form* $u = x^{-ia}(u_0 + u_1 \log(x))$, $v = x^{-ib}(v_0 + v_1 \log(x))$ *with extensions to* X *denoted by* U, V. *Then for real* λ

$$b_{(\mathsf{D}-\lambda)}(u,v) = i\int_X [\langle \mathsf{D}U, V\rangle_E - \langle U, \mathsf{D}V\rangle_E]\, dvol_{X,b} = \operatorname*{res}_{z=0} i\int_X x^z \langle U, \boldsymbol{\sigma} V\rangle\, dvol_{X,b},$$

which equals $i\langle u_0, \boldsymbol{\sigma} v_0\rangle_{\partial X}$ *when* $a = \overline{b}$.

Proof. This is a simple consequence of "Green's formula", which looks like

$$[\langle \mathsf{D}U, V\rangle_E - \langle U, \mathsf{D}V\rangle_E]\, dvol_b = [\, d^{*,d} x^{-v} \langle U, \mathbf{c}_d(\cdot) V\rangle_E]\, dvol_d$$

in our context. ■

The boundary pairing can be used to obtain relations between the formal and true null spaces of $\mathsf{D}-\lambda$. One can show (as in Chapter 6 in [Me3]):

Proposition 3.35 $G'(\mathsf{D}-\lambda, 0)^{\perp, b_{\mathsf{D}}} = G'(\mathsf{D}-\overline{\lambda}, 0)$, *especially* $G'(\mathsf{D}-r, 0)$, *with* r *real, is a Lagrangian subspace for the symplectic form* b_{D}. ■

Corollary 3.36 *The set (61) is a basis in* $\mathrm{null}_-(\mathsf{D}-r)/\mathrm{null}_{L^2_b}(\mathsf{D}-r)$. ■

Proposition 3.35 can also be used to obtain more information about the scattering operators $A(r)$, $B(r)$ in (60). We start with a simple

Lemma 3.37 *Let* ξ_j, η_k *be eigensections of* D_Y *with eigenvalues* c_j, c_k *and* $0 \neq c_j, c_k < |r|$. *Then w.r.t. the* L^2*-scalar product over* Y

(a) $\langle \pi^\pm(\xi_j, r), \pi^\pm(\eta_k, r)\rangle_{L^2(Y)} = \langle \xi_j, \eta_k\rangle_{L^2(Y)}(|r+c_j| + |r-c_j|)$

(b) $\langle \pi^\pm(\xi_j, r), \boldsymbol{\sigma}\pi^\pm(\eta_k, r)\rangle_{L^2(Y)} = \mp 2i \langle \xi_j, \eta_k\rangle_{L^2(Y)} \sqrt{r^2 - c_j^2}$ ■

Thus introduce

$$w_0^\pm(r) := \mp 1, \qquad w_j^\pm(r) := \pm 2 \frac{\sqrt{r^2 - c_j^2}}{|r+c_j| + |r-c_j|},$$

and define the scalar product $\langle\!\langle\, ,\, \rangle\!\rangle_r$ on $\Pi_{|r|}$ such that $\Pi_r^+ \perp \Pi_r^-$ and

$$\langle\!\langle \varphi_j^\pm, \eta_k^\pm \rangle\!\rangle_r := i\langle \varphi_j^\pm, \boldsymbol{\sigma}\eta_k^\pm\rangle_{L^2(Y)} = w_j^\pm(r) \langle \varphi_j^\pm, \eta_k^\pm\rangle_{L^2(Y)} \quad \text{for} \quad \varphi_i^\pm, \eta_i^\pm \in \langle \pi^\pm(\xi_i, r)\rangle.$$

This is a well defined scalar product away from $r = \pm c_j$ and

Proposition 3.38 $A(r)$, $B(-r)$ *are unitary w.r.t.* $\langle\!\langle\, ,\, \rangle\!\rangle_r$ *and* $A(r)^* = B(-r)$.

Proof. First, the above Corollary implies that the $U^-(\xi_j, r)$ and the $U^+(\xi_j, -r)$ span the generalized eigenspace of $\mathsf{D}-r$. The boundary values of any element in the generalized eigenspace have a unique decomposition into their $\Pi_r^\pm$-parts, which can be written in two ways according to (60):

$$(1+A(r))B(-r)\pi^+(\xi_j,r) = (1+B(-r))\pi^+(\xi_j,r).$$

This shows that $A(r)$ and $B(-r)$ are inverses of each other!

To prove unitarity, we use Proposition 3.35 and Lemma 3.34. For any $\eta \in \Pi_r^-$, for instance $\eta \in \langle \pi^-(\xi_j,r)\rangle$ for a fixed eigenvalue c_j, we get

$$\begin{aligned} 0 &= b_{\mathsf{D}-r}(\mathbb{X}(r)(1+A(r))\eta, \mathbb{X}(r)(1+A(r))\eta) \\ &= i\langle \eta, \boldsymbol{\sigma}\eta\rangle_{L^2(Y)} + i\langle A(r)\eta, \boldsymbol{\sigma}A(r)\eta\rangle_{L^2(Y)} = \langle\!\langle \eta,\eta\rangle\!\rangle_r + \langle\!\langle A(r)\eta, A(r)\eta\rangle\!\rangle_r \end{aligned}$$

and so on. ■

Spectral Measure around 0

Let $f \in C(\mathbf{R})$ have support in around 0. Then, according to Stone's formula

$$\begin{aligned} f(\mathsf{D})\varphi &= \lim_{\varepsilon\to 0}\frac{1}{2\pi i}\int_{\mathbf{R}} f(r)\left[(\mathsf{D}-r-i\varepsilon)^{-1} - (\mathsf{D}-r+i\varepsilon)^{-1}\right]\varphi\, dr \qquad (62) \\ &= \lim_{\varepsilon\to 0}\frac{1}{2\pi i}\int_{\mathbf{R}} f(r)[G^-(r+i\varepsilon) - G^+(-r+i\varepsilon)]\varphi\, dr \\ &= \frac{1}{2\pi i}\int_{\mathbf{R}} f(r)[G^-(r) - G^+(-r)]\varphi\, dr \end{aligned}$$

The last integrand can have singularities. However, fixing $\varphi \in x^\varepsilon L_b^2$ *orthogonal* to the L_b^2-null space of $(\mathsf{D}-r)$, we get, according to Proposition 3.32

$$[G^-(r) - G^+(-r)]\varphi = [B_r^- - B_{-r}^+]\varphi = (\mathsf{D}-r)[C_r^- - C_{-r}^+]\varphi$$

which shows that the LHS exists and lies in $\text{null}_-(\mathsf{D}-r)$ and is orthogonal to the L_b^2-null space of $(\mathsf{D}-r)$. This allows us to make a general *Ansatz* of the form

$$[G^-(r) - G^+(-r)]\varphi = \sum_{|c_j|,|c_k|<r;a,b} a_{j,k}^{a,b}(r)\langle \varphi, U^-(\xi_j^a, r)\rangle U^-(\xi_k^b, r), \qquad (63)$$

where we have used the orthonormal basis introduced before (61). This is a linear combination of finite rank linear operators, continuous on $x^\varepsilon L_b^2$, and has null space in $x^\varepsilon L_b^2$ of codimension at least $\dim(\Pi_{|r|})$. Thus it will be enough to calculate (63) on a (special) finite set of sections φ as follows:

Lemma 3.39 *Let φ be the test function of the form $\varphi = (\mathsf{D}-r)x^{-\theta_j(r)}\pi^-(\xi_j,r)$. Then*

(a) $[G^-(r) - G^+(-r)]\varphi = U^-(\xi_j, r)$

(b) $\langle \varphi, U^-(\eta_k,r)\rangle = \langle \pi^-(\xi_j,r), \boldsymbol{\sigma}\pi^-(\eta_k,r)\rangle_{L^2(Y)}$, *($= 2i\langle \xi_j, \eta_i\rangle_{L^2(Y)}\sqrt{r^2-c_j^2}$ for $k,j \neq 0$)*

Proof. This proof is a good exercise to test the reader's versality with the concepts introduced in this Section. We give the details for reference.

First, using Lemma 3.18(b) and (56), we know that

$$-\theta_j(r) = \theta_j(-r), \qquad \pi^-(\xi_j, r) = \mathrm{sgn}(r)\boldsymbol{\sigma}\pi^-(\xi_j, -r), \qquad \text{for } |r| > c_j. \tag{64}$$

Therefore φ can be written as the limiting function of a function on PD in two ways:

$$\begin{aligned} \varphi &= \lim_{\mathrm{PD}\ni\lambda\to r} (\mathsf{D}-\lambda)x^{-\theta_j(\lambda)}\pi^-(\xi_j,\lambda) && (65)\\ \varphi &= \lim_{\mathrm{PD}\ni\lambda\to -r} (\mathsf{D}+\lambda)x^{\theta_j(\lambda)}\,\mathrm{sgn}(r)\boldsymbol{\sigma}\;\pi^-(\xi_j,\lambda). && (66)\end{aligned}$$

Note that in (66) the Dirac operator is applied to a function which is already in L^2_b! Thus

$$\begin{aligned} G^+(-r)\varphi &= \lim_{\mathrm{PD}\ni\lambda\to -r} (\mathsf{D}+\lambda)(\mathsf{D}+\lambda)x^{\theta_j(\lambda)}\,\mathrm{sgn}(r)\boldsymbol{\sigma}\;\pi^-(\xi_j,\lambda)\\ &= x^{\theta_j(-r)}\,\mathrm{sgn}(r)\boldsymbol{\sigma}\;\pi^-(\xi_j,-r) \overset{(64)}{=} x^{-\theta_j(r)}\pi^-(\xi_j,r). \end{aligned}$$

Recalling the definition (58) of $U^-(\xi_j, r)$, part (a) is proved. To prove (b), simply recall that $U^-(\eta_k, r)$ is in the extended L^2_b-null space of $(\mathsf{D}-r)$. Thus, from the definition of the boundary pairing

$$\langle\varphi, U^-(\eta_k,r)\rangle = -ib_{(\mathsf{D}-r)}(x^{-\theta_j(r)}\pi^-(\xi_j,r), x^{-\theta_k(r)}\pi^-(\eta_k,r)) = \langle\pi^-(\xi_j,r), \boldsymbol{\sigma}\pi^-(\eta_k,r)\rangle_{L^2(Y)},$$

as claimed. The last equation then follows from Lemma 3.37(b). ■

Plugging this into our Ansatz (63) we get

Theorem 3.40 *The continuous part of the spectral measure for* D *is of the form*

$$\begin{aligned} dE_{\mathsf{D},c}(r)\varphi &= \frac{1}{2\pi i}[G^-(r) - G^+(-r)]\varphi\, dr\\ &= \frac{1}{2\pi}\sum_a \langle\varphi, U^-(\xi_0^a,r)\rangle U^-(\xi_0^a,r)\,dr - \frac{1}{4\pi}\sum_{0\neq|c_j|<r;a}(r^2-c_j^2)_+^{-1/2}\langle\varphi, U^-(\xi_j^a,r)\rangle U^-(\xi_j^a,r)\,dr \end{aligned}$$

■

Regularized Trace of the Heat Kernel for Large Times and the Index

Using the spectral measure dE_{D}, the heat kernel for D^2 can be written

$$e^{-t\mathsf{D}^2} = \int_{\mathbf{R}} e^{-tr^2}\,dE_{\mathsf{D}}(r).$$

Choose a cut off function χ, which is 1 near 0 and whose support lies in $]-c_1, c_1[$. Then, using Theorem 3.40, we can calculate the regularized graded trace:

$$\begin{aligned} \mathrm{Str}_d(e^{-t\mathsf{D}^2}\chi(\mathsf{D})) &= \operatorname*{reg}_{z=0} \mathrm{Str}(x^z e^{t\mathsf{D}^2}\chi(\mathsf{D}))\\ &= \operatorname*{reg}_{z=0} \mathrm{Str}(x^z[P_{\mathsf{D}}(0) + \sum_a \frac{1}{2\pi}\int_{\mathbf{R}} e^{-tr^2}\chi(r)U^-(\xi_0^a,r)\langle\cdot, U^-(\xi_0^a,r)\rangle_E\,dr])\\ &= \mathrm{ind}_{L^2_b}(\mathsf{D}) + \operatorname*{reg}_{z=0}\sum_a \frac{1}{2\pi}\int_X x^z\int_{\mathbf{R}} e^{-tr^2}\chi(r)\;\mathrm{str}_E(U^-(\xi_0^a,r)\langle\cdot, U^-(\xi_0^a,r)\rangle_E)\,dr\,dvol_b \end{aligned}$$

Now, using the fact that the grading operator ϵ intertwines the $\pm i$-eigenspaces of $\boldsymbol{\sigma}$, the supertrace can be written as

$$\begin{aligned}\mathrm{str}_E(U^-(\xi_0^a,r)\langle\cdot,U^-(\xi_0^a,r)\rangle_E) &= \langle\epsilon\,(x^{ir}\xi_0^a+x^{-ir}A(r)\xi_0^a),x^{ir}\xi_0^a+x^{-ir}A(r)\xi_0^a\rangle_E+O_r(x^\delta)\\ &= x^{2ir}\langle\epsilon\,\xi_0^a,A(r)\xi_0^a\rangle_E+x^{-2ir}\langle\epsilon\,A(r)\xi_0^a,\xi_0^a\rangle_E+O_r(x^\delta)\end{aligned}$$

The term $O_r(x^\delta)$ is regular in $r=0$. Thus the corresponding part of the above integral vanishes for $t\to\infty$. The remaining part gives integrals of the type

$$\operatorname*{reg}_{z=0}\frac{1}{2\pi}\int_{\mathbf{R}_+}x^z\int_{\mathbf{R}}e^{-tr^2}\chi(r)x^{2ir}\langle\epsilon\,\xi_0^a,A(r)\xi_0^a\rangle_{\partial X}\chi(x)\,dr\frac{dx}{x}=\frac{1}{2}\langle\epsilon\,\xi_0^a,A(0)\xi_0^a\rangle_{\partial X},$$

since this is just the Mellin transform back and forward and evaluated in $z=0$. Thus, recalling that the $U(\xi_0^k,0)/\sqrt{2}$ are orthonormal in $\mathrm{null}(\mathsf{D}_Y)$, we finally obtain

$$\lim_{t\to\infty}\mathrm{Str}_d(e^{-t\mathsf{D}^2})=\mathrm{ind}_{L^2_b}(\mathsf{D})+\frac{1}{2}\sum_a\langle\epsilon\,U(\xi_0^a,0),U(\xi_0^a,0)\rangle_{\partial X}=\mathrm{sdim}(\mathrm{null}_-(\mathsf{D})),\qquad(67)$$

the extended L^2_b-index of D, which we also denote by $\mathrm{ind}_-(\mathsf{D})$. To calculate this index we will have to consider the limit $t\to 0$ of the heat supertrace. This problem will be attacked in the next two Chapters.

4 The Heat Kernel at Finite Time

In this Chapter we describe the construction of the heat kernel of D^2 for finite times. We introduce a heat space X^2_H and heat calculus adapted to our problem and show by direct construction that the heat kernel lies in that calculus. As in the case of the resolvent, this construction will be performed by first solving the heat equation for D^2 to first order (at the relevant faces in X^2_H) and then using an iteration argument, based on a composition formula for our calculus. The presentation given here leans on [DaMe].

4.1 d-Heat Space and d-Heat Calculus

In this Section we describe the heat space and the corresponding calculus adapted to our Dirac operator. We define the heat space as the parabolic blow up

$$X^2_H := \left[X^2_b \times [0,\infty[_t ; (F_\phi \times \{0\}),\ dt; \triangle_X \times \{0\},\ dt\right].$$

The invariant notion of parabolic blow up is explained in [Me0], [Me3] and [DaMe]. For our purposes here, let us just mention that local descriptions of the space X^2_H are given by the sets of coordinates (68, 69, 70, 71) below. See also Remark 4.3.

Now, the corresponding blow down map is denoted by $\beta_H : X^2_H \to X^2 \times [0,\infty[$ and the left and right projections by $\beta_{H,L} : X^2_H \to X$ and $\beta_{H,R} : X^2_H \to X$. The faces of X^2_H are denoted by rf, lf and

$$\begin{aligned} \mathrm{hbf} &:= \overline{\beta_H^{-1}(\partial X^2 \times]0,\infty[)} & \qquad \mathrm{tff} &:= \beta_H^{-1}(F_\phi \times \{0\}) \\ \mathrm{tf} &:= \beta_H^{-1}(\triangle_X \times \{0\}) & \qquad \mathrm{tb} &:= \overline{\beta_H^{-1}((X^2 - \triangle_X) \times \{0\})}. \end{aligned}$$



Figure 5: X^2_H

Using the coordinates $x, x', \mathbf{y}, \mathbf{y}', \mathbf{z}, \mathbf{z}', t$ on $X^2 \times [0,\infty[$ we can describe the properties of lifted vector fields under the blow down map β_H and near the above faces as follows: In the interior of **hbf** we can use projective coordinates w.r.t. x', where x is the boundary defining function in X, as usual:

$$s = \frac{x}{x'}, \quad x', \quad \mathbf{y}, \quad \mathbf{y}', \quad \mathbf{z}, \quad \mathbf{z}', \quad t. \tag{68}$$

Then

$$ds = \frac{dx}{x'} - \frac{1}{sx'}\,dx' \qquad \text{and} \quad \beta_H^* x\frac{\partial}{\partial x} = s\frac{\partial}{\partial s}.$$

Near **hbf** and **tff** we have to consider the dt-parabolic blow up of $F_\phi \times \{0\}$. We do this by introducing coordinates

$$S_t = \frac{s-1}{\sqrt{t}}, \quad X_t = \frac{x'}{\sqrt{t}}, \quad \mathbf{Y}_t = \frac{\mathbf{y}-\mathbf{y}'}{\sqrt{t}}, \quad \sqrt{t}, \quad \mathbf{y}', \quad \mathbf{z}, \quad \mathbf{z}', \tag{69}$$

where now $\mathbf{y}$, $\mathbf{y}'$ are assumed to be **paired**. Then

$$dS_t = \frac{ds}{\sqrt{t}} - \frac{s-1}{2t^{3/2}}\,dt, \quad dX_t = \frac{dx'}{\sqrt{t}} - \frac{x'}{2t^{3/2}}\,dt, \quad d\mathbf{Y}_t = \frac{d\mathbf{y} - d\mathbf{y}'}{\sqrt{t}} - \frac{\mathbf{y}-\mathbf{y}'}{2t^{3/2}}\,dt,$$

and therefore

$$\beta_H^* t\frac{\partial}{\partial t} = \frac{1}{2}\left(\sqrt{t}\frac{\partial}{\partial\sqrt{t}} - S_t\frac{\partial}{\partial S_t} - X_t\frac{\partial}{\partial X_t} - \mathbf{Y}_t\frac{\partial}{\partial \mathbf{Y}_t}\right),$$

$$\beta_H^* x\frac{\partial}{\partial x} = (S_t + \frac{1}{\sqrt{t}})\frac{\partial}{\partial S_t}, \quad \beta_H^*\frac{\partial}{\partial \mathbf{y}} = \frac{1}{\sqrt{t}}\frac{\partial}{\partial \mathbf{Y}_t}, \quad \frac{1}{x}\frac{\partial}{\partial \mathbf{z}} = \frac{1}{sX_t\sqrt{t}}\frac{\partial}{\partial \mathbf{z}}.$$

In the interior of **tff** we can use projective coordinates w.r.t. x':

$$S_x = \frac{s-1}{x'}, \quad \mathbf{Y}_x = \frac{\mathbf{y}-\mathbf{y}'}{x'}, \quad x', \quad \tau = \frac{t}{(x')^2}, \quad \mathbf{y}', \quad \mathbf{z}, \quad \mathbf{z}'. \tag{70}$$

We then have

$$dS_x = \frac{ds}{x'} - \frac{s-1}{(x')^2}\,dx', \quad d\mathbf{Y}_x = \frac{d\mathbf{y} - d\mathbf{y}'}{x'} - \frac{\mathbf{y}-\mathbf{y}'}{(x')^2}\,dx', \quad d\tau = \frac{dt}{(x')^2} - \frac{t}{2(x')^3}\,dx',$$

and

$$\beta_H^* t\frac{\partial}{\partial t} = \tau\frac{\partial}{\partial \tau}, \quad \beta_H^*\frac{\partial}{\partial \mathbf{y}} = \frac{1}{x'}\frac{\partial}{\partial \mathbf{Y}_x}, \quad \beta_H^*\frac{1}{x}\frac{\partial}{\partial \mathbf{z}} = \frac{1}{x}\frac{\partial}{\partial \mathbf{z}}, \quad \beta_H^* x\frac{\partial}{\partial x} = (S_x + \frac{1}{x'})\frac{\partial}{\partial S_x}$$

At **tf** and near **tff** (i.e. after parabolically blowing up $\triangle_X$) we have projective coordinates w.r.t. $\tau^{1/2}$

$$\mathsf{S} = \frac{S_x}{\tau^{1/2}} = S_t, \quad \mathsf{Y} = \frac{\mathbf{Y}_x}{\tau^{1/2}} = \mathbf{Y}_t, \quad \mathsf{Z} = \frac{\mathbf{z}-\mathbf{z}'}{\tau^{1/2}} = \frac{x'(\mathbf{z}-\mathbf{z}')}{t^{1/2}}, \quad \sqrt{\tau}, \quad x', \quad \mathbf{y}', \quad \mathbf{z}'. \tag{71}$$

Here, we have added the requirement that $\mathbf{z}$, $\mathbf{z}'$ also be **paired**. Now

$$d\mathsf{S} = \frac{ds}{\sqrt{t}} - \frac{1}{2t^{3/2}}(s-1)\,dt, \quad d\mathsf{Y}_t = \frac{d\mathbf{y} - d\mathbf{y}'}{\sqrt{t}} - \frac{1}{2t^{3/2}}(\mathbf{y}-\mathbf{y}')\,dt,$$

$$d\mathsf{Z} = \frac{\mathbf{z}-\mathbf{z}'}{\sqrt{t}}\,dx' - \frac{x'(\mathbf{z}-\mathbf{z}')}{2t^{3/2}}\,dt + \frac{x'}{\sqrt{t}}(\,d\mathbf{z} - \,d\mathbf{z}'), \qquad d\sqrt{\tau} = \frac{1}{2x'\sqrt{t}}\,dt + \frac{\sqrt{t}}{2(x')^3}\,dx',$$

and the standard vector fields lift as

$$\beta_H^* x\frac{\partial}{\partial x} = (\mathsf{S} + \frac{1}{\sqrt{t}})\frac{\partial}{\partial \mathsf{S}}, \quad \beta_H^*\frac{\partial}{\partial \mathbf{y}} = \frac{1}{\sqrt{t}}\frac{\partial}{\partial \mathsf{Y}}, \quad \beta_H^*\frac{1}{x}\frac{\partial}{\partial \mathbf{z}} = \frac{1}{s\sqrt{t}}\frac{\partial}{\partial \mathsf{Z}},$$

$$\beta_H^* t\frac{\partial}{\partial t} = \frac{1}{2}\left(\tau^{1/2}\frac{\partial}{\partial \tau^{1/2}} - \mathsf{Z}\frac{\partial}{\partial \mathsf{Z}} - \mathsf{S}\frac{\partial}{\partial \mathsf{S}} - \mathsf{Y}\frac{\partial}{\partial \mathsf{Y}}\right).$$

These calculations show that for $V \in \Gamma({}^\phi TX)$ the vector field $\frac{t^{1/2}}{x}\beta_{H,L}^* V$ is tangent to (the fibres under $\beta_{H,R}$ of) tff and tf. We thus get the following analogue of Lemma 2.1:

Lemma 4.1 *The faces of X_H^2 are isomorphic to standard spaces:*

(a) $\overset{\circ}{\mathrm{tf}} \cong {}^\phi TX$
$\mathrm{tf} \cong \mathrm{RC}^2({}^\phi TX)$

(b) $\overset{\circ}{\mathrm{tff}} \cong {}^\phi N\partial X \times_Y \partial X \times]0,\infty[_\tau$
$\mathrm{tff} \cong \left[\mathrm{RC}^2({}^\phi N\partial X \times_Y \partial X) \times [0,\infty[_\tau ; \triangle_{\partial X} \times \{0\},\ d\tau\right]$

(c) $\overset{\circ}{\mathrm{hbf}} \cong (\partial X)^2 \times]0,\infty[_s \times]0,\infty[_t$
$\mathrm{hbf} \cong [(\partial X)^2 \times [-1,1]_\sigma \times [0,\infty[_t ; F_\phi \times \{0\},\ dt]$ ■

Here, given a euclidean vectorspace (W,g), we denote by $\mathrm{RC}^2(W)$ the "quadratic" radial compactification of W, obtained by introducing $\rho = |\cdot|^{-2}$ as a boundary defining function at infinity. The following gives a description of the restriction of ϕ-vector fields to these faces:

Lemma 4.2 *The lifts of vector fields $V \in \Gamma({}^\phi TX)$, $t\frac{\partial}{\partial t}$, restricted to the different faces are as follows:*

(a) *At* tf : $\beta_{H,L}^*\, t\frac{\partial}{\partial t}|_{\mathrm{tf}} = -\frac{1}{2}R^\phi \in T^\phi TX/X, \qquad \beta_{H,L}^*\, \frac{t^{1/2}}{x}V|_{\mathrm{tf}} = V \in T^\phi TX/X.$

(b) *At* $\overset{\circ}{\mathrm{tff}}$: $\beta_{H,L}^*\, t\frac{\partial}{\partial t}|_{\mathrm{tff}} = \tau\frac{\partial}{\partial \tau}, \quad \beta_{H,L}^*\, \frac{t^{1/2}}{x}V|_{\mathrm{tff}} = \tau^{1/2}N_\phi(V) \in T({}^\phi N\partial X \times_Y \partial X/Y).$ ■

Now define the density and the coefficient bundle for the heat kernel as

$$\mathrm{KD}_H := \frac{(x')^n}{t^{n/2}}\frac{dt}{t}\beta_{H,R}^*{}^\phi\Omega(X), \quad \mathrm{CB}_H := \mathrm{KD}_H \otimes \beta_{H,L}^* E \otimes \beta_{H,R}^* E^* =: \mathrm{KD}_H \otimes \mathrm{END}(E).$$

Since we have defined the density KD_H to be lifted from the right, its Lie derivative w.r.t. a vector field lifted from the left is trivial:

$$L_{\beta_{H,L}^* V}\mathrm{KD}_H = 0, \qquad L_{\beta_H^* t\frac{\partial}{\partial t}}\mathrm{KD}_H = -\frac{n}{2}\mathrm{KD}_H.$$

This will again prove useful in Section 4.3, when we calculate the action of such vector fields on our calculus.

The *d*-heat calculus is defined to be

$$\Psi^{l,m,p}_{H,\mathrm{cl}}(X,E) := \rho_{\mathrm{tf}}^{l}\,\rho_{\mathrm{tff}}^{m}\,\rho_{\mathrm{hbf}}^{p-v}\,\dot{C}^{\infty}_{\mathrm{tf,tff,hbf}}(X^2_H,\mathrm{CB}_H)$$

for $l > 0, m \geq 0, p \geq 0$. For $l = 0$, we set

$$\Psi^{0,m,p}_{H,\mathrm{cl}}(X,E) := \rho_{\mathrm{tff}}^{m}\,\rho_{\mathrm{hbf}}^{p-v}\,\dot{C}^{\infty}_{\mathrm{tf,tff,hbf}}(X^2_H,\mathrm{CB}_H)^{\mathrm{tf}} \oplus \mathbf{C},$$

where, using (79), we require for $A \in \dot{C}^{\infty}_{\mathrm{tf,tff,hbf}}(X^2_H,\mathrm{CB}_H)^{\mathrm{tf}}$ that the following mean value condition holds

$$\int_{{}^{\phi}TX/X} \boldsymbol{\iota}(t\frac{\partial}{\partial t})A|_{\mathrm{tf}} = 0. \tag{72}$$

Let us show, how we can define the action of an element A in the calculus on some function $g \in \dot{C}^{\infty}_{c}(X \times [0,\infty[, E)$. First, for any function $f \in \dot{C}^{\infty}_{c}(X \times [0,\infty[, \Omega(X)\, dt \otimes E^*)$ we consider the convolution product

$$f * g(t) = \int_0^\infty f(t+s)g(s) \in \dot{C}^{\infty}_{c}(X \times X \times [0,\infty[, \beta^*_{H,L}\Omega(X) \otimes \beta^*_{H,L}E^* \otimes \beta^*_{H,R}E).$$

Note that for functions vanishing rapidly at the faces $\mathrm{rf}, \mathrm{lf}, \mathrm{tb}$ we have

$$\beta^*_{H,L}\Omega(X) \otimes \mathrm{KD}_H \sim \rho^{0}_{\mathrm{tff}}\,\rho^{-1}_{\mathrm{tf}}\,\rho^{v}_{\mathrm{hbf}}\Omega(X^2_H),$$

which implies that for an operator $A \in \Psi^{l,m,p}_{H,\mathrm{cl}}(X,E)$ the pairing

$$(A, \beta^*_H(f * g)) \in \rho^{m}_{\mathrm{tff}}\,\rho^{l-1}_{\mathrm{tf}}\,\rho^{p}_{\mathrm{hbf}}\,\dot{C}^{\infty}_{\mathrm{tf,tff,hbf}}(X^2_H, \Omega(X^2_H)),$$

is integrable for $l > 0$. In the case $l = 0$ we have to use the mean value condition (72) to make sense of the integral. This is done by writing, in the neighborhood $U(\mathrm{tf}) := \{\rho_{\mathrm{tf}} < \varepsilon\} \cong [0,\varepsilon[\times {}^{\phi}TX$:

$$\int_{U(\mathrm{tf})} (A, \beta^*_H(f * g)) = \lim_{\delta \to 0} \int_\delta^\varepsilon \int_X \int_{{}^{\phi}TX/X} a\, dvol_{{}^{\phi}TX/X}\, dvol_X\, \frac{d\rho_{\mathrm{tf}}}{\rho_{\mathrm{tf}}}.$$

Since the innermost integral converges to 0 for $\rho_{\mathrm{tf}} \to 0$, this limit exists.

Now we can define the action of A on g by setting

$$\langle Ag, f\rangle_{X \times [0,\infty[} := \langle A, \beta^*_H(f * g)\rangle_{X^2_H}.$$

As usual, the task of writing down the definitions for the extended heat calculus is left to the reader. It is important to note, however, that we can only allow conormal functions for which the above mean value condition makes sense.

4.2 Composition Formula

To explain our constructions in this Section consider simplified convolution operators $A, B \in \dot{C}^\infty_c(X^2 \times [0,\infty[, \Omega_R(X)\, dt)$. Their composition is given by

$$A \circ B(\mathbf{w}, \mathbf{w}'', t) = \int_{\mathbf{w}', t'} A(\mathbf{w}, \mathbf{w}', t - t') B(\mathbf{w}', \mathbf{w}'', t') \in \dot{C}^\infty_c(X^2 \times [0,\infty[, \Omega_R(X)\, dt). \tag{73}$$

Using the space $X^3 \times [0,\infty[_{t'}[\times[0,\infty[_{t-t'}$ and the three projections

$$\begin{tikzcd} X^2 \times [0,\infty[_{t-t'} & X^3 \times [0,\infty[_{t'}[\times[0,\infty[_{t-t'} \arrow[l, "\pi_L"'] \arrow[r, "\pi_R"] \arrow[d, "\pi_M"] & X^2 \times [0,\infty[_{t'} \\ & X^2 \times [0,\infty[_t & \end{tikzcd}$$

this can also be written as:

$$A \circ B = \pi_{M,*}(\pi_L^* A \cdot \pi_R^* B).$$

Now, in our case, the image spaces of the projections will really have to be X^2_H. To be able to use the pullback- and pushforward-results in [Me0] (see also Appendix A.6) we will have to modify the "triple space" such that the corresponding lifted projections become b-fibrations.

Remark 4.3 To simplify things a bit, we agree to define the d-heat space X^2_H using $\sqrt{t}$ as a *global* variable, i.e. as the simple blow up

$$X^2_H = \left[X^2_b \times [0,\infty[_{\sqrt{t}}; (F_\phi \times \{0\}); \triangle_X \times \{0\}\right] \quad \text{away from tb.}$$

Since elements in the heat calculus vanish at tb by definition, nothing is lost by considering the RHS above, instead of the original space X^2_H.

We now define

$$\mathsf{T}^2 := [0,\infty[_{\sqrt{t'}} \times [0,\infty[_{\sqrt{t-t'}},$$

and denote its faces by lt and rt. The first thing to note now, is that the map

$$\pi_M : \mathsf{T}^2 \longrightarrow [0,\infty[_{\sqrt{t}} \quad (a,b) \mapsto \sqrt{a^2 + b^2}$$

is *not* a b-fibration. Thus we will have to take care to also blow up the corner bt $=$ lt $\cap$ rt whenever this situation arises. So our starting point is the diagram

$$\begin{tikzcd} X^2_b \times [0,\infty[_{\sqrt{t-t'}} & X^3_b \times \mathsf{T}^2 \arrow[l, "\pi_L"'] \arrow[r, "\pi_R"] \arrow[d, "\pi_M"] & X^2_b \times [0,\infty[_{\sqrt{t'}} \\ & X^2_b \times [0,\infty[_{\sqrt{t}} & \end{tikzcd} \tag{74}$$

where we keep in mind that the horizontal maps are still b-fibrations but π_M is not.

The definition of the heat triple space now proceeds in two steps. The first step consists in replacing the image spaces in (74) by the partial blow up $\widetilde{X}^2_H = [X^2_b \times [0,\infty[; F_\phi \times \{0\}, dt]$, which we interpret as simple blow up w.r.t. the global variable $\sqrt{t}$.



Figure 7: The T^2- and X^3_b-factor

As in the case of the ϕ-triple space we denote the faces to be blown up (as well as the resulting faces) by

$$\mathrm{bt} \times \phi_C,\ \mathrm{bt} \times \phi_{CT},\ \mathrm{bt} \times \phi_{TT},\ \mathrm{rt} \times \phi_F,\ \mathrm{lt} \times \phi_S,\ \mathrm{rt} \times \phi_{FT},\ \mathrm{lt} \times \phi_{ST},$$

and define the (intermediate) heat triple space to be

$$\widetilde{X}^3_H := \left[X^3_b \times \mathsf{T}^2; \mathrm{bt}\phi_{TT}; \mathrm{bt}\phi_{CT}; \mathrm{rt}\phi_{FT}; \mathrm{lt}\phi_{ST}; \mathrm{bt}\phi_C; \mathrm{rt}\phi_F; \mathrm{lt}\phi_S\right].$$

To avoid any misunderstandings, recall that this is the space $X^3_b \times \mathsf{T}^2$, with faces blown up as listed from left to right. Each face in that list thus really represents the *preimage* of the corresponding face in $X^3_b \times \mathsf{T}^2$ under the previous blow ups.

For the second step, denote the different lifts of the diagonal $\triangle_\phi \subset X^2_\phi$ to X^3_ϕ by $\triangle_L$, $\triangle_M$, $\triangle_R$ and analogously their intersection by $\triangle_{LMR}$. Then the true "triple space" is defined as

$$X^3_H := [\widetilde{X}^3_H; \mathrm{bt}\triangle_{LRM}; \mathrm{bt}\triangle_M; \mathrm{lt}\triangle_L, \mathrm{rt}\triangle_R].$$

Besides those faces obtained by the above blow ups, this space has the faces

$$M \overset{\triangle}{=} \mathsf{T}^2 M, L, R, S, C, T, \quad \text{and} \quad \mathrm{bt} \overset{\triangle}{=} \mathrm{bt}X^3_b, \mathrm{rt}, \mathrm{lt}.$$

Again, these are understood to be the lifts of the corresponding faces in $X^3_b \times \mathsf{T}^2$ minus those faces, which were obtained in the previous blow up. The important result now is

Lemma 4.4 *There are induced b-maps $X^3_H \xrightarrow{\pi_{H,o}} X^2_H$, $o = L, M, R$, (where we agreed to define the RHS to be defined with global time variable $\sqrt{t}$) such that the diagrams*

$$\begin{tikzcd}
X^3_H \arrow[r, "\pi_{H,o}"] \arrow[d] & X^2_H \arrow[d] \\
X^3_b \times \mathsf{T}^2 \arrow[r, "\pi_o"] & X^2_b \times \mathsf{T}
\end{tikzcd}$$

commute.

Proof. We show how the map $\pi_{H,L}$ is defined. Starting with X^3_H we first blow down all those faces which do not intersect any of the faces lifted from the left factor:

$$\mathrm{rt}\triangle_R,\ \mathrm{bt}\triangle_M,\ \mathrm{rt}\phi_F,\ \mathrm{rt}\phi_{FT},\ \mathrm{bt}\phi_C,\ \mathrm{bt}\phi_{CT},\ F,\ C,$$

thus getting

$$[X^2_b \times X \times \mathsf{T}^2; \mathrm{bf} \times \partial X \times \mathsf{T}^2; \mathrm{bt}\phi_{TT}; \mathrm{lt}\phi_{ST}; \mathrm{lt}\phi_S; \mathrm{bt}\triangle_{LRM}; \mathrm{lt}\triangle_L].$$

Now, obviously, $\mathrm{bt}\triangle_{LRM}$ is a p-submanifold of $\mathrm{lt}\triangle_L$, i.e. their blow ups can be exchanged and the triple diagonal can be blown down. The same argument let us us exchange the blow ups of $\mathrm{bt}\phi_{TT}$ and $\mathrm{lt}\phi_{ST}$, yielding

$$[X^2_b \times X \times \mathsf{T}^2; \mathrm{bf} \times \partial X \times \mathsf{T}^2; \mathrm{lt}\phi_{ST}; \mathrm{lt}\phi_S; \mathrm{bt}\phi_{TT}; \mathrm{lt}\triangle_L].$$

Now the face $\mathrm{bt}\phi_{TT}$ in question here is really the preimage of this face under the blow up of $\mathrm{lt}\phi_{ST}$, and one easily checks that it is transversal to $\mathrm{lt}\triangle_L$. Being disjoint from the other blow ups, we can switch the corresponding blow up to the front and then blow down $\mathrm{bt}\phi_{TT}$ altogether. Following this, it is easy to see that the blow up of $\mathrm{bf} \times \partial X \times \mathsf{T}^2$ can be "pulled to the front", if one switches the order of the blow ups $\mathrm{lt}\phi_{ST}$ and $\mathrm{lt}\phi_S$. Thus this face can be blown down, too, leaving us with

$$[X^2_b \times X \times \mathsf{T}; F_\phi \times X \times \{0\}; F_\phi \times \partial X \times \{0\}; \triangle_b \times X \times \{0\}] \times \mathsf{T},$$

where we have written the faces ϕ_S, ϕ_{ST} in their blown down form in $X^2_b \times X$. It is important to note that, again, $\phi_{ST} \times \{0\}$ is really the preimage of that face under the blow up of $\phi_S \times \{0\}$, making it transversal to the lifted left diagonal. Thus $\phi_{ST}\{0\}$ can be blown down and we are left with

$$[X^2_b \times \mathsf{T}; F_\phi \times \{0\}; \triangle_b \times \{0\}] \times X \times \mathsf{T},$$

which projects nicely to X^2_H. ■

Since all their entries are either 0 or 1, we can write down the "coefficient matrices" for π_L, π_M, π_R (see Appendix A.6) by just listing the preimages of the different faces of X^2_H:

Lemma 4.5 (a) π_L *maps the faces* $\{L, \mathrm{rt}, \mathrm{rt}\triangle_R\}$ *to the interior of* X^2_H. *Also*

$$\begin{aligned}
\pi_L^*\mathrm{hbf} &= \{D, T, \mathrm{bt}\phi_{CT}, \mathrm{rt}\phi_{FT}\}\\
\pi_L^*\mathrm{tff} &= \{\mathrm{lt}\phi_S, \mathrm{lt}\phi_{ST}, \mathrm{bt}\phi_{TT}\}\\
\pi_L^*\mathrm{tf} &= \{\mathrm{lt}\triangle_L, \mathrm{bt}\triangle_{LMR}\}\\
\pi_L^*\mathrm{lbf} &= \{R, C, \mathrm{bt}\phi_C\}\\
\pi_L^*\mathrm{rbf} &= \{M, F, \mathrm{rt}\phi_F\}\\
\pi_L^*\mathrm{tb} &= \{\mathrm{bt}, \mathrm{lt}, \mathrm{bt}\phi_C, \mathrm{bt}\phi_{CT}, \mathrm{bt}\triangle_M\}
\end{aligned}$$

(b) π_R *maps the faces* $\{R, \mathrm{lt}, \mathrm{lt}\triangle_L\}$ *to the interior of* X^2_H. *Also*

$$\begin{aligned}
\pi_R^*\mathrm{hbf} &= \{F, T, \mathrm{bt}\phi_{CT}, \mathrm{lt}\phi_{ST}\}\\
\pi_R^*\mathrm{tff} &= \{\mathrm{rt}\phi_F, \mathrm{rt}\phi_{FT}, \mathrm{bt}\phi_{TT}\}\\
\pi_R^*\mathrm{tf} &= \{\mathrm{rt}\triangle_R, \mathrm{bt}\triangle_{LMR}\}\\
\pi_R^*\mathrm{lbf} &= \{M, S, \mathrm{lt}\phi_S\}\\
\pi_R^*\mathrm{rbf} &= \{L, C, \mathrm{bt}\phi_C\}\\
\pi_R^*\mathrm{tb} &= \{\mathrm{bt}, \mathrm{rt}, \mathrm{bt}\phi_C, \mathrm{bt}\phi_{CT}, \mathrm{bt}\triangle_M\}
\end{aligned}$$

(c) π_M *maps the faces* $\{S, \mathrm{rt}, \mathrm{lt}, \mathrm{rt}\triangle_R, \mathrm{lt}\triangle_L\}$ *to the interior of* X_H^2. *Also*

$$\begin{aligned}
\pi_M^*\mathrm{hbf} &= \{C, T, \mathrm{rt}\phi_{FT}, \mathrm{lt}\phi_{ST}\}\\
\pi_M^*\mathrm{tff} &= \{\mathrm{bt}\phi_C, \mathrm{bt}\phi_{CT}, \mathrm{bt}\phi_{TT}\}\\
\pi_M^*\mathrm{tf} &= \{\mathrm{bt}\triangle_M, \mathrm{bt}\triangle_{LMR}\}\\
\pi_M^*\mathrm{lbf} &= \{R, M, \mathrm{lt}\phi_S\}\\
\pi_M^*\mathrm{rbf} &= \{L, F, \mathrm{rt}\phi_F\}\\
\pi_M^*\mathrm{tb} &= \{\mathrm{bt}\}
\end{aligned}$$

■

This Lemma shows that π_L, π_R, are not b-normal, and thus cannot be b-fibrations, since the faces $\mathrm{bt}\phi_C$, $\mathrm{bt}\phi_{CT}$ appear in the preimage of several faces of X_H^2. However, for two heat operators

$$A \in \Psi_H^{\mathcal{L}}(X) = \mathcal{A}^{\mathcal{L}}(X_H^2, \rho_{\mathrm{hbf}}^{-v}\mathrm{KD}_H), \quad B \in \Psi_H^{\mathcal{R}}(X) = \mathcal{A}^{\mathcal{R}}(X_H^2, \rho_{\mathrm{hbf}}^{-v}\mathrm{KD}_H)$$

their lift from the left and the right to X_H^3 is

$$\pi_L^*A \cdot \pi_R^*B \in \mathcal{A}^{\mathcal{H}}(X_H^3, \Omega(X_H^3)),$$

and the above Lemma tells us that

$$\begin{aligned}
H_L = H_M = H_R = H_F = H_S = H_C = H_{\mathrm{bt}} = H_{\mathrm{rt}} = H_{\mathrm{lt}}&\\
= H_{\mathrm{bt}\phi_C} = H_{\mathrm{lt}\phi_S} = H_{\mathrm{rt}\phi_F} = H_{\mathrm{bt}\phi_{CT}} = H_{\mathrm{bt}\triangle_M} = \infty.& \qquad (75)
\end{aligned}$$

To describe the expansions at the remaining faces, we need to caculate the lifts of our densities under the blow down map $\gamma_H : X_H^3 \to X^3 \times [0,\infty[^2$. Writing $s = t - t'$ for the (right) variable in T^2 note first the coefficients which appear when lifting to $X_b^3 \times \mathsf{T}^2$:

$$\gamma_b^*\Omega(X^3) \otimes {}^b\Omega(\mathsf{T}^2) = \rho^2_{T\mathsf{T}^2_b}\,\rho_{F\mathsf{T}^2_b}\,\rho_{S\mathsf{T}^2_b}\,\rho_{C\mathsf{T}^2_b}\,\rho^{-1}_{X^3_b\mathrm{rt}}\,\rho^{-1}_{X^3_b\mathrm{lt}}\,\gamma^*_{H,b}\Omega(X^3_b \times \mathsf{T}^2). \qquad (76)$$

To determine the relevant coefficients of the lift of

$$\Omega_L(X)\pi_L^*(\rho_{\mathrm{hbf}}^{-v}\mathrm{KD}_H)\pi_R^*(\rho_{\mathrm{hbf}}^{-v}\mathrm{KD}_H) = (s^{1/2}t'^{1/2})^{-n}(x'x'')^{v-1}\,\pi_L^*\rho_{\mathrm{hbf}}^{-v}\,\pi_R^*\rho_{\mathrm{hbf}}^{-v}\Omega(X^3)^b\Omega(\mathsf{T}^2),$$

we use Lemma 2.2, i.e we need to list the contributions of the factors in (76) at the faces of X_H^3 and to keep track of the codimensions of the faces blown up in $\partial(X_b^3 \times \mathsf{T}^2)$:

face	codim	(76)	$(s^{1/2}t'^{1/2})^{-n}$	$(x'x'')^{v-1}$	$\pi_L^*\rho_{\text{hbf}}^{-v}\pi_R^*\rho_{\text{hbf}}^{-v}$	Total
T	0	2	0	$2v-2$	$-2v$	0
$\text{bt}\phi_{TT}$	$2h+4$	0	$-2n$	$2v-2$		0
$\text{rt}\phi_{FT}$	$h+2$	1	$-n$	$2v-2$	$-v$	0
$\text{lt}\phi_{ST}$	$h+2$	1	$-n$	$2v-2$	$-v$	0
$\text{bt}\triangle_{LMR}$	$2n+1$	-2	$-2n$			-1
$\text{rt}\triangle_R$	n	-1	$-n$			-1
$\text{lt}\triangle_L$	n	-1	$-n$			-1

Thus, at the relevant faces, the density is

$$\begin{aligned}\gamma_b^*\Omega_L(X)(\pi_L^*\rho_{\text{hbf}}^{-v}\mathrm{KD}_H)(\pi_R^*\rho_{\text{hbf}}^{-v}\mathrm{KD}_H)\\ = \rho_T^0\,\rho_{\text{rt}\phi_{FT}}^0\,\rho_{\text{lt}\phi_{ST}}^0\,\rho_{\text{bt}\phi_{TT}}^0\,\rho_{\text{lt}\triangle_L}^{-1}\,\rho_{\text{rt}\triangle_R}^{-1}\,\rho_{\text{bt}\triangle_{LMR}}^{-1}\,\Omega(X_H^3),\end{aligned} \tag{77}$$

and the nontrivial terms of the index set $\mathcal{H}$ are given by

$$\begin{array}{lcllcl} H_T & = & L_{\text{hbf}} + R_{\text{hbf}} & & & \\ H_{\text{rt}\phi_{FT}} & = & L_{\text{hbf}} + R_{\text{tff}} & H_{\text{lt}\phi_{ST}} & = & L_{\text{tff}} + R_{\text{hbf}} \\ H_{\text{bt}\phi_{TT}} & = & L_{\text{tff}} + R_{\text{tff}} & & & \\ H_{\text{lt}\triangle_L} & = & L_{\text{tf}} - 1 & H_{\text{rt}\triangle_R} & = & R_{\text{tf}} - 1 \\ H_{\text{bt}\triangle_{LMR}} & = & L_{\text{tf}} + R_{\text{tf}} - 1 & & & \end{array}$$

Though π_M is not a b-fibration it can be made into one in this special case, because (75) allows the faces causing the nuisance to be blown down. Thus the pushforward

$$\pi_{M,*}(\gamma_{H,L}^* f \pi_L^* A \cdot \pi_R^* B) \in \mathcal{A}^{\pi_{M\sharp}\mathcal{H}}(X_H^2, \Omega(X_H^2))$$

is well defined when it is integrable at those faces mapped to the interior by π_M, i.e. whenever

$$H_{\text{rt}\triangle_R}, H_{\text{lt}\triangle_L} > -1 \quad \text{or equivalently} \quad R_{\text{tf}}, L_{\text{tf}} > 0$$

(the condition at the other faces again is trivial due to (75)), and the pushforward of the index set is given by:

$$\pi_{M\sharp}\mathcal{H}(\text{hbf}) = H_T \overline{\cup} H_{\text{lt}\phi_{ST}} \overline{\cup} H_{\text{rt}\phi_{TF}}, \quad \pi_{M\sharp}\mathcal{H}(\text{tff}) = H_{\text{bt}\phi_{TT}}, \quad \pi_{M\sharp}\mathcal{H}(\text{tf}) = H_{\text{bt}\triangle_{LMR}}.$$

Also, in the case $R_{\text{tf}}, L_{\text{tf}} \geq 0$, the pushforward can still be defined using the mean value condition (72). Rewriting everything in terms of $\beta_{H,L}^*\Omega(X)\rho_{\text{hbf}}^{-v}\mathrm{KD}_H = \rho_{\text{tf}}^{-1}\Omega(X_H^2)$ we finally get

Theorem 4.6 (Composition Formula) *Let $R_{\text{tf}}, L_{\text{tf}} \geq 0$. Then composition gives a map*

$$\Psi_H^{\mathcal{L}}(X) \times \Psi_H^{\mathcal{R}}(X) \xrightarrow{\circ} \Psi_H^{\mathcal{M}}(X)$$

with index sets given by

$$M_{\text{hbf}} = (L_{\text{hbf}} + R_{\text{hbf}}) \overline{\cup} (L_{\text{tff}} + R_{\text{hbf}}) \overline{\cup} (L_{\text{hbf}} + R_{\text{tff}}),$$

$$M_{\text{tff}} = L_{\text{tff}} + R_{\text{tff}}, \quad M_{\text{tf}} = L_{\text{tf}} + R_{\text{tf}}.$$

■

4.3 Normal Operators and Construction of the Heat Kernel

Using the local coordinate representations described in Section 4.1 it is easy to see that the restriction of the density KD_H to the different faces has the following form.

$$\begin{aligned}
\mathrm{KD}_H|_{\overset{\circ}{\mathrm{tf}}} &\cong \frac{dt}{t}\Omega({}^{\phi}TX/X) && (78)\\
\mathrm{KD}_H|_{\overset{\circ}{\mathrm{tff}}} &\cong \frac{d\tau}{\tau}\tau^{-n/2}\Omega_{\mathrm{fib}}({}^{\phi}N\partial X \times_Y \partial X \xrightarrow{\beta_{H,L}} \partial X) && (79)\\
\mathrm{KD}_H|_{\overset{\circ}{\mathrm{hbf}}} &\cong \frac{dt}{t}X_t^{v}\,{}^{b}\Omega_{\mathrm{fib}}(\mathrm{hbf} \overset{\beta_{H,L}}{\to} \partial X) && (80)
\end{aligned}$$

For $A \in \Psi^{l,m,p}_{H,\mathrm{cl}}(X)$ we now define the normal operators at the faces tf and tff and hbf by

$$\begin{aligned}
N_{H,\mathrm{tf},l>0}(A) &:= t^{1-l/2}A|_{\mathrm{tf}} \in \rho_{\mathrm{tff}}^{m-l}\,\dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{tf}, t\mathrm{KD}_H) \cong x^{m-l}\mathcal{S}_{\mathrm{fib}}({}^{\phi}TX, \Omega({}^{\phi}TX/X)\,dt)\\
N_{H,\mathrm{tf},0}(A \oplus c) &:= tA|_{\mathrm{tf}} \oplus c \in \rho_{\mathrm{tff}}^{m}\,\dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{tf}, t\mathrm{KD}_H)^{\mathrm{tf}} \oplus \mathbf{C}\\
N_{H,\mathrm{tff},m}(A) &:= (x')^{-m}A|_{\mathrm{tff}} \in \rho_{\mathrm{tf}}^{l}\,\rho_{\mathrm{hbf}}^{p-v-m}\ \dot{C}^{\infty}_{\mathrm{tf,hbf}}(\mathrm{tff}, \mathrm{KD}_H)\\
N_{H,\mathrm{hbf},p}(A) &:= (x')^{-p}A|_{\mathrm{hbf}} \in \rho_{\mathrm{tff}}^{m-p}\,\dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{hbf}, {}^{b}\Omega_{\mathrm{fib}}(\mathrm{hbf})\,dt)
\end{aligned}$$

For $l \geq 1$ these normal operators fit into the usual short exact sequences:

$$0 \longrightarrow \Psi^{l-1,m,p}_{H,\mathrm{cl}}(X) \longrightarrow \Psi^{l,m,p}_{H,\mathrm{cl}}(X) \overset{N_{H,\mathrm{tf},l}}{\longrightarrow} x^{m-l}\mathcal{S}_{\mathrm{fib}}({}^{\phi}TX, \Omega({}^{\phi}TX/X)\,dt) \longrightarrow 0$$

$$0 \longrightarrow \Psi^{l,m-1,p}_{H,\mathrm{cl}}(X) \longrightarrow \Psi^{l,m,p}_{H,\mathrm{cl}}(X) \overset{N_{H,\mathrm{tff},m}}{\longrightarrow} \rho_{\mathrm{tf}}^{l}\,\rho_{\mathrm{hbf}}^{p-v-m}\ \dot{C}^{\infty}_{\mathrm{tf,hbf}}(\mathrm{tff}, \mathrm{KD}_H) \longrightarrow 0 \qquad (81)$$

$$0 \longrightarrow \Psi^{l,m,p-1}_{H,\mathrm{cl}}(X) \longrightarrow \Psi^{l,m,p}_{H,\mathrm{cl}}(X) \overset{N_{H,\mathrm{hbf},p}}{\longrightarrow} \rho_{\mathrm{tff}}^{m-p}\ \dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{hbf}, {}^{b}\Omega_{\mathrm{fib}}(\mathrm{hbf})\,dt) \longrightarrow 0,$$

and we leave the formulation of the case $l = 0$ to the reader, as well as the usual extension to the calculus with coefficients.

As usual, the spaces on the RHS in the above sequences represent some calculi of operators whose composition rules are induced by the composition rules for operators in the "big" calculus $\Psi_{H,\mathrm{cl}}(X)$ (compare Lemma 3.5). For instance, in the case (81) we can define

$$\Psi^{l,p}_{H,\mathrm{sus}}({}^{\phi}N\partial X) := \rho_{\mathrm{tf}}^{l}\,\rho_{\mathrm{hbf}}^{p-v}\ \dot{C}^{\infty}_{\mathrm{tf,hbf}}(\mathrm{tff}, \mathrm{KD}_H).$$

This is the heat space associated with the suspended calculus introduced in Section 3.3. Some related calculi are described in more detail in Appendices A.4 and A.5.

At this point we are not interested in the general algebraic properties of the normal maps. All we really need is to describe the normal action of the Dirac operator D at the different faces, for which it will suffice – as usual – to analyze compositions of vector fields with elements in the heat calculus. To do this, we remind the reader of the following simple fact about Lie derivatives of forms and densities: Let $j_F : F \hookrightarrow M$ be a face of a manifold with corners with defining function f, W a vector field on M tangent to F, and ω a form on M. Then

$$j_F^* L_{fW}\omega = j_F^*(f L_W \omega + \boldsymbol{\varepsilon}(\,df)\,\boldsymbol{\iota}(W)\omega) = 0,$$

which means that

$$j_F^* L_W \omega = L_{W|_F} j_F^* \omega,$$

i.e. the Lie derivative restricted to the face can be calculated *on* the face. The normal action of vector fields on the elements in the calculus is now described in

Proposition 4.7 *Let V be in $\Gamma({}^\phi TX)$. Composition gives maps*

$$\Psi_{H,\mathrm{cl}}^{l,m,p}(X) \xrightarrow{\frac{1}{x}V\circ} \Psi_{H,\mathrm{cl}}^{l-1,m-1,p-1}(X), \qquad \Psi_{H,\mathrm{cl}}^{l,m,p}(X) \xrightarrow{\frac{\partial}{\partial t}\circ} \Psi_{H,\mathrm{cl}}^{l-2,m-2,p}(X)$$

Applied to $A \in \Psi_{H,\mathrm{cl}}^{l,m,p}(X)$, we get at the different faces

(a) $N_{H,\mathrm{tf},l-2}(\frac{\partial}{\partial t}\circ A) = \begin{cases} -\frac{1}{2}(L_{R^\phi} + 2 - l) N_{H,\mathrm{tf},l}(A) & \text{for} \quad l > 2 \\ -\frac{1}{2}(L_{R^\phi} + 2 - l) N_{H,\mathrm{tf},2}(A) \oplus -\int_{{}^\phi TX/X} N_{H,\mathrm{tf},2}(A) & \text{for} \quad l = 2 \end{cases}$

(b) $N_{H,\mathrm{tf},l-1}(\frac{1}{x}V \circ A) = L_V N_{H,\mathrm{tf},l}(A)$

(c) $N_{H,\mathrm{tff},m-2}(\frac{\partial}{\partial t} \circ A) = L_{\frac{\partial}{\partial \tau}} N_{H,\mathrm{tff},m}(A)$

(d) $N_{H,\mathrm{tff},m-1}(\frac{1}{x}V \circ A) = L_{N_\phi(V)} N_{H,\mathrm{tff},m}(A)$

(e) $N_{H,\mathrm{hbf},p}(\frac{\partial}{\partial t} \circ A) = L_{\frac{\partial}{\partial t}} N_{H,\mathrm{hbf},p}(A)$

(f) $N_{H,\mathrm{hbf},p}(\frac{1}{x}V \circ A) = L_{\beta_H^* \frac{1}{x}V} N_{H,\mathrm{hbf},p}(A)$*, for* $V \in \Gamma([{}^\phi N\partial X])$.

Proof. First, recall the partial integration formula for the Lie derivative w.r.t. a vector field V on a manifold with boundary N:

$$\int_N \alpha \wedge L_V \beta = -\int_N (L_V \alpha) \wedge \beta + \int_{\partial N} \boldsymbol{\iota}(V)\alpha \wedge \beta.$$

Now let $f \in \dot{C}_c^\infty(X \times [0,\infty[, \Omega(X)\,dt)$ and $g \in \dot{C}_c^\infty(X \times [0,\infty[)$. We then have

$$\begin{aligned} \langle L_{\frac{\partial}{\partial t}}(A * g), f\rangle_{X\times[0,\infty[} &= -\langle A * g, L_{\frac{\partial}{\partial t}} f\rangle_{X\times[0,\infty[} + \int_{X\times\{0\}} \boldsymbol{\iota}(\frac{\partial}{\partial t}) A * g \cdot f \\ &= -\langle A, \beta_H^*(L_{\frac{\partial}{\partial t}} f) * g\rangle_{X_H^2} = -\langle A, L_{\beta_H^* \frac{\partial}{\partial t}} \beta_H^*(f * g)\rangle_{X_H^2} \\ &= \langle L_{\beta_H^* \frac{\partial}{\partial t}} A, \beta_H^*(f * g)\rangle_{X_H^2} - \int_{\mathrm{tf}\cup\mathrm{tff}} \boldsymbol{\iota}(\frac{\partial}{\partial t}) A \cdot \beta_H^*(f * g). \end{aligned}$$

Since near tf the density factor in $A \cdot \beta_H^*(f * g)$ is of the form $\rho_{\mathrm{tf}}^l\, \rho_{\mathrm{tff}}^m\, \frac{dt}{t} \beta_{H,L}^* \Omega(X) \Omega^\phi TX/X$ we see that the last integral in the above expression vanishes for $l > 2$, but contributes the constant

$$\int_X \left(\int_{{}^\phi TX/X} N_{H,\mathrm{tf},2}(A) \right) \langle f, g\rangle_{[0,\infty[}$$

for $l = 2$. Thus, recalling that

$$tL_{\frac{\partial}{\partial t}} = L_{t\frac{\partial}{\partial t}} - 1, \qquad L_{\frac{\partial}{\partial t}}\, dt = \, dt, \qquad \text{and} \quad L_{R_\phi}\, dt = 0,$$

and using Lemma 4.2, we find for $l > 2$:

$$\begin{aligned} N_{H,\mathrm{tf},l-2}(L_{\frac{\partial}{\partial t}} \circ A) &= t^{1-l/2} t L_{\beta_H^* \frac{\partial}{\partial t}} A|_{\mathrm{tf}} = t^{1-l/2}(L_{\beta_H^* t\frac{\partial}{\partial t}} - 1)A|_{\mathrm{tf}} \\ &= (L_{\beta_H^* t\frac{\partial}{\partial t}} - 2 + l/2)t^{1-l/2}A|_{\mathrm{tf}} = -\frac{1}{2}(L_{R^\phi} + 2 - l)N_{H,\mathrm{tf},l}(A). \end{aligned}$$

Note also that in the case $l = 2$ we have

$$N_{H,\mathrm{tf},0}(\frac{\partial}{\partial t} \circ A) = -\frac{1}{2} L_{R^\phi} N_{H,\mathrm{tf},2}(A) = -\frac{1}{2}\, dt L_{R^\phi}(N_{H,\mathrm{tf},2}(A)/\, dt),$$

therefore

$$\int_{{}^\phi TX/X} N_{H,\mathrm{tf},0}(\frac{\partial}{\partial t} \circ A) = 0,$$

i.e. the mean value condition is fulfilled. ■

As usual, the above results can all be formulated for operators with coefficients in a vector bundle. We leave this to the reader. Let us just make the remark that the construction is independent of the choice of connection made. Also, for $W \in \Gamma({}^\phi TX)$, the normal operators of $\tau^{1/2}\nabla^d_W$ and $\tau^{1/2}\nabla^\phi_W$ at tf and tff are the same.

Corollary 4.8 (a) $N_{H,\mathrm{tf}}(t^{1/2}\mathsf{D}) = N_{H,\mathrm{tf}}(t^{1/2}\mathsf{D}^d) = N_{H,\mathrm{tf}}(t^{1/2}\frac{1}{x}\mathsf{D}^\phi) = \mathbf{c}_{{}^\phi TX} \circ\, d_{{}^\phi TX}$

(b) $N_{H,\mathrm{tf}}(t\mathsf{D}^2) = \Delta_{{}^\phi TX}$, *the fibrewise Laplacian for the fibre metric* $g_{{}^\phi TX/X}$.

(c) $N_{H,\mathrm{tff}}(t^{1/2}\mathsf{D}) = t^{1/2}\frac{1}{x}N_{H,\mathrm{tff}}(\mathsf{D}^\phi) = \tau^{1/2}N_\phi(\mathsf{D}^\phi)$

(d) $N_{H,\mathrm{tff}}(t\mathsf{D}^2) = \tau N_\phi(\mathsf{D}^\phi)^2 = \tau\Delta_{{}^\phi N\partial X} + \tau(\mathsf{D}^{\phi,V})^2$. ■

The description of the action of D^2 can be described in the same way using Lemma 3.14 (d). We present this as part of the proof of

Theorem 4.9 (Parametrix for the Heat Equation) *There exists an element* $K \in \Psi^{2,2,0}_{H,\mathrm{cl}}(X, E)$ *such that*

$$\left(\frac{\partial}{\partial t} + \mathsf{D}^2\right) * K = I + R, \qquad R \in \Psi^{\infty,1,1}_{H,\mathrm{cl}}(X, E). \tag{82}$$

Proof. We start by solving the above equation to first order at tf. Applying $N_{H,\mathrm{tf},0}$ to both sides, we get first

$$\left(-\frac{1}{2}L_{R^\phi} + \Delta_{{}^\phi TX}\right) N_{H,\mathrm{tf},2}(K_2) = 0 \quad \text{and} \quad \int_{{}^\phi TX/X}(N_{H,\mathrm{tf},2}(K_2)/\, dt) = 1.$$

This is to be understood as an equation on the fibres of ${}^\phi TX$. Writing $N_{H,\mathrm{tf},2}(K_2) = k_2^{\mathrm{tf}}\, dt\, dvol_{{}^\phi TX/X}$ (and forgetting about the ${}^{\mathrm{tf}}$-index until it is really needed) we get

$$(-\frac{1}{2}(R_\phi + n) + \Delta_{{}^\phi TX})k_2 = 0, \qquad \int_{{}^\phi TX/X} k_2\, dvol_{{}^\phi TX/X} = 1.$$

Taking the Fourier transform in ${}^{\phi}TX$ yields

$$(\frac{1}{2}(\xi\frac{\partial}{\partial\xi}) + |\xi|^2)\widehat{k_2} = 0, \qquad \widehat{k_2}(0) = 1$$

the solution of which is easily seen to be

$$k_2 = \frac{1}{\sqrt{4\pi}^n}\exp(-\left\|.\right\|^2_{\phi}/4) \in x^0\mathcal{S}_{\text{fib}}({}^{\phi}TX),$$

which also shows that the solution is of order 2 at tff in the calculus. Thus the compatibility condition at the corner reads

$$N_{H,\text{tff},2}(K_2)_{\text{tf}\cap\text{tff}} = (x')^{-2}N_{H,\text{tf},2}(K_2)_{\text{tf}\cap\text{tff}} = k_2^{\text{tf}}\ d\tau\ dvol_{{}^{\phi}TX/X}$$

and the heat equation on the interior of tff is

$$\tau(L_{\tau\frac{\partial}{\partial\tau}} + \Delta_{{}^{\phi}N\partial X} + (\mathsf{D}^{\phi,V})^2)N_{H,\text{tff},2}(K_2) = 0 \tag{83}$$

Writing $N_{H,\text{tff},2}(K_2) = k_2^{\text{tff}}\ dvol_{{}^{\phi}N\partial X/Y}\ d\tau$ we get the equation

$$(\frac{\partial}{\partial\tau} + \Delta_{{}^{\phi}N\partial X} + (\mathsf{D}^{\phi,V})^2)k_2^{\text{tff}} = 0,$$

which is solved by

$$k_2^{\text{tff}} = \sqrt{4\pi\tau}^{-h-1}e^{-\|\cdot\|^2_{{}^{\phi}N\partial X}/4\tau}([\exp(-\tau(\mathsf{D}^{\phi,V})^2)]/\ d\tau\beta^*_{H,R}\ dvol_{M/Y}).$$

The compatibility condition at $\text{hbf}\cap\text{tff}$ is then

$$N_{H,\text{hbf},0}(K_2)|_{\text{tff}} = (x')^2N_{H,\text{tff},2}(K_2)|_{\text{hbf}} = \frac{e^{-\|\cdot\|^2_{{}^{\phi}N\partial X}/4}}{\sqrt{4\pi}^{h+1}}[\Pi_\circ]\frac{ds}{s}\ dvol_{{}^{\phi}N\partial X/\partial X}\ dt, \tag{84}$$

where $[\Pi_\circ] \in C^\infty(M\times_Y M,\ \text{END}\otimes\beta_R^*\Omega(M/Y))$ is the kernel of the projection onto the null space of $\mathsf{D}^{\phi,V}$. Then (84) is the initial value for the problem at hbf.

In order to solve the heat equation at hbf, recall that

$$x\mathsf{D} : C^\infty(X,E) \to C^\infty(X,E), \quad x\mathsf{D}|_{\partial X} = \mathsf{D}^{\phi,V}, \quad \text{and} \quad \mathsf{D} : C^\infty(X,E)^\circ \to C^\infty(X,E).$$

Thus for $G \in \Psi^{2,2,0}_{H,\text{cl}}(X)$ the expression D^2G a priori only lies in $\Psi^{0,0,-2}_{H,\text{cl}}(X)$. However, it follows from Lemma 3.14 (d) that $G|_{\text{hbf}}$ can be extended to the interior in such a way that $\mathsf{D}^2G \in \Psi^{\infty,1,0}_{H,\text{cl}}(X)$ and

$$\mathsf{D}^2G|_{\text{hbf}} = (\Pi_0\mathsf{D}\Pi_0)^2G_0|_{\text{hbf}} = I_b(\mathsf{D})^2G|_{\text{hbf}} \tag{85}$$

It is then clear that the heat equation at hbf can be solved by choosing $G|_{\text{hbf}}$ more concretely, namely

$$G|_{\text{hbf}} = [\exp(-tI_b(\mathsf{D})^2)] = [\exp(-t\mathsf{D}_Y^2)]\frac{1}{\sqrt{4\pi t}}e^{-(\ln(s))^2/4t}\frac{ds}{s},$$

and then use the extension which fulfills (85).
Since the solutions at the faces tf, tff, hbf are compatible, we can choose an extension $K_2 \in \Psi^{2,2,0}_{H,\mathrm{cl}}(X)$ with

$$N_{H,\mathrm{tf},2}(K_2) = k_2^{\mathrm{tf}}\, dt\, dvol_\phi, \quad N_{H,\mathrm{tff},2}(K_2) = k_2^{\mathrm{tff}}\, d\tau\, dvol_{\phi N\partial X/Y}, \quad N_{H,\mathrm{hbf},-v}(K_2) = g_0.$$

This gives a parametrix, which solves the heat equation to *first* order at all the faces

$$\left(\frac{\partial}{\partial t} + \mathsf{D}^2\right) * K_2 = I + R_2, \qquad R_2 \in \Psi^{1,1,1}_{H,\mathrm{cl}}(X).$$

In order to further improve our parametrix at tf, we have to successively find $K_l \in \Psi^{l,3,\infty}_{H,\mathrm{cl}}(X)$, $l = 3, 4, \ldots$ with

$$N_{\mathrm{tf},l-1}\left((\frac{\partial}{\partial t} + \mathsf{D}^2) * K_{l+1}\right) = -N_{\mathrm{tf},l-1}(R_l) \in (x')^{2-l}\mathcal{S}({}^\phi TX,\, dt\, dvol_\phi),$$

and

$$(\frac{\partial}{\partial t} + \mathsf{D}^2)(K_2 + \ldots + K_l) = I + R_l, \qquad R_l \in \Psi^{l-1,1,\infty}_{H,\mathrm{cl}}(X),$$

Again writing $N_{H,\mathrm{tf},l}(K_l) = k_l\, dt\, dvol_{\phi TX/X}$ and so on, we have

$$(\frac{1}{2}(\xi\frac{\partial}{\partial \xi}) + |\xi|^2 + l/2 - 1/2 + \Lambda)\widehat{k_{l+1}} = \widehat{r_l} \in (x')^{2-l}\mathcal{S}({}^\phi TX).$$

That this can be uniquely solved is the content of the following

Lemma 4.10 *Let $f \in \mathcal{S}(\mathbf{R})$ and $m > 0$. Then there is a unique solution $u \in \mathcal{S}(\mathbf{R})$ with*

$$((s\frac{\partial}{\partial s}) + 2s^2 + m)u = f.$$

Proof. First, the homogeneous equation

$$((s\frac{\partial}{\partial s}) + 2s^2 + m)u_0 = 0$$

is uniquely solved by $u_0 = e^{-s^2}s^{-m}$. Writing $u = u_0u_1$ we see that

$$f = ((s\frac{\partial}{\partial s}) + 2s^2 + m)u_0u_1 = u_0 s\frac{\partial}{\partial s}u_1.$$

Thus u_1 solves the equation

$$\frac{\partial}{\partial s}u_1 = (su_0)^{-1}f,$$

i.e. u is of the form

$$u = e^{-s^2}s^{-m}\int_0^s f(r)e^{r^2}r^{m-1}\, dr,$$

which is obviously in $\mathcal{S}(\mathbf{R})$. ∎

This also finishes the proof of the Theorem. ■

This result can be further improved by formally inverting the term $I + R$. From the composition formula, Theorem 4.6, we get:

$$R^j \in \Psi_H^{\mathcal{G}_j}(X), \quad G_{\mathrm{hbf},j} \geq j, \quad G_{\mathrm{tff}} = j + \mathbf{N}, \quad G_{\mathrm{tf},j} = \infty.$$

Thus the powers R^j can be asymptotically summed, and the Neumann series

$$(1+R)^{-1} = \sum_{k=0}^{\infty} (-R)^k = 1 + \widetilde{R}$$

is valid in that sense. We can then define

$$\begin{aligned} K_{\mathrm{sm}} &:= K(1+R)^{-1} = K + K\widetilde{R} \in \Psi_H^{\mathcal{H}}(X), \quad \text{with} \\ & H_{\mathrm{tf}} = 2 + \mathbf{N}, \quad H_{\mathrm{tff}} = 2 + \mathbf{N}, \quad H_{\mathrm{hbf}} \subset \mathbf{N} \times \mathbf{N}, \quad T(H_{\mathrm{hbf}}) = \{0\}. \end{aligned} \tag{86}$$

This gives a parametrix of the heat equation up to a smoothing remainder

$$(\frac{\partial}{\partial t} + \mathsf{D}^2) K_{\mathrm{sm}} = 1 + R_{\mathrm{sm}}, \qquad R_{\mathrm{sm}} \in \dot{C}^{\infty}(X_H^2, \mathrm{KD}_H).$$

Now, as in the final step of the resolvent construction, the operator $1 + R_{\mathrm{sm}}$ is a *Volterra* operator, which can be inverted and whose inverse $(1 + R_{\mathrm{sm}})^{-1}$ is again of that type. We have therefore shown

Theorem 4.11 (Heat Kernel) *The heat kernel for* D^2 *is given by*

$$\mathsf{K} = K_{\mathrm{sm}}(1 + R_{\mathrm{sm}})^{-1} \in \Psi_H^{\mathcal{H}}(X, E), \qquad \mathcal{H} \quad \textit{as in} \quad (86).$$

Its normal operators at the different faces are (well defined and) given by

(a) $N_{H,\mathrm{tf},2}(\mathsf{K}) = \frac{1}{\sqrt{4\pi}^n} \exp(- \|.\|_{\phi}^2 /4) \, dvol_{\phi TX/X} \;\; dt$

(b) $N_{H,\mathrm{tff},2}(\mathsf{K}) = \frac{1}{\sqrt{4\pi\tau}^{h+1}} \exp(- \|.\|_{\phi N\partial X}^2 /4\tau) \, dvol_{\phi N\partial X/\partial X}([\exp(-\tau(\mathsf{D}^{\phi,V})^2)]$

(c) $N_{H,\mathrm{hbf},0}(\mathsf{K}) = [\Pi_{\circ}][\exp(-t\mathsf{D}_Y^2)] \frac{1}{\sqrt{4\pi t}} e^{-(\ln(s))^2/4t} \frac{ds}{s}$

■

Remark 4.12 There are several possible extensions of this result which are not presented here, since they are not crucial to our argument. First, one can eliminate half of the coefficients in the expansion of the heat kernel at tf, by analyzing their behavior under the "flip map" in X_H^2. This procedure is described in Chapter 7 of [Me3]. Also, direct analysis of the model problem at hbf shows that the heat kernel expansion does not exhibit log-terms at that face. Thus really, $H_{\mathrm{hbf}} = \mathbf{N}$

Still, Theorem 4.11 gives a rather crude description of the expansions of the heat kernel at tf and tff. Especially, it does not yet allow us to calculate the regularized heat supertrace at small times. For this, we have to introduce a refinement of the heat calculus, which keeps track of the Clifford degree of the coefficients in the expansions. This will be done in the next Chapter.

5 Rescaled d-Heat Calculus

In this Chapter we present a refinement of the d-heat calculus presented in Chapter 4. This "rescaled" calculus encodes the behavior of the Taylor coefficients of the heat kernel at tf and tff w.r.t. the Clifford filtration of the endomorphism bundle at these faces. This generalization of Getzler's technique (compare [BGV]) was introduced in [Me3]. A good description of this philosophy can also be found in [DaMe].

While the results in earlier Sections are independent of the parity of the dimension n of X we now assume that $n = v + h + 1$ is *even*.

5.1 The Double Rescaling

In this Section we show how the natural filtration by Clifford degree of the bundle $\mathrm{END}(E)$ over the faces tf and tff can be used to define a new bundle $\mathrm{Gr}(\mathrm{END})$ *rescaled* at these faces. First, we have to describe the Clifford action on $\mathrm{END}(E)$ in more detail.

Elements $\alpha \in \mathrm{Cl}({}^dT^*X)$ act on β_L^*E via Clifford left multiplication and on $\beta_R^*E^*$ via Clifford right multiplication

$$\mathbf{c}_{d,L}(\alpha)\beta_L^*\xi = \beta_L^*\mathbf{c}_d(\alpha)\xi, \qquad \mathbf{c}_{d,R}(\alpha)\beta_R^*\xi^* = \beta_R^*\xi^* \circ \mathbf{c}_d(\alpha).$$

The right action can be turned into a left action, using the transpose map $(\cdot)^\dagger$ on $\mathrm{Cl}({}^dT^*X)$:

$$\overline{\mathbf{c}}_{d,R}(\alpha)\beta_R^*\xi^* = (-1)^{|\alpha||\xi^*|}\beta_R^*\xi^* \circ \mathbf{c}_d(\alpha)^\dagger.$$

Since the fibres of ${}^dT^*X$ are even dimensional, the Clifford modules E and E^*, as well as $\mathrm{END}(E)$ are naturally $\mathbf{Z}_2$-graded by the action of the volume element in $\mathrm{Cl}({}^dT^*X)$. The idea of introducing a rescaling into our heat calculus is based on the observation that the bundle $\mathrm{END}(E)$ over the heat space X_H^2 carries a filtration when restricted to the different faces.

Lemma 5.1 *At* tf, tff *and* hbf *the bundle* END *has the following form*

(a) $\mathrm{END}(E)|_{\mathrm{hbf}} = \mathrm{Cl}(\frac{dx}{x}) \otimes \mathrm{END}_{\mathrm{Cl}(\frac{dx}{x})}(E)$

(b) $\mathrm{END}(E)|_{\mathrm{tff}} = \phi^*\,\mathrm{Cl}({}^{\mathrm{b}}T^*B|_Y) \otimes \mathrm{END}_{\mathrm{Cl}({}^{\mathrm{b}}T^*B|_Y)}(E)$.

(c) $\mathrm{END}(E)|_{\mathrm{tf}} = \mathrm{Cl}({}^{\mathrm{d}}T^*X) \otimes \mathrm{End}_{\mathrm{Cl}({}^{\mathrm{d}}T^*X)}(E)$

The restrictions of the connection $\nabla^{\mathrm{END},d}$ split accordingly. Note that (b) is true independent of the parity of the dimension h of the base Y. Also, part (a) will be of no importance to us. We state it because its proof is exemplary for the proofs of (b) and (c).

Proof. First, since the bundle END is just pulled back from $X^2 \times [0,\infty[$, it suffices to prove the above statements in the blown down picture.

To prove (a), restrict to the face hbf. There, we can assume the coordinates x, x' to be *paired* (compare Section 2.1). For a section $Q \in \Gamma(X_H^2, \mathrm{END})$, homogeneous w.r.t. to the grading of END, we can therefore define

$$[\mathbf{c}_d(\frac{dx}{x}), Q] := \mathbf{c}_d(\frac{dx}{x}) \circ Q - (-1)^{|Q|}Q \circ \mathbf{c}_d(\frac{dx'}{x'}).$$

Then

$$\Big[\mathbf{c}_d(\frac{dx}{x}), [\mathbf{c}_d(\frac{dx}{x}), Q]\Big] = 0, \qquad \Big[\mathbf{c}_d(\frac{dx}{x}), Q + \frac{1}{2}\mathbf{c}_d(\frac{dx}{x}) \circ [\mathbf{c}_d(\frac{dx}{x}), Q]\Big] = 0,$$

which shows that the map

$$Q \longmapsto \Big(-\frac{1}{2}\mathbf{c}_d(\frac{dx}{x}) \circ [\mathbf{c}_d(\frac{dx}{x}), Q]\Big) + \Big(Q + \frac{1}{2}\mathbf{c}_d(\frac{dx}{x}) \circ [\mathbf{c}_d(\frac{dx}{x}), Q]\Big)$$

has image in the RHS of (a), and is an isomorphism of algebras. This proves (a).

Parts (b) and (c) can be obtained by iteration of this scheme using an orthonormal basis in $\mathrm{Cl}({}^bT^*B)$ and $\mathrm{Cl}({}^dT^*X)$ respectively. The important observation is that an element α in $\Gamma(Y, {}^bT^*B)$ can be lifted from the left or the right to tff, and we can define as above

$$[\mathbf{c}_d(\alpha), Q] := \mathbf{c}_d(\beta^*_{H,L}\alpha) \circ Q - (-1)^{|Q|} Q \circ \mathbf{c}_d(\beta^*_{H,R}\alpha),$$

and the same can be done for *any* $\alpha \in \Gamma({}^dT^*X)$ at tf. ■

To rescale the bundle $\mathrm{End}(E)$ we want to continue the filtrations at tf and tff to the interior using parallel transport along a normal vector field, and define the space of sections of our rescaled bundle as the set of sections, whose lth normal derivative has degree at most l at the corresponding face. This encodes the (so far: supposed) behavior of the heat kernel at these faces.

To explain the construction, assume for a moment that the bundle and the connection are trivial. A local basis for the rescaled bundle near tf ∩ tff could then look like

$$\rho_{\mathrm{tf}}\, \rho_{\mathrm{tff}}\, \frac{dx}{x}, \quad \rho_{\mathrm{tf}}\, \rho_{\mathrm{tff}}\ d\mathbf{y}, \quad \rho_{\mathrm{tf}}\, x\, d\mathbf{z}\, (!), \tag{87}$$

and (wedge-) products thereof. For simplicity of notation, we have also dropped the "coefficients" $\mathrm{END}_{\,\mathrm{Cl}({}^dT^*X)}(E)$, i.e. we have taken E to be the spinor bundle.

In reality however, neither our bundle nor the connection are trivial. Also, we would like to avoid the choice of a normal vector field, at least for the mere definition of the rescaled bundle. The relevant connection on $\mathrm{END}(E)$ in our context is the lifted connection

$$\nabla = \beta^*_{H,L}\nabla^{E,d} \otimes \beta^*_{H,R}\nabla^{E^*,d} \otimes \beta^*_H\, dt\frac{\partial}{\partial t} : \Gamma(\,\mathrm{END}(E)) \longrightarrow \Gamma(T^*X^2_H \otimes \mathrm{END}(E)),$$

and the associated curvature will be denoted by K. Also, recall that the identification of the faces tf and tff was based on the following identifications for $A \in \Gamma({}^\phi TX)$:

$$\tau^{1/2}\beta^*_L A|_{\mathrm{tf}} \in T\mathrm{tf}/X \quad \text{with} \quad A \in {}^\phi TX \subset T^\phi TX/X \tag{88}$$

$$\beta^*_L A|_{\mathrm{tff}} \in T\mathrm{tff}/\partial X \quad \text{with} \quad N_\phi(A) \in T^\phi N\partial X/Y. \tag{89}$$

Rescaling at tff

We agree to write

$$\mathrm{Cl}^k_{\mathrm{tff}} \quad \hat{=} \quad \phi^*\ \mathrm{Cl}^k({}^{\mathrm{b}}T^*B|_Y) \otimes \mathrm{END}_{\,\mathrm{Cl}({}^{\mathrm{b}}T^*B|_Y)}(E) \quad \text{at} \quad \mathrm{tff}$$

for the filtration space in Lemma 5.1. Our construction then starts from the following simple observations (not all of which are going to be used right away):

Lemma 5.2 (Basic Properties of the Curvature and the Filtration at tff)

(a) *Let* A *be a vector tangent to* tff. *Then* ∇_A *preserves the filtration of* END *at* tff.

(b) $K(T_{\mathrm{tff}}, \cdot)|_{\mathrm{tff}} \in N^*\mathrm{tff} \otimes \mathrm{Cl}^1_{\mathrm{tff}}$ *whenever* T_{tff} *is tangent to the fibres* tff/Y.

(c) $(\nabla_N K)(T_{\mathrm{tff}}, U_{\mathrm{tff}}) \in \mathrm{Cl}^1_{\mathrm{tff}}$, *for* $T_{\mathrm{tff}}, U_{\mathrm{tff}}$ *tangent to the fibres* tff/Y.

(d) *The covariant derivatives of the curvature* $\nabla^l K$ *are at most of order 2 at* tf *and* tff.

Proof. Since the connection ∇ is the connection on END pulled back from X^2, it suffices to prove (a) for vectors A, tangent to the fibre diagonal F_ϕ. In view of the proof of Lemma 5.1 and the fact, shown in Lemma 1.5(c) and (15), that ∇^d_A preserves the decomposition ${}^dT^*X = {}^dN^*\partial X \oplus {}^dV^*\partial X$ over the boundary, the assertion is then clear.

For part (b) to (d), look at the decomposition of the curvature (and see the remark at the end of Appendix A.1)

$$\begin{aligned} K &= \beta_L^*(R^{S,d} + F^{E/S,d}) + \beta_R^*(R^{S^*,d} + F^{E/S,d}) \\ &\qquad = \frac{1}{2}\beta_L^*\mathbf{c}_d(R^d) - \frac{1}{2}\beta_R^*\mathbf{c}_d(R^d) + \beta_L^* F^{E/S,d} + \beta_R^* F^{E/S,d}. \end{aligned} \tag{90}$$

Parts (b)-(d) then follow immediately from Proposition 1.14. ∎

As a first step in the rescaling procedure, we can now define the space of sections, *rescaled* at tff as

$$\mathcal{D}_{\mathrm{tff}} := \left\{ u \in \Gamma(X^2_H, \mathrm{END}) |\, \nabla^k u|_{\mathrm{tff}} \in (T^*X^2_H)^k \otimes \mathrm{Cl}^k_{\mathrm{tff}} \right\}.$$

First note that, although it looks like this definition uses a connection on TX^2_H, a second look convinces us that no such choice is necessary. In fact, by choosing any vector field N_{tff} (strictly) normal to tff, and using the compatibility of the connection and the filtration at tff as stated in Lemma 5.2(a) and (d), we could also write

$$\mathcal{D}_{\mathrm{tff}} = \left\{ u \in \Gamma(X^2_H, \mathrm{END}) | \quad (\nabla_{N_{\mathrm{tff}}})^k u|_{\mathrm{tff}} \in \mathrm{Cl}^k_{\mathrm{tff}} \right\}. \tag{91}$$

As in [Me3] this is the space of sections of a vector bundle $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})$. We note this as

Proposition 5.3 (a) $\mathcal{D}_{\mathrm{tff}} = \Gamma(X^2_H, \mathrm{Gr}_{\mathrm{tff}}(\mathrm{END}))$.

(b) $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})|_{\mathrm{tf}} = \mathrm{Cl}([{}^dV^*\partial X]) \otimes \mathrm{End}_{\mathrm{Cl}({}^dT^*X)}(E) = \mathrm{Cl}(x^{\mathrm{b}}T^*X) \otimes \mathrm{End}_{\mathrm{Cl}({}^dT^*X)}(E)$

Proof. (a) is proved as in Chapter 7 of [Me3] by writing down a local basis for $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})$. To prove (b), we first check that sections in $\mathrm{Cl}([{}^dV^*\partial X])$ satisfy the condition in the definition (91) of $\mathcal{D}_{\mathrm{tff}}$. Note that for sections $\alpha_i \in \Gamma([{}^dV^*\partial X])$ the derivative $\nabla^d_N \alpha_i$ still lies in ${}^dT^*X$. Thus

$$(\nabla_N)^k(\alpha_1 \wedge \ldots \wedge \alpha_l)|_{\partial X} \in \Lambda^{k \wedge l\, d}T^*X \otimes \Lambda^{0 \vee l-k}[{}^dV^*\partial X],$$

and $(\nabla_N)^k(\mathbf{c}_d(\alpha_1)\ldots\mathbf{c}_d(\alpha_l))$ is of order at most k at tff. The proof of the reverse inclusion proceeds by induction and is left to the reader. ∎

Let us describe the structure of the bundle $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$ in more detail. To see what the bundle looks like at tff, note that the space of sections over tff is

$$\mathcal{D}_{\mathrm{tff}}/\rho_{\mathrm{tff}}\mathcal{D}_{\mathrm{tff}},$$

and the "denominator" space can be described as follows

Lemma 5.4 $\rho_{\mathrm{tff}}\mathcal{D}_{\mathrm{tff}} = \{u \in \mathcal{D}_{\mathrm{tff}} \mid \nabla^k u|_{\mathrm{tff}} \in (T^*X^2_H|_{\mathrm{tff}})^k \otimes \mathrm{Cl}^{k-1}_{\mathrm{tff}}\}$. ■

Thus *choosing* a projection P_{tff} of $TX^2_H|_{\mathrm{tff}}$ onto a subspace (or, if one prefers, a vector field $\nu_{\mathrm{tff}} = P_{\mathrm{tff}}/\,dx$) normal to tff, the map

$$u \longmapsto u|_{\mathrm{tff,Gr}} = (u|_{\mathrm{tff}}, P_{\mathrm{tff}}\nabla u|_{\mathrm{tff}}, \ldots\,, \frac{1}{(h+1)!}P_{\mathrm{tff}}(\nabla^{h+1}u)|_{\mathrm{tff}})$$

yields an exact sequence

$$0 \longrightarrow \mathcal{D}_{\mathrm{tff}}/\rho_{\mathrm{tff}}\,\mathcal{D}_{\mathrm{tff}} \longrightarrow \mathcal{D}_{\mathrm{tff}} \overset{|_{\mathrm{tff,Gr}}}{\longrightarrow} C^\infty(\mathrm{tff}, \textstyle\bigoplus_k (N^*\mathrm{tff})^k \otimes \mathrm{Cl}^k_{\mathrm{tff}}\,/\,\mathrm{Cl}^{k-1}_{\mathrm{tff}}) \longrightarrow 0.$$

The vector bundle appearing on the right represents the restriction of $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$ to tff. It will be denoted by

$$\mathcal{N}\Lambda(\mathrm{tff}) = \textstyle\bigoplus_k \mathcal{N}\Lambda^k(\mathrm{tff}).$$

However, the reader should keep in mind that the restriction *map* depends on the choice of projection P_{tff}.

We have chosen to keep track of the factors $N^*\mathrm{tff}$ for a while (and to remind the reader of this fact by introducing "$\mathcal{N}$" into the notation as above) because this allows for an easy description of the induced connections: Denote by ${}^bT_{\mathrm{tff}}X^2_H$ the vector bundle whose sections are vector fields over X^2_H which are tangent to tff. Also ${}^bT_{\mathrm{tff}/Y}X^2_H$ is defined as the vector bundle of vector fields tangent to the *fibres* of $\mathrm{tff} \to Y$. On $N\mathrm{tff}$ we will use the (b-) connection

$$\nabla^{N\mathrm{tff}} : \Gamma(\mathrm{tff}, N\mathrm{tff}) \longrightarrow \Gamma(\mathrm{tff}, {}^bT^*_{\mathrm{tff}}X^2_H|_{\mathrm{tff}} \otimes N\mathrm{tff}) \quad \nabla^{N\mathrm{tff}}_T N := [T, N]. \tag{92}$$

The reader should note that this is only a b-connection, since for instance, $[xN', N] = -(Nx)N'$. However, we can restrict it to an action of $T\mathrm{tff}/Y$ on $N\mathrm{tff}$ in a reasonably natural way, by using the inclusion map

$$[] : T\mathrm{tff}/Y \longrightarrow {}^bT^*_{\mathrm{tff}}X^2_H|_{\mathrm{tff}}, \quad T \longmapsto [T], \quad \text{with} \quad \frac{dx}{x}([T]) = O(x).$$

We can now describe the connection induced on the rescaled bundle $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$ and its representation under restriction to the boundary.

Remark 5.5 It has been above this author's capabilities to devise a good notation for the following developments. A connection ∇ will appear in two distinct functions, first as the connection appearing in the definition of the space of sections (e.g. $\mathcal{D}_{\mathrm{tff}}$) or of the restriction map (e.g. $|_{\mathrm{tff,Gr}}$), second as the connection *acting* on sections of the rescaled bundle in question (e.g. $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$). In the latter case, the connection will usually have to be restricted to a specific class of vectors (e.g. ${}^bT_{\mathrm{tff}/Y}X^2_H$) and will then be written in *boldface* $\boldsymbol{\nabla}$. As an additional piece of notation, we will write $\mathbb{K} := [\boldsymbol{\nabla}, \nabla]$ and $\mathbf{K} := [\boldsymbol{\nabla}, \boldsymbol{\nabla}]$. The complication with this notation arises from the fact that we have to rescale *twice*, thus the connections change their role in the process.

Using these conventions we get

Proposition 5.6 *Let* $T, U \in \Gamma({}^bT_{\mathrm{tff}/Y}X^2_H)$*. Then*

(a) $\boldsymbol{\nabla}_T$ *maps* $\mathcal{D}_{\mathrm{tff}}$ *to itself and thus induces a connection* $\boldsymbol{\nabla}_T \overset{\wedge}{=} \boldsymbol{\nabla}^{\mathrm{Gr(tff)}}_T$ *on* $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$ *with*

$$(\boldsymbol{\nabla}_T u)|_{\mathrm{tff,Gr}} = (\boldsymbol{\nabla}_T - P_{\mathrm{tff}}\mathbb{K}(T,\cdot) - \frac{1}{2}P_{\mathrm{tff}}\nabla.\mathbb{K}(T,\cdot))(u|_{\mathrm{tff,Gr}}) =: \boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_T(u|_{\mathrm{tff,Gr}}),$$

where $P_{\mathrm{tff}}\mathbb{K}(T,\cdot) \in \mathcal{N}\Lambda^1(\mathrm{tff})$ *and* $P_{\mathrm{tff}}\nabla.\mathbb{K}(T,\cdot) \in \mathcal{N}\Lambda^2(\mathrm{tff})$*.*

(b) $\mathbf{K}(T,U) \in \mathcal{D}_{\mathrm{tff}}$*. Its (rescaled) restriction to* tff *is just the curvature of* $\boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}$*:*

$$(\mathbf{K}(T,U)u)|_{\mathrm{tff,Gr}} = \mathbf{K}(T,U)|_{\mathrm{tff,Gr}} \cdot u|_{\mathrm{tff,Gr}} = \mathbf{K}^{\mathcal{N}\Lambda(\mathrm{tff})}(T,U) \cdot u|_{\mathrm{tff,Gr}}.$$

Proof. We make abundant use of the following commutator identity:

Lemma 5.7 *Let* $T \in \Gamma({}^bT_{\mathrm{tff}/Y}X^2_H)$*,* $N \in TX^2_H$*. Then over* X^2_H

(a) $\nabla_T\boldsymbol{\nabla}(N) = \boldsymbol{\nabla}_T\nabla_N - \nabla_{[T,N]} - \mathbb{K}(T,N)$ *(* $= \boldsymbol{\nabla}\nabla(T,N) - \mathbb{K}(T,N)$ *restricted to* tff*)*

(b) $P_{\mathrm{tff}}\boldsymbol{\nabla}_T\nabla - \boldsymbol{\nabla}_T P_{\mathrm{tff}}\nabla$ *is of order* 0 *at* tff

Proof. (a) is just a consequence of our definition of $\boldsymbol{\nabla}, \mathbb{K}$ and $\boldsymbol{\nabla}^{N^*\mathrm{tff}}$. For the proof of (b) consider

$$\boldsymbol{\nabla}_T\nabla(P_{\mathrm{tff}}N) - \boldsymbol{\nabla}_T P_{\mathrm{tff}}\nabla(N) = \nabla_{P_{\mathrm{tff}}[T,N]} - \nabla_{[T,P_{\mathrm{tff}}N]}$$

This is of order 0 at tff, since it is covariant differentiation w.r.t. a vector tangent to tff as can be seen from

$$P_{\mathrm{tff}}([T,P_{\mathrm{tff}}N] - P_{\mathrm{tff}}[T,N]) = P_{\mathrm{tff}}[T,P_{\mathrm{tff}}N - N] = 0.$$

This finishes the proof of the Lemma. ■

The proof of part (a) of the Proposition now proceeds as follows: Let $u \in \mathcal{D}$. Then using the identies from Lemma 5.7 we get modulo lower order at tff:

$$\frac{1}{k!}P(\nabla^k)\boldsymbol{\nabla}_T u = \frac{1}{k!}\boldsymbol{\nabla}_T P(\nabla^k)u - P\mathbb{K}(T,\cdot)\frac{1}{(k-1)!}P(\nabla^{k-1})u - \frac{1}{2}(P\nabla\mathbb{K}(T,\cdot))\frac{1}{(k-2)!}P(\nabla^{k-2})u.$$

Higher derivatives of the curvature do not contribute because of Lemma 5.2(d).

To show (b), note that is clear from Lemma 5.2 that

$$\mathbf{K}(T,U)|_{\mathrm{tff}} \in \mathrm{Cl}^0_{\mathrm{tff}}, \qquad (\nabla_N\mathbf{K})(T,U) \in \mathrm{Cl}^1_{\mathrm{tff}},$$

all the higher derivatives being of order at most 2. Thus $\mathbf{K}(T,U)$ is in fact an element in $\mathcal{D}_{\mathrm{tff}}$. The first equality in (b) is now an immediate consequence of Leibniz' rule while the second equality can be obtained by playing around with the definition in part (a):

$$\begin{aligned}\mathbf{K}^{\mathcal{N}\Lambda(\mathrm{tff})}(T,U)\cdot u|_{\mathrm{tff,Gr}} &= (\boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_T\boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_U - \boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_U\boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_T - \boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_{[T,U]})\cdot u|_{\mathrm{tff,Gr}} \\ &= [(\boldsymbol{\nabla}_T\boldsymbol{\nabla}_U - \boldsymbol{\nabla}_U\boldsymbol{\nabla}_T - \boldsymbol{\nabla}_{[T,U]})u]|_{\mathrm{tff,Gr}}\end{aligned}$$

which is just $(\mathbf{K}(T,U)u)|_{\mathrm{tff,Gr}}$. ■

Rescaling at tf

In a second step, we now want to rescale the bundle $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})$ at tf using its induced filtration as described in Proposition 5.3(b)

$$\mathrm{Cl}^k_{\mathrm{tf}} \quad \overset{\triangle}{=} \quad \mathrm{Cl}^k(x^bT^*X) \otimes \mathrm{End}_{\mathrm{Cl}(^{\mathrm{d}}T^*X)}(E),$$

over tf and the induced (partial) connection $\boldsymbol{\nabla}$ described in Proposition 5.6. To do this, we need the following addendum to Lemma 5.2, which shows that the filtration and the connection are compatible:

Lemma 5.8 (Basic Properties of the Curvature and the Filtration at tf) *Let $A, T_{\mathrm{tf}}, N \in \Gamma(^bT_{\mathrm{tff}/Y}X^2_H)$. Then*

(a) *Let B be tangent to* tf. *Then $\boldsymbol{\nabla}_B \overset{\triangle}{=} \boldsymbol{\nabla}^{\mathrm{Gr(tff)}}_B$ preserves the filtration of* $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})$ *at* tf.

(b) $\mathbf{K}(T_{\mathrm{tf}},N)|_{\mathrm{tf}} = 0$ *whenever T_{tf} is tangent to the fibres of* tf $\to X$.

Proof. Again, since $\nabla^{\mathrm{Gr(tff)}}$ is induced from ∇^{END}, away from tff it suffices to prove (a) for vectors B, tangent to the diagonal $\triangle \times 0$. In view of the proof of Lemma 5.1 and Lemma 1.5(c) and (15)) the assertion is then clear. For (b), just note that T_{tf} can be thought of as being of the form $\tau^{1/2}\beta_L^*V$ for $V \in \Gamma(^\phi TX)$. Away from tff (90) then implies

$$\mathbf{K}(T_{\mathrm{tf}},N) = \mathbf{K}(\beta_L^*\tau^{1/2}V,N) = \tau^{1/2}K(\beta_L^*V,N),$$

which is obviously 0 when restricted to tf. The assertions extend to tff by continuity. ■

Now the space of sections of the bundle rescaled at tff and tf can be defined, using a vector field N_{tf} normal to tf but tangent to the fibres of tff:

$$\mathcal{D} \equiv \mathcal{D}_{\mathrm{tff,tf}} := \left\{ u \in \mathcal{D}_{\mathrm{tff}} |\, (\nabla^{\mathrm{Gr(tff)}}_{N_{\mathrm{tf}}})^k u|_{\mathrm{tf}} \in \mathrm{Cl}^k_{\mathrm{tf}} \right\}.$$

Again, this definition gives the space of sections of a vector bundle $\mathrm{Gr}(\mathrm{END})$ and is independent of the choices made:

Proposition 5.9 (a) $\mathcal{D} = \Gamma(X^2_H, \mathrm{Gr}(\mathrm{END}))$.

(b) $\mathcal{D} = \left\{ u \in \Gamma(\mathrm{END}) |\, \nabla^k u|_{\mathrm{tff}} \in (T^*X^2_H)^k \otimes \mathrm{Cl}^k_{\mathrm{tff}}, \quad \nabla^l u|_{\mathrm{tf}} \in (T^*X^2_H)^l \otimes \mathrm{Cl}^l_{\mathrm{tf}} \right\}.$

Proof. (a) follows from iterating the argument in the case of a simple rescaling, (b) follows from Lemma 5.8 and the fact that $\nabla^{\mathrm{Gr(tff)}}$ on $\mathrm{Gr}_{\mathrm{tff}}(\mathrm{END})$ is just induced from the connection on END. ■

From now on, part (b) in this Proposition will be our preferred definition of $\mathcal{D}$.

The rescaling $\mathrm{Gr}_{\mathrm{tf}}(\mathcal{N}\Lambda(\mathrm{tff}))$ of the bundle $\mathcal{N}\Lambda_{\mathrm{tff}}$ at tff $\cap$ tf can be defined in the same way, using the connection $\nabla^{\mathcal{N}\Lambda(\mathrm{tff})}$ and the filtration at tf$\cap$tff induced from Lemma 5.3(b). The description of the bundle Gr(END) over the faces tf and tff then follows the lines set out above: The space of sections over the faces $F = $ tf, or $F = $ tff is given by

$$\mathcal{D}/\rho_F\mathcal{D},$$

which can be described using

Lemma 5.10 $\rho_F\mathcal{D} = \{u \in \mathcal{D} | \quad \nabla^k u|_F \in (T^*X^2_H|_F)^k \otimes \mathrm{Cl}_F^{k-1}\}$. ■

Then *choosing* projections $P = P_F$ of $TX^2_H|_F$ onto a normal subspace to F – of course P_{tf} needs to be tangent to the fibres of tff – the maps

$$u \longmapsto \begin{cases} u|_{\mathrm{tf,Gr}} = (u|_{\mathrm{tf}}, P_{\mathrm{tf}}\nabla u|_{\mathrm{tf}}, \dots, \frac{1}{n!}P_{\mathrm{tf}}(\nabla^n u)|_{\mathrm{tf}}) \\ u|_{\mathrm{tff,Gr}} = (u|_{\mathrm{tff}}, P_{\mathrm{tff}}\nabla u|_{\mathrm{tff}}, \dots, \frac{1}{(h+1)!}P_{\mathrm{tff}}(\nabla^{h+1} u)|_{\mathrm{tff}}) \end{cases}$$

give exact sequences

$$0 \longrightarrow \mathcal{D}/\rho_{\mathrm{tff}}\,\mathcal{D} \longrightarrow \mathcal{D} \overset{|_{\mathrm{tff,Gr}}}{\longrightarrow} C^\infty(\mathrm{tff}, \mathrm{Gr}_{\mathrm{tf}}(\mathcal{N}\Lambda(\mathrm{tff}))) \longrightarrow 0,$$

$$0 \longrightarrow \mathcal{D}/\rho_{\mathrm{tf}}\,\mathcal{D} \longrightarrow \mathcal{D} \overset{|_{\mathrm{tf,Gr}}}{\longrightarrow} C^\infty(\mathrm{tf}, \mathcal{N}\Lambda(\mathrm{tf})) \longrightarrow 0,$$

where we have written $\mathcal{N}\Lambda(\mathrm{tf}) := \bigoplus_k (N^*\mathrm{tf})^k \otimes \mathrm{Cl}^k_{\mathrm{tf}} \,/\, \mathrm{Cl}^{k-1}_{\mathrm{tf}}$. Of course, there are also induced restriction maps

$$|_{\mathrm{tff,Gr}} : C^\infty(\mathrm{tf}, \mathcal{N}\Lambda(\mathrm{tf})) \to C^\infty(\mathrm{tf} \cap \mathrm{tff}, \mathcal{N}\mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff})),$$

$$|_{\mathrm{tf,Gr}} : C^\infty(\mathrm{tff}, \mathrm{Gr}_{\mathrm{tf}}(\mathcal{N}\Lambda(\mathrm{tff}))) \to C^\infty(\mathrm{tf} \cap \mathrm{tff}, \mathcal{N}\mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff})),$$

defined by

$$u|_{\mathrm{tff,Gr}} = (\frac{1}{j!}P_{\mathrm{tff}}(\nabla^j u)|_{\mathrm{tff}})_{j=1,\dots,h+1} \quad \text{and} \quad u|_{\mathrm{tf,Gr}} = (\frac{1}{j!}P_{\mathrm{tf}}(\nabla^{\mathcal{N}\Lambda(\mathrm{tff})})^j u|_{\mathrm{tf}})_{j=1\dots,n}.$$

The vector bundle on the RHS is $\mathcal{N}\mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff}) = \bigoplus_{m=1}^n \mathcal{N}\mathcal{N}\Lambda^m(\mathrm{tf}, \mathrm{tff})$. Its part of grade m is defined as

$$(N^*\mathrm{tf})^m \otimes \Big[\bigoplus\nolimits_{j+l=m} \Lambda^{jd}V^*\partial X \otimes (N^*\mathrm{tff})^l \otimes \Lambda^l \phi^{*b}T^*B\Big] \otimes \mathrm{END}_{\mathrm{Cl}({}^dT^*X)}(E). \tag{93}$$

The restriction maps have been defined in such a way that restriction to the corner $\mathrm{tf} \cap \mathrm{tff}$ is well defined independent of the order

$$u|_{\mathrm{tf}\cap\mathrm{tf,Gr}} := u|_{\mathrm{tff,Gr}}|_{\mathrm{tf,Gr}} = u|_{\mathrm{tf,Gr}}|_{\mathrm{tff,Gr}}. \tag{94}$$

Defining the (b-) connection on $N\mathrm{tf}$ in analogy with (92) the analogue of Proposition 5.6 reads

Proposition 5.11 *Let* $T \in \Gamma({}^bT_{\mathrm{tff}/Y,\mathrm{tf}/X}X^2_H)$, *i.e.* T *is tangent to the fibres* tf/X *and* tff/Y. *Then* $\boldsymbol{\nabla}_T$ *maps* $\mathcal{D}$ *to itself and thus induces a connection* $\boldsymbol{\nabla}_T \overset{\wedge}{=} \boldsymbol{\nabla}^{\mathrm{Gr}}_T$ *on* $\mathrm{Gr}(\,\mathrm{END})$. *Restricted to the faces* tf *and* tff *this has the form*

(a) $(\boldsymbol{\nabla}_T u)|_{\mathrm{tf,Gr}} = (\boldsymbol{\nabla}_T - \frac{1}{2}P_{\mathrm{tf}}\nabla.\mathbb{K}(T,\cdot))(u|_{\mathrm{tf,Gr}}) =: \nabla^{\mathcal{N}\Lambda(\mathrm{tf})}_T(u|_{\mathrm{tf,Gr}})$

(b) $(\boldsymbol{\nabla}_T u)|_{\mathrm{tff,Gr}} = (\boldsymbol{\nabla}_T - P_{\mathrm{tff}}\mathbb{K}(T,\cdot) - \frac{1}{2}P_{\mathrm{tff}}\nabla.\mathbb{K}(T,\cdot))(u|_{\mathrm{tff,Gr}}) = \nabla^{\mathcal{N}\Lambda(\mathrm{tff})}_T(u|_{\mathrm{tff,Gr}})$

(c) $(\boldsymbol{\nabla}_T u)|_{\mathrm{tff}\cap\mathrm{tf}\mathrm{Gr}} = (\boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}_T - \frac{1}{2}P_{\mathrm{tf}}\nabla^{\mathcal{N}\Lambda(\mathrm{tff})}_{.}\mathbb{K}^{\mathcal{N}\Lambda(\mathrm{tff})}(T,\cdot))(u|_{\mathrm{tff}\cap\mathrm{tf,Gr}})$

Proof. As in Proposition 5.6 this follows from the commutator identity in Lemma 5.7 and the properties of the curvature described in Lemma 5.8. In addition, one has to verify that the term $\nabla.\mathbb{K}(T,\cdot)$ (is of order 2 and) preserves $\mathrm{Cl}(x^bT^*X)$ at tf! This will follow from the more concrete calculation of this term in Proposition 5.16 below. Note also that $\mathbb{K}(T,\cdot)$ *vanishes* at tf. ■

5.2 The Rescaled Calculus and Normal Operators

Using the rescaled heat bundle Gr(END) we can introduce the corresponding "coefficient bundle"

$$\mathrm{CB}_G := \mathrm{Gr}(\,\mathrm{END}) \otimes \mathrm{KD}_H.$$

It is then straightforward to define the rescaled heat calculus as

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X,E) := \rho^l_{\mathrm{tf}}\, \rho^m_{\mathrm{tff}}\, \rho^{p-v}_{\mathrm{hbf}}\ \dot{C}^\infty_{\mathrm{tf,tff,hbf}}(X^2_H, \mathrm{CB}_G),$$

for $l > 0, m \geq 0, p \geq -v$. For $l = 0$, we set

$$\Psi^{0,m,p}_{G,\mathrm{cl}}(X,E) := \rho^m_{\mathrm{tff}}\, \rho^{p-v}_{\mathrm{hbf}}\ \dot{C}^\infty_{\mathrm{tf,tff,hbf}}(X^2_H, \mathrm{CB}_G)$$

and we remind the reader that $A \in \dot{C}^\infty_{\mathrm{tf,tff,hbf}}(X^2_H, \mathrm{CB}_G)^{\mathrm{tf}}$ satisfies the mean value condition

$$\int_{{}^\phi TX/X} \boldsymbol{\iota}(t\frac{\partial}{\partial t})[A|_{\mathrm{tf}}]^0 = 0.$$

The notation $[]^0$ is meant to indicate the order 0-part of the restriction. As usual, there are definitions for the extended (rescaled) calculi.

The rescaled heat calculus is a refinement of the usual heat calulus in the sense that we have an inclusion of the spaces of sections

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X,E) \subset \Psi^{l,m,p}_{H,\mathrm{cl}}(X,E).$$

Therefore the action on sections and the composition of rescaled operators is well defined. It is important to know, however, that the composition of rescaled operators is again a rescaled operator. It is an advantage of our invariant definition of $\mathcal{D}$ that it allows for a relatively easy proof of the following refinement of Theorem 4.6 for rescaled operators:

Theorem 5.12 (Rescaled Composition Formula) *Let $R_{\mathrm{tf}}, L_{\mathrm{tf}} \geq 0$. Then composition gives a map*

$$\Psi^{\mathcal{L}}_G(X) \times \Psi^{\mathcal{R}}_G(X) \xrightarrow{\circ} \Psi^{\mathcal{M}}_G(X)$$

with index set given by

$$M_{\mathrm{hbf}} = (L_{\mathrm{hbf}} + R_{\mathrm{hbf}})\overline{\cup}(L_{\mathrm{tff}} + R_{\mathrm{hbf}})\overline{\cup}(L_{\mathrm{hbf}} + R_{\mathrm{tff}})$$

$$M_{\mathrm{tff}} = L_{\mathrm{tff}} + R_{\mathrm{tff}}, \qquad M_{\mathrm{tf}} = L_{\mathrm{tf}} + R_{\mathrm{tf}}$$

Proof. We do not give the details of this proof here. Basically, the bundle END(E) on X^2_H can be lifted to X^3_H from the left and the right, using $\pi_{H,L}$, $\pi_{H,R}$. Composition of endomorphisms gives a well defined map

$$\pi^*_{H,L}\ \mathrm{END}(E) \otimes \pi^*_{H,R}\ \mathrm{END}(E) \longrightarrow \pi^*_{H,M}\ \mathrm{END}(E).$$

The connections used in the definition of the rescaling can be lifted to X^3_H in the same way. Using compatibility of these connections with this product, as well as with the pullback and pushforward operations, the proof then boils down to Leibniz' rule. ■

Normal operators for our rescaled heat calculus can now be defined in complete analogy with Section 4.3. For $A \in \Psi^{l,m,p}_{G,\mathrm{cl}}(X)$ define the normal operators of A at the faces tf and tff by

$$\begin{aligned} N_{G,\mathrm{tf},l>0}(A) &:= t^{1-l/2}A|_{\mathrm{tf},\mathrm{Gr}} \in \rho_{\mathrm{tff}}^{m-l}\, \dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{tf}, t\mathrm{KD}_H \otimes \Lambda_{\mathrm{tf}}) \\ N_{G,\mathrm{tf},0}(A \oplus c) &:= tA|_{\mathrm{tf},\mathrm{Gr}} \oplus c \in \rho_{\mathrm{tff}}^{m}\, \dot{C}^{\infty}_{\mathrm{tff}}(\mathrm{tf}, t\mathrm{KD}_H \otimes \Lambda_{\mathrm{tf}})^{\mathrm{tf}} \oplus \mathbf{C} \\ N_{G,\mathrm{tff},m}(A) &:= (x')^{-m}A|_{\mathrm{tff},\mathrm{Gr}} \in \rho_{\mathrm{tf}}^{l}\, \rho_{\mathrm{hbf}}^{p-v-m}\, \dot{C}^{\infty}_{\mathrm{tf},\mathrm{hbf}}(\mathrm{tff}, \mathrm{KD}_H \otimes \mathrm{Gr}_{\mathrm{tf}}(\Lambda_{\mathrm{tff}})). \end{aligned}$$

Our aim is to construct the heat kernel for D^2 in the rescaled calculus. This means that we have to describe the action of $t\mathsf{D}^2$ and $t\frac{\partial}{\partial t}$ on $\mathcal{D}$ at the faces tf and tff. To see what this requires, consider the Lichnerowicz formula for D^d

$$t(\mathsf{D}^d)^2 \triangleq t\beta_L^* g_d^{-1}(\boldsymbol{\nabla}, \boldsymbol{\nabla}) + t\beta_L^* \frac{\kappa_X^d}{4} + t\mathbf{c}_{d,L}(\beta_L^* F^{E/S,d}), \tag{95}$$

where the notation is meant to emphasize that we are looking at the operator lifted to X^2_H from the left. Thus we will have to calculate the normal operators for

$$t\frac{\partial}{\partial t}, \quad \boldsymbol{\nabla}, \quad t\beta_L^* g_d^{-1}, \quad t^{1/2}\mathbf{c}_{d,L}, \quad t\kappa_X^d, \quad t\mathbf{c}_{d,L}(\beta_L^* F^{E/S,d}). \tag{96}$$

As in Proposition 5.11 we will show that all of these operators map $\mathcal{D}$ to $\mathcal{D}$ in an appropriate sense by calculating their commutators with ∇^l.

Let us start with the analysis of the action of $\boldsymbol{\nabla}$. The main point is to calculate the curvature terms in Proposition 5.11. However, instead of considering the action of general sections in ${}^bT_{\mathrm{tff}/Y,\mathrm{tf}/X}X^2_H$ on $\mathcal{D}$ we now restrict to the special class of sections of the type

$$\mathcal{W} := \langle t^{1/2}\beta_{H,L}^{*}\,{}^{\mathrm{d}}TX, t\frac{\partial}{\partial t}\rangle = \langle \tau^{1/2}\beta_{H,L}^{*}\,{}^{\phi}TX, t\frac{\partial}{\partial t}\rangle. \tag{97}$$

The essential property of sections in $\mathcal{W}$ is that they are tangent to the fibres in tf$/X$ and tff$/Y$. At tff this is also true for sections in $\beta_{H,L}^{*}\,{}^{\phi}TX$, which is why we will occasionaly drop the factor $\tau^{1/2}$ on the RHS in (97). Restricting to the class $\mathcal{W}$ allows us to make sense of the formulas in Proposition 5.11 on all of X^2_H and to follow the conventions described in Remark 5.5 more consistently. Thus, allowing from now on the shorthand $\beta_L \sim \beta_{H,L}$, we introduce the following notation for our connections:

- ∇ describes the action of TX^2_H on $\mathcal{W}$ and $\mathrm{END}(E)$ and represents the connection used in the definition of $\mathcal{D}$.

  $$\nabla : \Gamma(\mathcal{W}) \to \Gamma(T^*X^2_H \otimes \mathcal{W}) \text{ with } \nabla_N t^{1/2}\beta_L^* V = t^{1/2}\beta_L^* \nabla^d \beta_L^* V \quad \text{and} \quad \nabla_N t\frac{\partial}{\partial t} = 0.$$

- $\boldsymbol{\nabla}$ describes the action of $\mathcal{W}$ on TX^2_H and $\mathrm{END}(E)$. It represents the connection used in the definition of our operators acting on $\mathcal{D}$.

  $$\begin{aligned} &\boldsymbol{\nabla} : \Gamma(TX^2_H) \to \Gamma(\mathcal{W}^* \otimes TX^2_H) \text{ with} \\ &\boldsymbol{\nabla}_{t\frac{\partial}{\partial t}} N = [t\frac{\partial}{\partial t}, N] \text{ and } \boldsymbol{\nabla}_{t^{1/2}\beta_L^* V} N = t^{1/2}\, \beta_L^* \nabla^d_N\, \beta_L^* V + [t^{1/2}\beta_L^* V, N] \end{aligned} \tag{98}$$

It is easy to see that these definitions are compatible with the definition (92) of $\nabla^{N\mathrm{tff}}$ in the last Section. Note that the (b-) connections on $N\mathrm{tf}$ and $N\mathrm{tff}$ are curvature-free by the Jacobi identity. Also in some cases they are rather easy to calculate:

Lemma 5.13 *Let $\alpha \in \Gamma(T^*X)$, $[\phi^*\omega] \in \Gamma([H^*\partial X])$, let $W \in \Gamma({}^\phi TX)$ and $A \in \Gamma([{}^\phi N\partial X])$. Then $P_{\mathrm{tf}}\beta_L^*\alpha \in N^*\mathrm{tf}$, $P_{\mathrm{tff}}\beta_L^*[\phi^*\omega] \in N^*\mathrm{tff}$ and*

(a) $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{\tau^{1/2}\beta_L^* W}(P_{\mathrm{tf}}\beta_L^*\alpha) = \alpha(W)\,d\tau^{1/2}$, *and* $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{t\frac{\partial}{\partial t}}(P_{\mathrm{tf}}\beta_L^*\alpha) = 0$

(b) $\boldsymbol{\nabla}^{N^*\mathrm{tff}}_{\beta_L^* A}(P_{\mathrm{tff}}\beta_L^*[\phi^*\omega]) = \phi^*\omega(A/x)\,dx$, *and* $\boldsymbol{\nabla}^{N^*\mathrm{tff}}_{t\frac{\partial}{\partial t}}(P_{\mathrm{tff}}\beta_L^*\alpha) = 0$

(c) $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{\tau^{1/2}\beta_L^* W}\,d\tau^{1/2} = 0$ *and* $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{t\frac{\partial}{\partial t}}\,d\tau^{1/2} = \frac{1}{2}\,d\tau^{1/2}$

(d) $\boldsymbol{\nabla}^{N^*\mathrm{tff}}_{\beta_L^* W}\,dx = 0$ *and* $\boldsymbol{\nabla}^{N^*\mathrm{tff}}_{t\frac{\partial}{\partial t}}\,dx = 0.$

Proof. As an example, let us do the first part in (a) (leaving out the ubiquitous β_L^*):

$$\begin{aligned}\boldsymbol{\nabla}_{\tau^{1/2}W}\alpha\,(N) &\overset{(98)}{=} \tau^{1/2}W\,\alpha(N) - \alpha\left(\tau^{1/2}x\nabla^d_N\frac{1}{x}W + [\tau^{1/2}W, N]\right)\\ &\overset{(*)}{=} d\alpha(\tau^{1/2}W, N) + N\,\alpha(\tau^{1/2}W) \overset{(**)}{=} d\tau^{1/2}(N)\alpha(W),\end{aligned}$$

where, in $(*)$ and $(**)$, we have used the fact that $\tau^{1/2}$ vanishes at the face tf. ■

In order to make the calculation of normal operators look as invariant as possible, we will consider rescaled "calculi" with coefficients in $\mathcal{W}$ and $\mathcal{W}^*$, the definition proceeding just as before. Also write $\mathbb{K} = [\boldsymbol{\nabla}, \nabla]$, which is the restriction of the curvature K to $\Gamma(\mathcal{W}^* \otimes T^*X_H^2 \otimes \mathrm{END}(E))$, and similarly $\mathbf{K} = [\boldsymbol{\nabla}, \boldsymbol{\nabla}]$. We can now restate Proposition 5.11 as a refinement of Proposition 4.7:

Proposition 5.14 *The maps*

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X) \xrightarrow{\boldsymbol{\nabla}} \Psi^{l,m,p}_{G,\mathrm{cl}}(X, \mathcal{W}^*), \qquad \Psi^{l,m,p}_{G,\mathrm{cl}}(X) \xrightarrow{\frac{\partial}{\partial t}} \Psi^{l-2,m-2,p}_{G,\mathrm{cl}}(X)$$

have the following general form when restricted to the different faces

(a) $N_{G,\mathrm{tf},l}(\boldsymbol{\nabla}A) = (\boldsymbol{\nabla} - \frac{1}{2}P_{\mathrm{tf}}\nabla\mathbb{K})N_{G,\mathrm{tf},l}(A) = \boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tf})}N_{G,\mathrm{tf},l}(A)$

(b) $N_{G,\mathrm{tff},m}(\boldsymbol{\nabla}A) = (\boldsymbol{\nabla} - P_{\mathrm{tff}}\mathbb{K} - \frac{1}{2}P_{\mathrm{tff}}\nabla\mathbb{K})N_{G,\mathrm{tff},m}(A) = \boldsymbol{\nabla}^{\mathcal{N}\Lambda(\mathrm{tff})}N_{G,\mathrm{tff},m}(A)$

(c) $N_{G,\mathrm{tf},l-2}(\frac{\partial}{\partial t}A) = \begin{cases} -\frac{1}{2}(L_{R^\phi} + 2 - l - \mathsf{N})N_{G,\mathrm{tf},l}(A) & \textit{for} \quad l > 2 \\ -\frac{1}{2}(L_{R^\phi} + 2 - l - \mathsf{N})N_{G,\mathrm{tf},2}(A) \oplus -\int_{{}^\phi TX/X} N_{H,\mathrm{tf},2}(A) & \textit{for} \quad l = 2 \end{cases}$

(d) $N_{G,\mathrm{tff},m-2}(\frac{\partial}{\partial t}A) = L_{\frac{\partial}{\partial \tau}}N_{G,\mathrm{tff},m}(A)$

Here N denotes the number operator in Λ_{tf}. Note also that $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{R^\phi}\,d\tau^{1/2} = 0$.

Proof. This is really just a reformulation of Proposition 5.11. For (c) and (d) just recall that we now also have to consider the action of $\boldsymbol{\nabla}_{t\frac{\partial}{\partial t}}$ on $N^*\mathrm{tf}$ and $N^*\mathrm{tff}$, which is 1/2 on $N^*\mathrm{tf}$ and 0 on $N^*\mathrm{tff}$ according to Lemma 5.13. ■

For the calculation of these curvature terms it will be useful to list some further properties of the curvature $\mathbf{K}$ at the faces tf and tff. It has been shown in Lemmas 5.2(a) and 5.8(a) that

$$K(T_{\mathrm{tf}},\,\cdot\,)|_{\mathrm{tf}} \in N^*\mathrm{tf} \otimes \mathrm{Cl}^2_{\mathrm{tf}} \,/\, \mathrm{Cl}^1_{\mathrm{tf}} \quad \text{and} \quad K(T_{\mathrm{tff}},\,\cdot\,)|_{\mathrm{tff}} \in N^*\mathrm{tff} \otimes \mathrm{Cl}^2_{\mathrm{tff}} \,/\, \mathrm{Cl}^1_{\mathrm{tff}} \tag{99}$$

are well defined for vectors T_{tf}, T_{tff} tangent to tf, tff. Using the identifications (88,89) we now show a more precise result:

Lemma 5.15 (More Properties of the Curvature) *For the faces $F = \mathrm{tf}, \mathrm{tff}$, let T_F, N_F be vector fields on X^2_H tangent respectively normal to F. Also, we assume that T_F are lifted from vector fields on $X^2_b \times [0,\infty[$ tangent to the blown down faces F.*

(a) $K(T_{\mathrm{tf}}, N_{\mathrm{tf}})|_{A\in {}^\phi TX} = \, d\tau^{1/2}(N_{\mathrm{tf}})K(T_{\mathrm{tf}}, \beta_L^* A) \quad$ *modulo* $\mathrm{Cl}^1_{\mathrm{tf}}$

(b) $K(T_{\mathrm{tff}}, N_{\mathrm{tff}})|_{B\in {}^\phi N\partial X} = \, dx(N_{\mathrm{tff}})K(T_{\mathrm{tff}}, \frac{1}{x}\beta_L^*[B]) \quad$ *modulo* $\mathrm{Cl}^1_{\mathrm{tff}}$.

Here $[B]$ denotes a choice of extension of B to $[{}^\phi N\partial X]$.

Proof. The proofs of (a) and (b) follow the same pattern. To prove (b), note that

$$K(T_{\mathrm{tff}}, N_{\mathrm{tff}})|_{0\in {}^\phi N\partial X} \in \mathrm{Cl}^1_{\mathrm{tff}}\,. \tag{100}$$

This can be seen by looking at the map π flipping sides in X^2_H. Since, by assumption, the vector T_{tff} is lifted from a vector on $X^2_b \times [0,\infty[$, tangent to $F_\phi \times \{0\}$, we can restrict ourselves to the following situation:

$$\pi(N_{\mathrm{tff}}) = N_{\mathrm{tff}} \ \text{ and } \ \pi(T_{\mathrm{tff}}) = T_{\mathrm{tff}} + VF_\phi \quad \text{at the zero-section of } \mathrm{tff} \to F_\phi.$$

Using (b) above, we get

$$K(T_{\mathrm{tff}}, N_{\mathrm{tff}}) = -K(\pi(T_{\mathrm{tff}}), \pi(N_{\mathrm{tff}})) \in -K(T_{\mathrm{tff}}, N_{\mathrm{tff}}) + \mathrm{Cl}^1_{\mathrm{tff}},$$

which proves (100). It now suffices to calculate the derivative of $K(T_{\mathrm{tff}},\,\cdot\,)$ in Λ^2_{tff} along the fibres of tff. But from (90), the assumption on T_{tff} and Corollary 1.7, this is just a linear combination of sections of the type

$$\beta_L^*[\phi^*\omega] \otimes \beta_L^*\xi, \quad \beta_R^*[\phi^*\omega] \otimes \beta_R^*\xi, \quad \text{with} \quad [\phi^*\omega] \in \Gamma(X, [H^*\partial X]), \quad \xi \in \Gamma(X, \mathrm{End}(E))$$

Thus for $B \in {}^\phi N\partial X$ representing a point in the fibre, we can use Lemma 5.13(b) (and its analogue for sections lifted from the right) to get

$$(\boldsymbol{\nabla}_{\beta_L^* B} K(T_{\mathrm{tff}},\,\cdot\,))(N_{\mathrm{tff}}) = \, dx(N_{\mathrm{tff}})K(T_{\mathrm{tff}}, \frac{1}{x}[\beta_L^* B]),$$

which proves the claim. ■

The curvature terms in $N_{G,\mathrm{tf}}(\boldsymbol{\nabla})$, $N_{G,\mathrm{tff}}(\boldsymbol{\nabla})$ can now be calculated. Readers, who feel uneasy about the signs or the different factors of x in the formulas should (hope that the author has made no mistake and) consult Definition 1.12 and Appendix A.1.

Proposition 5.16 (a) *Using the identification (88) we have for* $A \in {}^{\phi}TX$

$$(P_{\mathrm{tf}}\nabla\mathbb{K})(\cdot, \tau^{1/2}\beta_L^* A, \cdot)|_{B\in {}^{\phi}TX} = (\, d\tau^{1/2})^2 \frac{1}{2}\, \boldsymbol{\varepsilon}_d[\langle x^2 R^d(\cdot,\cdot)(B + \nu_{\mathrm{tf}}|_0), A\rangle_\phi]$$
$$=: (\, d\tau^{1/2})^2 \left(\frac{1}{2}\, \boldsymbol{\varepsilon}_d[\langle x^2 R^d(\cdot,\cdot)B, A\rangle_\phi] + Q_{\mathrm{tf}}(A)\right) \in \Lambda^2(\mathrm{tf}).$$

Here $\nu_{\mathrm{tf}}|_0$ *is the vector field* $P_{\mathrm{tf}}/\, d\tau^{1/2}$ *at the zero section in* ${}^{\phi}TX$*.*

(b) *Using the identification of (89) we find for* $W \in {}^{\phi}TX$

$$(P_{\mathrm{tff}}\mathbb{K})(\beta_L^* W, \cdot)|_{B\in {}^{\phi}N\partial X} = -\frac{1}{2}\, dx(\boldsymbol{\varepsilon}_d, \mathbf{c}_d)(\langle x S_d(W)\cdot, \cdot\rangle_\phi)$$
$$= -\frac{1}{2}\, dx\left[(\boldsymbol{\varepsilon}_d, \mathbf{c}_d)\langle S_\phi(W)\cdot, \cdot\rangle_d + \boldsymbol{\varepsilon}(\frac{dx}{x})\mathbf{c}_d(W/x)\right] \in \Lambda^1(\mathrm{tff})$$

covariant constant along the fibres tff/F_ϕ*.*

(c) *Let* $A \in \Gamma({}^{\phi}N\partial X)$*. Then*

$$(P_{\mathrm{tff}}\nabla\mathbb{K})(\cdot, \beta_L^* A, \cdot)|_{B\in {}^{\phi}N\partial X} = (\, dx)^2 \frac{1}{2}\, \boldsymbol{\varepsilon}_d[\langle \phi^* R^{B,b}(\cdot,\cdot)([B] + x\nu_{\mathrm{tff}}|_0), [A]\rangle_\phi]$$
$$=: (\, dx)^2 \left(\frac{1}{2}\, \boldsymbol{\varepsilon}_d[\langle \phi^* R^{B,b}(\cdot,\cdot)[B], [A]\rangle_\phi] + Q_{\mathrm{tff}}(A)\right) \in \Lambda^2(\mathrm{tff}).$$

Here, $\nu_{\mathrm{tff}}|_0$*, is the vector field* $P_{\mathrm{tff}}/\, dx$ *evaluated at the zero section in* $\mathrm{tff} \to F_\phi$ *and* $[A], [B]$ *denote choices of extension of* A, B *into* $[{}^{\phi}N\partial X]$*. The result is clearly independent of that choice.*

(d) *Now let* $V \in \Gamma(V\partial X)$*. Then*

$$\begin{aligned}(P_{\mathrm{tff}}\nabla\mathbb{K})(\cdot, \beta_L^* V, \cdot)_{B\in {}^{\phi}N\partial X} &= \frac{1}{4}(\, dx)^2(\boldsymbol{\varepsilon}_d, \boldsymbol{\varepsilon}_d)\langle x^2\Omega_\phi(\cdot,\cdot), V\rangle_\phi \\ &= \frac{1}{4}(\, dx)^2(\boldsymbol{\varepsilon}_d, \boldsymbol{\varepsilon}_d)\langle \Omega_M(\cdot,\cdot), V\rangle_{M,b}] \in \Lambda^2(\mathrm{tff}).\end{aligned}$$

Again, this term is covariant constant along the fibres tff/F_ϕ*.*

Note that, except for the term in (b), none of these terms contain a $\boldsymbol{\varepsilon}(\frac{dx}{x})$*-contribution.*

Proof. The proof of (a) at tf is easy. First, recall again that

$$\mathbb{K}(\tau^{1/2}\beta_L^* A, N) = \tau^{1/2} K(\beta_L^* A, N)$$

vanishes at tf since $\tau^{1/2}$ vanishes there. Thus

$$(P_{\mathrm{tf}}\nabla\mathbb{K})(N, \tau^{1/2}\beta_L^* A, N) = \, d\tau^{1/2}(N) K(\beta_L^* A, P_{\mathrm{tf}} N),$$

and we can calculate the derivative along the fibre as in the proof of Lemmas 5.8(a) and 5.15. In this case, however, the constant term along the fibres does not vanish, accounting for the term $Q_{\mathrm{tff}}(A)$ in the formula. Now using (90) and writing

$$g_d(R^d(A, B)\cdot, \cdot) = g_\phi(x^2 R^d(\cdot,\cdot)B, A),$$

gives the result.

It remains to show that the RHS in (a) really is an element in $\mathrm{Cl}^2_{\mathrm{tf}}$. This should be clear for the linear term in the fibres by Proposition 5.6(c). It is also true for $Q_{\mathrm{tf}}(A)$, since ν_{tf} is tangent to the fibres tff/Y, and can therefore be written in the form $\nu_{\mathrm{tf}} = \beta_L^* C + xN$ for $C \in \Gamma({}^{\phi}TX), N \in \Gamma(TX)$. Then

$$Q_{\mathrm{tf}}(A) = K(\beta_L^* A, (\beta_L^* C + xN)|_{0 \in {}^{\phi}TX}) = xK(\beta_L^* A, N) + \mathrm{Cl}^2_{\mathrm{tf}},$$

but the order-2 part of $K(\beta_L^* A, N)$ is seen to be in $x\Lambda^{2b}T^*X$ in part (b)!

Part (b) follows from (90) and Proposition 1.14 (a). The proof of (c) is completely analogous to (a) once we restrict ourselves to $A, B \in {}^{\phi}N\partial X$ with extensions $[A], [B]$ in $[{}^{\phi}N\partial X]$ as stated. Then again

$$(P_{\mathrm{tff}} \nabla \mathbb{K})(N, \beta_L^* A, N)|_{\mathrm{tff}} = \ dx(N) K(\frac{1}{x}\beta_L^*[A], P_{\mathrm{tff}} N), \tag{101}$$

and the result follows by calculating the derivative along the fibres as in Lemma 5.8(d) and Lemma 5.13. Finally (d): For $V \in V\partial X$, $A_1, A_2 \in {}^{\phi}N\partial X$ the term at tff,

$$\nabla_N \mathbb{K}(N, \beta_L^* V) = \frac{1}{2} \beta_L^* \mathbf{c}_d((\nabla^d_N R^d)(V, N)),$$

was calculated in Proposition 1.14 (b). ■

Having described the role of the conormal bundles $N^*\mathrm{tf}$ and $N^*\mathrm{tff}$, and used them in our calculations, we allow ourselves to assume from now on that

N^*tf is trivialized by $d\tau^{1/2}$ and N^*tff is trivialized by dx ,

to simplify our formulas a bit. Then, recalling Lemma 4.2, the rescaled bundle tf looks like

$$\Lambda(\mathrm{tf}) = \Lambda(x^b T^* X) \otimes \mathrm{END}_{\mathrm{Cl}({}^d T^* X)}(E) \longrightarrow {}^{\phi}TX.$$

Also, $\boldsymbol{\nabla}$ restricts to the fibres of tf → X as the trivial connection (d_{END}, $d_{{}^{\phi}TX/X}$). The rescaled bundle over the *interior* of tff (i.e. without the second rescaling) looks like

$$\Lambda(\mathrm{tff}) = \Lambda(\phi^{*b} T^* B) \otimes \mathrm{END}_{\mathrm{Cl}(\phi^{*b} T^* X)}(E) \longrightarrow {}^{\phi}N\partial X \times_Y \partial X \times]0, \infty[_{\tau^{1/2}}.$$

The induced connection on the fibres tff/Y is trivial along the fibres of p : ${}^{\phi}N\partial X \times]0, \infty[_{\tau^{1/2}} \to \partial X$. One could write it as $(p^* \nabla^{\mathrm{END},d}_{\mathsf{v}\cdot}, p^* \nabla^{{}^{\phi}N\partial X}_{\mathsf{v}\cdot})$.

Corollary 5.17 (Explicit Form of the Induced Rescaled Connection)
Denote by $\mathbf{x}$, $\mathbf{y}$ *the linear variables in the fibres of* ${}^{\phi}TX$ *and* ${}^{\phi}N\partial X$ *respectively. Then for* $A \in {}^{\phi}TX$, $B \in {}^{\phi}N\partial X$ *and* $W \in V\partial X$

(a) $\boldsymbol{\nabla}^{\Lambda(\mathrm{tf})}_A = \boldsymbol{\iota}(A)\, d_{\mathrm{END}} - \frac{1}{4} \boldsymbol{\varepsilon}[\langle x^2 R^d(\cdot,\cdot)\mathbf{x}, A\rangle_{\phi}] - \frac{1}{2} Q_{\mathrm{tf}}(A))$

(b) $$\begin{aligned}\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}_W &= \boldsymbol{\nabla}^{\mathrm{END},d}_{\mathsf{v}W} - \tfrac{1}{2}(\boldsymbol{\varepsilon}_d, \mathsf{v}\mathbf{c}_d)\langle S_{\phi}(W)\cdot, \cdot\rangle_d - \tfrac{1}{4}\boldsymbol{\varepsilon}_d(x^2 \langle \Omega_{\phi}(\cdot,\cdot), W\rangle_{\phi}) + \tfrac{1}{2}\boldsymbol{\varepsilon}_d(\tfrac{dx}{x})\mathbf{c}_d(\tfrac{1}{x}\mathsf{v}W) \\ &=: \widehat{\boldsymbol{\nabla}}^{\Lambda(\mathrm{tff})}_W + \tfrac{1}{2}\boldsymbol{\varepsilon}_d(\tfrac{dx}{x})\mathbf{c}_d(\tfrac{1}{x}\mathsf{v}W)\end{aligned}$$

(c) $\boldsymbol{\nabla}_B^{\Lambda(\mathrm{tff})} = \boldsymbol{\iota}(B)\, d_{\,\mathrm{END}} - \frac{1}{4}\,\boldsymbol{\varepsilon}_d[\langle \phi^* R^{B,b}(\cdot,\cdot)\mathbf{y}, B\rangle_\phi] - \frac{1}{2} Q_{\mathrm{tff}}(B))$. ■

Let us now describe the action of the Clifford action and of the metric in (96).

Proposition 5.18 *The metric gives a map*

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X) \xrightarrow{t\beta_L^* g_d^{-1}} \Psi^{l,m,p}_{G,\mathrm{cl}}(X, \mathcal{W}^* \otimes \mathcal{W}^*)$$

such that, in view of (88, 89), $N_{G,\mathrm{tf}}(tg_d^{-1}) = g^{-1}_{\phi TX/X}$ *and* $N_{G,\mathrm{tff}}(tg_d^{-1}) = \tau g^{-1}_{\phi N\partial X/Y}$. ■

To describe the Clifford action at tf and tff, we use the partial Clifford action m_d on $\Lambda(\mathrm{tff})$, given for $\alpha \in {}^dT^*X$ by

$$\mathsf{m}_d(\alpha) := \mathbf{c}_d(\mathsf{v}\alpha) + \boldsymbol{\varepsilon}_d(\mathsf{n}\alpha).$$

Also, we recall that the map $\delta_h^* : x^bT^*X|_{\partial X} \to {}^dT^*X|_{\partial X}$ was described in Lemma 1.9.

Proposition 5.19 *The (left) Clifford action of an element* $x\alpha \in \Gamma(x^bT^*X) = \Gamma([{}^dV^*\partial X])$ *gives a map*

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X) \xrightarrow{\tau^{1/2}\beta_L^*\mathbf{c}_d(x\alpha)} \Psi^{l,m,p}_{G,\mathrm{cl}}(X).$$

It restricts to the faces as

(a) $N_{G,\mathrm{tf},l}(\tau^{1/2}\beta_L^*\mathbf{c}_d(x\alpha)A) = d\tau^{1/2}\,\boldsymbol{\varepsilon}(x\alpha) \circ N_{G,\mathrm{tf},l}(A) \mathrel{\hat{=}} \boldsymbol{\varepsilon}(x\alpha) \circ N_{G,\mathrm{tf},l}(A)$

(b) $N_{G,\mathrm{tff},m}(\tau^{1/2}\beta_L^*\mathbf{c}_d(x\alpha)A) = \tau^{1/2}(\mathbf{c}_d(x\mathsf{v}\alpha) + dx\,\boldsymbol{\varepsilon}(\mathsf{h}\alpha)) \circ N_{G,\mathrm{tff},m}(A)$
$\mathrel{\hat{=}} \tau^{1/2}\mathsf{m}_d(\delta_h(x\alpha)) \circ N_{G,\mathrm{tff},m}(A)$.

Proof. Part (a) follows from the fact that $\tau^{1/2}\beta_L^*\mathbf{c}_d(x\alpha)$ vanishes at tf and

$$\nabla_N^{\mathrm{End},d}\tau^{1/2}\beta_L^*\mathbf{c}_d(x\alpha)|_{\mathrm{tf}} = d\tau^{1/2}(N)\beta_L^*\mathbf{c}_d(x\alpha)|_{\mathrm{tf}} + \tau^{1/2}\nabla^{\mathrm{End},d}\beta_L^*\mathbf{c}_d(x\alpha)|_{\mathrm{tf}},$$

where the last summand again vanishes at tf. To prove (b), first note that $\tau^{1/2}\mathbf{c}_d(x\alpha)$ restricts as $\tau^{1/2}\mathbf{c}_d(x\mathsf{v}\alpha)$ at tff. The first order contribution of $\tau^{1/2}\mathbf{c}_d(x\alpha)$ at tff can be calculated using Lemma 1.11

$$\nabla_N^{\mathrm{End},d}\tau^{1/2}\mathbf{c}_{d,L}(x\alpha) = \tau^{1/2}\mathbf{c}_{d,L}(\nabla_N^d x\alpha) = \tau^{1/2}\,dx(N)\mathbf{c}_{d,L}(\mathsf{h}\alpha),$$

as claimed. ■

A slightly different formulation of the last result will also be of some use:

Corollary 5.20 *At* tf, *the rescaled restriction to the corner is given by*

(a) $|_{\mathrm{tff,Gr}} = \mathsf{m}_d \circ \delta_h : \mathrm{Cl}(x^bT^*X) \longrightarrow \mathrm{Cl}({}^dV^*\partial X) \otimes \Lambda\phi^{*b}T^*B|_{\partial X}$ *in* $\mathrm{Gr}_{\mathrm{tff}}(\,\mathrm{END})$,

(b) $|_{\mathrm{tff,Gr}} = \delta_h : \Lambda(x^bT^*X) \longrightarrow \Lambda^dT^*X|_{\partial X}$ *in* $\mathrm{Gr}(\,\mathrm{END})$. ■

We are left with the analysis of the behavior of the scalar curvature term $t\kappa_X$ in (96). Now near ∂X we can write, using Corollary 1.7

$$\begin{aligned} t\kappa_X &= t\ \mathrm{tr}_{aa'}(\langle\langle (g_d^{-1})_{bb'}, {}_aR^d(\cdot_{a'},\cdot_b)_{b'}\rangle\rangle) \\ &= \tau\ \mathrm{tr}_{aa'}(\langle\langle (g_\phi^{-1})_{bb'}, {}_a\mathsf{v}R^\phi(\cdot_{a'},\cdot_b)\mathsf{v}_{b'}\rangle\rangle) + \tau\ \mathrm{tr}_{aa'}(\langle\langle (g_\phi^{-1})_{bb'}, {}_a\phi^*R^{B,b}(\cdot_{a'},\cdot_b)_{b'}\rangle\rangle). \end{aligned} \tag{102}$$

This is a C^∞-function on X^2_H, which vanishes at tf. At tff the second summand on the RHS vanishes, since the curvature $\phi^*R^{B,b}$ vanishes when evaluated on a vertical vector. Since the B_ϕ-term in (20) also vanishes on vertical vectors, we are left with

$$t\kappa_X|_{\mathrm{tf}} = 0, \qquad t\kappa_X|_{\mathrm{tff}} = \tau\kappa_{M/Y}.$$

Putting everything together, we have found the following description of D^2 at the different faces:

Corollary 5.21 *The square of the Dirac operator gives a map*

$$\Psi^{l,m,p}_{G,\mathrm{cl}}(X) \xrightarrow{t\mathsf{D}^2} \Psi^{l,m,p}_{G,\mathrm{cl}}(X)$$

with

(a) $N_{G,\mathrm{tf}}(t\mathsf{D}^2) = N_{G,\mathrm{tf}}(t(\mathsf{D}^d)^2) \triangleq g^{-1}_{\phi TX/X}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tf})}, \boldsymbol{\nabla}^{\Lambda(\mathrm{tf})}) + \boldsymbol{\varepsilon}_d(x^2F^{E/S,d}(\cdot,\cdot)) \qquad =: \mathcal{H}_X$

(b) $N_{G,\mathrm{tff}}(t\mathsf{D}^2) = N_{G,\mathrm{tff}}(t(\mathsf{D}^d)^2)$

$$\begin{aligned} &= \tau g^{-1}_{\phi N\partial X/\partial X}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}, \boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}) && =: \tau\mathcal{H}_B \\ &\quad + \tau\left(g^{-1}_{M/Y}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}, \boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}) + \tfrac{\kappa_{M/Y}}{4} + \mathsf{m}_d(\delta_h(x^2F^{E/S,d}(\cdot,\cdot)))\right) && =: \tau\mathbf{B}^2 \end{aligned}$$

Remark 5.22 Using the definition of $\widehat{\boldsymbol{\nabla}}^{\Lambda(\mathrm{tff})}$ given in Corollary 5.17 we set

$$\widehat{\mathbf{B}}^2 := g^{-1}_{M/Y}(\widehat{\boldsymbol{\nabla}}^{\Lambda(\mathrm{tff})}, \widehat{\boldsymbol{\nabla}}^{\Lambda(\mathrm{tff})}) + \frac{\kappa_{M/Y}}{4} + \mathsf{m}_d(\delta_h(x^2F^{E/S,d}(\cdot,\cdot))).$$

Then $\widehat{\mathbf{B}}^2$ (truly) commutes with $\boldsymbol{\varepsilon}(\frac{dx}{x})$ and we can write $\mathbf{B}^2$ as

$$\widehat{\mathbf{B}}^2 + \boldsymbol{\varepsilon}(\frac{dx}{x})\left[\mathsf{D}^{\phi,V} - \frac{1}{2}(\boldsymbol{\varepsilon}_d, \boldsymbol{\varepsilon}_d, \mathbf{c}_d)\langle xB_\phi(\cdot,\cdot),\cdot\rangle_d - \frac{1}{8}(\boldsymbol{\varepsilon}_d, \boldsymbol{\varepsilon}_d, \mathbf{c}_d)\langle x\Omega_\phi(\cdot,\cdot),\cdot\rangle_d\right]. \tag{103}$$

Recall that our model space ${}^\phi N\partial X \to Y$ can be identified with ${}^bHM \to Y$ using the map δ_h. Also the Clifford algebra $\mathrm{Cl}({}^dT^*X|_{\partial X})$ will be identified with $\mathrm{Cl}({}^bT^*X|_M)$ in that way. Using Definition 1.12 and Lemma 1.13 we find

$$\Omega_M = x^2\Omega_\phi, \quad \text{"}xS_M = S_\phi + \frac{1}{2}B_\phi\text{"}, \quad R^Y = \phi^*R^{B,b}, \quad \mathsf{m} = \mathsf{m}_d \circ \delta_h \circ x.$$

In the case $B_\phi = 0$, for instance if g_ϕ is a product near the boundary, we thus have $S_\phi = xS_M$ and $\widehat{\mathbf{B}}^2 = \mathbf{A}^2_M$, i.e. (103) is of the classical form.

Proof. These are direct consequences of the Lichnerowicz formula (95) for the Dirac operator D^d and the calculations made in the above series of Propositions, especially 5.14 and 5.16. To see that we can use the Dirac operator D^d instead of D, note that by the Lichnerowicz formula and Lemma 1.6(c) the difference of their squares:

$$\begin{aligned} t(\mathsf{D}^2 - (\mathsf{D}^d)^2) &= t x^{v/2} (\mathsf{D}^d)^2 x^{-v/2} - t(\mathsf{D}^d)^2 = t g_d^{-1}(x^{v/2} \boldsymbol{\nabla} x^{-v/2}, x^{v/2} \boldsymbol{\nabla} x^{-v/2}) - t g_d^{-1}(\boldsymbol{\nabla}, \boldsymbol{\nabla}) \\ &= -vt\boldsymbol{\nabla}_{X_d} + t\frac{v^2}{4} - t\frac{v}{2}\langle g_d^{-1}, (\boldsymbol{\nabla}\frac{dx}{x})\rangle = -vt\boldsymbol{\nabla}_{X_d} - \frac{v^2}{4}t + O(x,t), \end{aligned}$$

vanishes at tff and tf. ■

5.3 The Rescaled Heat Kernel and the Index

The aim of this Section is to prove an extension of Theorem 4.9 in the rescaled heat calculus. We start writing down the solutions of the model problems at tf, tff by adapting the results in Appendices A.4 and A.5 to our setting.

Proposition 5.23 (a) *The heat kernel for* $\mathcal{H}_X$

$$\mathsf{K}_{\mathcal{H}_X}(\mathbf{x}, t) := p(x^2 R^d, \mathbf{x}, t) \exp(\frac{1}{2}\langle x^2 R^d \cdot \nu_{\mathrm{tf}}|_0, \mathbf{x}\rangle_\phi) \exp(-x^2 F^{E/S,d})\, dt\, dvol_{^\phi TX/X}$$

is in $\Psi^{2,0}_{G,\mathrm{s,cl}}({}^\phi TX/X, \Lambda(\mathrm{tf}))$ *with normal operator at* $\mathrm{tf}({}^\phi TX) = \mathrm{tf}(X^2_H)$ *given by*

$$N_{G,\mathrm{tf},2}(\mathsf{K}_{\mathcal{H}_X})(\mathbf{x}) = p(x^2 R^d, \mathbf{x}, 1) \exp(\frac{1}{2} Q_{\mathrm{tf}}(\mathbf{x})) \exp(-x^2 F^{E/S,d})\, dt\, dvol_{^\phi TX/X}\,.$$

Recall that $Q_{\mathrm{tf}}(\mathbf{x}) = \langle x^2 R^d \cdot \nu_{\mathrm{tf}}|_0, \mathbf{x}\rangle_\phi$*.*

(b) *The heat kernel for* $\mathcal{H}_B$

$$\mathsf{K}_{\mathcal{H}_B}(\mathbf{y}, \tau) := p(\phi^* R^{B,b}, \mathbf{y}, \tau) \exp(\frac{1}{2} Q_{\mathrm{tff}}(\mathbf{y}))\, d\tau\, dvol_{^\phi N\partial X/\partial X}$$

is in $\Psi^{2,0}_{G,\mathrm{s,cl}}({}^\phi N\partial X/\partial X, \Lambda(\mathrm{tff}))$ *with normal operators at* $\mathrm{tf} = {}^\phi N\partial X$ *and* ti *given by*

$$N'_{G,\mathrm{tf},2}(\mathsf{K}_{\mathcal{H}_B})(\mathbf{y}) = p(\phi^* R^{B,b}, \mathbf{y}, 1) \exp(\frac{1}{2} Q_{\mathrm{tff}}(\mathbf{y}))\, d\tau\, dvol_{^\phi N\partial X/\partial X},$$

$$N'_{H,\mathrm{ti},0}(\mathsf{K}_{\mathcal{H}_B})(\mathbf{y}) = \sqrt{4\pi}^{\,-h-1} \exp(-\|\mathbf{y}\|^2/4)\, d\tau\, dvol_{^\phi N\partial X/\partial X}\,.$$

Recall that $Q_{\mathrm{tff}}(\mathbf{y}) = \langle x\phi^* R^{B,b} \cdot \nu_{\mathrm{tff}}|_0, \mathbf{y}\rangle_\phi$*. Also, the notation* $N'_{G,\mathrm{tf}}$ *is used to remind the reader that these are the normal operators defined using* τ *as the temporal variable as in Appendix A.4. Also, we emphasize that all the normal operators above commute with* $\boldsymbol{\varepsilon}(\frac{dx}{x})$*.*

Proof. We prove (a) using (the generalization to families of) Proposition A.8. The connection $\boldsymbol{\nabla}_A^{\Lambda(\mathrm{tf})}$, for $A \in {}^\phi TX$, appearing in the definition of $\mathcal{H}_X$ was calculated in Corollary 5.17(a):

$$\boldsymbol{\nabla}_A^{\Lambda(\mathrm{tf})} \quad = \quad \boldsymbol{\iota}(A)\, d_{\,\mathrm{END}} - \frac{1}{4} g_{{}^\phi TX/X}(x^2 R^d(\cdot,\cdot)\mathbf{x}, A) - \frac{1}{2} Q_{\mathrm{tf}}(A).$$

This is of the form required in Theorem A.7 with $R = x^2 R^d$ and $q = Q_{\mathrm{tf}}$. The underlying vector space W is any fibre of the bundle ${}^\phi TX \to X$, the metric g_W is given by the fibre metric $g_{{}^\phi TX/X}$ and the coefficient algebra $\mathcal{A}$ is $\Lambda^{\mathrm{ev}}(x\,{}^bT^*X) \otimes \mathrm{END}_{\,\mathrm{Cl}({}^dT^*X)}(E)$. ■

Theorem 5.24 (Rescaled Heat Kernel) *The heat kernel for* D^2 *is an element*

$$\mathsf{K} \in \Psi_G^{\mathcal{H}}(X, E), \qquad \mathcal{H} = (2 + \mathbf{N}, 2 + \mathbf{N}, 0 + H_{\mathrm{hbf}}) \quad \textit{as in} \quad (86).$$

Its rescaled normal operators at the different faces are (well defined and) given by

(a) $N_{G,\mathrm{tf},2}(\mathsf{K})(\mathbf{x}) = p(x^2 R^d, \mathbf{x}, 1) \exp(\frac{1}{2} Q_{\mathrm{tff}}(\mathbf{x})) \exp(-x^2 F^{E/S,d})\; dt\; dvol_{{}^\phi TX/X}$

(b) $N_{G,\mathrm{tff},2}(\mathsf{K})(\mathbf{y}, \tau) = \mathsf{K}_{\mathcal{H}_B}(\mathbf{y}, \tau)/\, d\tau \cdot \mathsf{K}_{\mathbf{B}^2}(\tau)$

$\qquad = p(\phi^* R^{B,b}, \mathbf{y}, \tau) \exp(\frac{1}{2} Q_{\mathrm{tff}}(\mathbf{y}))\; dvol_{{}^\phi N\partial X/\partial X}\, \mathsf{K}_{\mathbf{B}^2}(\tau),$

(c) $N_{H,\mathrm{hbf},0}(\mathsf{K}) = [\Pi_\circ][\exp(-t\mathsf{D}_Y^2)] \frac{1}{\sqrt{4\pi t}} e^{-(\ln(s))^2/4t} \frac{ds}{s}$

Proof. The solutions at the faces patch together because of Corollaries 1.7 and 5.20. We thus get a parametrix which solves the heat equation to first order in the rescaled heat calculus. This is transformed into a parametrix with a smooting remainder using the Composition Formula 5.12. Then a final inversion step as in Section 4.3 is performed. ■

Remark 5.25 As in Theorem 4.11, we can show that $H_{\mathrm{hbf}} = \mathbf{N}$. In the following paragraphs we assume that we have shown the Theorem in this form, if only for reasons of notational simplicity.

The Regularized Supertrace

Let us note what we have achieved w.r.t. the local supertrace of the heat kernel by introducing the rescaling. Denote again by $\triangle_H$ the lift of the diagonal $\triangle_X \times [0, \infty[$ to X_H^2. First, using our notation for blowups we get

$$\triangle_H = [X \times [0, \infty[_{t^{1/2}}, \partial X \times \{0\}].$$

The behavior of general elements in the rescaled heat calculus w.r.t. the local supertrace is now described in parts (c) and (d) of the following

Lemma 5.26 (a) $t\mathrm{KD}_H|_{\triangle_H} = \tau^{-n/2} \beta_{\triangle_H,R}^{*}\,{}^\phi\Omega(X)\, dt = \rho_{\mathrm{hbf}}^{v-1} \rho_{\mathrm{tff}}^{-h-2} \rho_{\mathrm{tf}}^{-n} \beta_{\triangle_H,R}^{*}\Omega(X)\, dt$

(b) $\mathrm{str} : \Psi_H^{2,2,0}(X, E) \longrightarrow \rho_{\mathrm{hbf}}^{-1} \rho_{\mathrm{tff}}^{-h-2} \rho_{\mathrm{tf}}^{-n} C^\infty(\triangle_H, \beta_{\triangle_H,R}^{*}\Omega(X)\, dt)$

(c) $\mathrm{str} : \Psi_G^{2,2,0}(X) \longrightarrow C^\infty(\triangle_H, \beta^*_{\triangle_H,R}{}^b\Omega(X)\,dt) \quad \big(= tC^\infty(\triangle_H, {}^b\Omega(\triangle_H))\big)$

(d) $\beta^*_{\triangle_H,R}{}^b\Omega(X)|_{\mathrm{tf}\cap\triangle_H} \overset{\wedge}{=} {}^b\Omega(X), \qquad \beta^*_{\triangle_H,R}{}^b\Omega(X)|_{\mathrm{tff}\cap\triangle_H} \overset{\wedge}{=} \frac{d\tau^{1/2}}{\tau^{1/2}}\Omega(\partial X) = {}^b\Omega(\mathrm{tff}).$ ■

Of course, an element $L(t) \in C^\infty(\triangle_H, \beta^*_{\triangle_H,R}{}^b\Omega(X))$ will in general not be integrable for fixed $t = T$. We therefore have to consider the *regularized* integral

$$\operatorname*{reg}_{z=0} \int_{X,T} x^z L(T),$$

as in Section 3.8. There we showed that the limit of the regularized supertrace

$$\mathrm{Str}_d(\mathsf{K}(T)) = \operatorname*{reg}_{z=0} \mathrm{Str}(x^z\mathsf{K}(T)) = \operatorname*{reg}_{z=0} \int_{X,T} x^z \ \mathrm{str}(\mathsf{K}(T))/\,dt$$

of the heat kernel at *large* times equals the extended L^2_b-index of D, denoted by $\mathrm{ind}_-(\mathsf{D})$. To analyze the behavior of this supertrace for *small* times we have to improve Lemma 5.26 further:

Lemma 5.27 (Small Time Limit of the Heat Supertrace)

(a) *The local supertrace* $\mathrm{str}(\mathsf{K}(t))/\,dt \in C^\infty(\triangle_H, \beta^*_{\triangle_H,R}{}^b\Omega(X))$ *vanishes in* $\mathrm{tf}\cap\mathrm{tff}$ *and* $\mathrm{tff}\cap\mathrm{hbf}$.

(b) $\lim\limits_{T\to 0} \mathrm{Str}_d(\mathsf{K}(T)) = \lim\limits_{T\to 0} \operatorname*{reg}_{z=0} \int_X x^z \ \mathrm{str}(\mathsf{K}(T))/\,dt = \int_{\mathrm{tff}\cap\triangle_H} \mathrm{str}(\mathsf{K})|_{\mathrm{tff}} + \int_{\mathrm{tf}\cap\triangle_H} \mathrm{str}(\mathsf{K})|_{\mathrm{tf}},$

the RHS being well defined because of (a).

Proof. (a) follows from the fact, proved in Theorem 5.24(a) and (c) (see also the remark in Proposition 5.16), that the restriction of K to these corners contains no contribution of type $\boldsymbol{\varepsilon}_d(\frac{dx}{x})$. Thus the supertrace has to vanish there.

To prove (b) we localize around the corner $\mathrm{tf}\cap\mathrm{tff}$ (the argument at $\mathrm{tff}\cap\mathrm{hbf}$ is even easier), where we can use the coordinates x and $\tau^{1/2}$. Then, using the vanishing at the corner shown in (a), we can calculate the $T \to 0$-limit as follows:

$$\operatorname*{reg}_{z=0} \int x^z a(x, \frac{T^{1/2}}{x})\frac{dx}{x} = \int a(x, \frac{T^{1/2}}{x})\frac{dx}{x} = -\int a(\frac{T^{1/2}}{\tau^{1/2}}, \tau^{1/2})\frac{d\tau^{1/2}}{\tau^{1/2}},$$

for $a \in C_c^\infty([0,\infty[\times[0,\infty[)$ with compact support and $a(0,0) = 0$. We can decompose $a = b + a_1 + a_2$ such that b vanishes at $\{x = 0\} \cup \{\tau^{1/2} = 0\}$, and $a_1 = a(0, \tau^{1/2})$ and $a_2 = a(x, 0)$. Thus

$$\begin{aligned} \operatorname*{reg}_{z=0} \int x^z a(x, \frac{T^{1/2}}{x})\frac{dx}{x} \quad &= \quad \int b(x, \frac{T^{1/2}}{x})\frac{dx}{x} + \int a(x,0)\frac{dx}{x} - \int a(0, \tau^{1/2})\frac{d\tau^{1/2}}{\tau^{1/2}} \\ &\xrightarrow{T\to 0} \int a(x,0)\frac{dx}{x} - \int a(0, \tau^{1/2})\frac{d\tau^{1/2}}{\tau^{1/2}}, \end{aligned}$$

which proves the claim. ■

We can now calculate the (extended L^2_b-) index of D (resp. the extended L^2_d-index of D^d), using an extension of the McKean-Singer formula. Starting from (67) we have

$$\begin{aligned} \mathrm{ind}_-(\mathsf{D}) &= \lim_{T\to\infty} \operatorname*{reg}_{z=0} \mathrm{Str}(x^z \mathsf{K}(T)/\,dt) \\ &= \lim_{T\to 0} \operatorname*{reg}_{z=0} \mathrm{Str}(x^z \mathsf{K}(T)/\,dt) + \int_0^\infty \operatorname*{reg}_{z=0} \frac{\partial}{\partial T} \mathrm{Str}(x^z \mathsf{K}(T)) \end{aligned} \tag{104}$$

For the second summand, we apply the usual trick:

$$\begin{aligned} \frac{\partial}{\partial T} \mathrm{Str}(x^z \mathsf{K}(T)) &= \frac{1}{2} \mathrm{Str}(x^z[\mathsf{D},\mathsf{D}]\mathsf{K}(T)) = \frac{1}{2} \mathrm{Str}(x^z[\mathsf{D},\mathsf{D}\mathsf{K}(T)]) \\ &= \frac{1}{2} \mathrm{Str}([\mathsf{D}, x^z \mathsf{D}\mathsf{K}(T)] - [\mathsf{D}, x^z]\mathsf{D}\mathsf{K}(T)) = -\frac{1}{2} \mathrm{Str}(z x^z \mathbf{c}_d(\frac{dx}{x})\mathsf{D}\mathsf{K}(T)) \end{aligned}$$

The integrand in the second summand in (104) therefore becomes

$$\begin{aligned} -\frac{1}{2} \operatorname*{res}_{z=0} \mathrm{Str}(x^z \mathbf{c}_d(\frac{dx}{x})\mathsf{D}\mathsf{K}(T)) &= -\frac{1}{2} \int_{\partial X} \mathrm{str}(N_{H,\mathrm{hbf},0}(\mathbf{c}_d(\frac{dx}{x})\mathsf{D}\mathsf{K}(T))) \\ &= -\frac{1}{2} \int_{\partial X} \mathrm{tr} \left(\boldsymbol{\epsilon}\, \Pi_\circ \mathbf{c}_d(\frac{dx}{x})[\mathsf{D}_Y + \mathbf{c}_d(\frac{dx}{x}) s \frac{\partial}{\partial s}][\exp(-T\mathsf{D}_Y^2)] \frac{1}{\sqrt{4\pi T}} e^{-(\ln(s)^2)/4T} \right) |_{s=1} \\ &= -\frac{1}{4\sqrt{\pi T}} \int_{\partial X} \mathrm{tr}(\boldsymbol{\epsilon}\, \Pi_\circ \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]) \\ &= -\frac{1}{4\sqrt{\pi T}} \int_{Y} \mathrm{tr}_{\mathcal{K}}(\boldsymbol{\epsilon}\, \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]) =: \frac{1}{2}\eta(\mathsf{D}_Y, \mathcal{K}, T). \end{aligned}$$

This is the usual eta-integrand for the operator D_Y, see for instance [Va]. Note however that the underlying vector bundle $\mathcal{K} \to Y$ is graded by $\boldsymbol{\epsilon}$ and not just $\boldsymbol{\epsilon}_B \overset{\wedge}{=} \boldsymbol{\epsilon}_{{}^bT^*B}$. Also recall from Section 3.5 that D_Y might have a rather complicated structure in general. In any case the eta-term vanishes when $h+1$ is odd, since

$$\begin{aligned} \mathrm{tr}_{\mathcal{K}}(\boldsymbol{\epsilon}\, \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]) &= \mathrm{tr}_{\mathcal{K}}(\boldsymbol{\epsilon}\, \boldsymbol{\epsilon}_B^2\, \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]) \\ = \mathrm{tr}_{\mathcal{K}}(\boldsymbol{\epsilon}_B\, \boldsymbol{\epsilon}\, \boldsymbol{\epsilon}_B\, \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]) &= -\, \mathrm{tr}_{\mathcal{K}}(\boldsymbol{\epsilon}\, \mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]). \end{aligned}$$

We have used that $\mathbf{c}_d(\frac{dx}{x})\mathsf{D}_Y[\exp(-T\mathsf{D}_Y^2)]$ is even w.r.t. $\boldsymbol{\epsilon}_B$ and that $\boldsymbol{\epsilon}\,\boldsymbol{\epsilon}_B = -\,\boldsymbol{\epsilon}_B\,\boldsymbol{\epsilon}$, when $h+1$ is odd.

The calculation of the other summand in (104) is straightforward, using Lemma 5.27. Recall that $Q_{\mathrm{tf}}(0) = 0$ as well as $Q_{\mathrm{tff}}(0) = 0$, so the corresponding terms in Theorem 5.24 do not contribute in the supertrace:

Lemma 5.28 *The contributions at* tf *and* tff *are*

(a) $\mathrm{str}(\mathsf{K})/\,dt|_{\mathrm{tf}\cap\triangle_H} = \sqrt{2\pi i}^{\,-n} \mathrm{top}_{\Lambda^d T^*X}[\widehat{A}(R^d)\ \mathrm{Ch}(F^{E/S,d})]$

(b) $\mathrm{str}(\mathsf{K})/\,dt|_{\mathrm{tff}\cap\triangle_H} = \sqrt{2\pi i}^{\,-h-1} \mathrm{top}_{\Lambda^b T^*B}[\widehat{A}(R^Y)\tau^{-\mathsf{N}_B/2}\ \mathrm{str}_{E/B}(\boldsymbol{\epsilon}_{V^*\partial X}\, \mathsf{K}_{\mathbf{B}^2}(\tau))]$

Proof. First, at $\overset{\circ}{\mathrm{tf}}\cong {}^{\phi}TX$ the density factor looks like $\tau^{n/2}t\mathrm{KD}_H|_{\mathrm{tf}\cap\triangle_H} = {}^{\phi}\Omega(X)\,dt$. Therefore

$$\begin{aligned}
\mathrm{str}(\mathsf{K})/\,dt|_{\mathrm{tf}\cap\triangle_H} &= \sqrt{2/i}^{\,n}\,\frac{\mathrm{top}_{\Lambda^d T^*X}}{dvol_{{}^dT^*X}}(\tau^{n/2}N_{G,\mathrm{tf},2}(\mathsf{K})/\,dt) \\
&= \sqrt{2/i}^{\,n}\,\mathrm{top}_{\Lambda^d T^*X}[p(R^d,0,1)\ \mathrm{str}_{E/S}(\exp(F^{E/S,d}))] \\
&= \sqrt{2\pi i}^{\,-n}\,\mathrm{top}_{\Lambda^d T^*X}[\widehat{A}(R^d)\ \mathrm{Ch}(F^{E/S,d})].
\end{aligned}$$

In (b), using the counting function N_B on $\Lambda^b T^*B$ and Lemma A.1 we get:

$$\begin{aligned}
\mathrm{str}(\mathsf{K})/\,dt|_{\mathrm{tff}\cap\triangle_H} &= \sqrt{2/i}^{\,h+1}\,\frac{\mathrm{top}_{\Lambda^b T^*B}}{dvol_{{}^bT^*B}}(\frac{x^{h+1}}{d\tau}\ \mathrm{str}_{E/B}(N_{G,\mathrm{tff},2}(\mathsf{K}))|_{\triangle_H}) \\
&= \sqrt{2/i}^{\,h+1}\,\mathrm{top}_{\Lambda^b T^*B}\left[p(R^Y,\mathbf{y}=0,\tau)\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau))\right]|_{\triangle_H} \\
&= \sqrt{2\pi i\tau}^{\,-h-1}\,\mathrm{top}_{\Lambda^b T^*B}[\widehat{A}(\tau R^Y)\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau))] \\
&= \sqrt{2\pi i}^{\,-h-1}\,\mathrm{top}_{\Lambda^b T^*B}[\widehat{A}(R^Y)\tau^{-\mathsf{N}_B/2}\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau))].
\end{aligned}$$

■

Using the definition of the fibred eta invariant in (117) and putting everything together we get the following index theorem:

Theorem 5.29 (Index Theorem) *For the Dirac operator* D *fulfilling (39) the index formula*

$$\mathrm{ind}_-(\mathsf{D}) = \frac{1}{(2\pi i)^{n/2}}\int_X \widehat{A}(R^d)\ \mathrm{Ch}(F^{E/S,d}) + \frac{1}{(2\pi i)^{(h+1)/2}}\int_Y \widehat{A}(R^Y)\widehat{\eta}(\mathsf{D}^{\phi,V},E) + \frac{1}{2}\eta(\mathsf{D}_Y)$$

holds. ■

Note that since $R^d \in \Gamma(\Lambda^2T^*X \otimes \mathrm{End}({}^dT^*X))$, the first integrand on the RHS is a *true* form on the compact manifold X and integrability is automatic. The second and third term on the RHS should be compared with the adiabatic limit formula (see [BiCh1], [Da1]) for the eta invariant of a geometric Dirac operator on $\partial X \to Y$ under collapsing of the metric in the fibres. We describe this in the case of the signature operator in the next Section.

Theorem 5.29 contains the usual APS (or b-) index theorem (see [APS] or [Me3]) for Dirac operators as the special case $Z = \{\mathrm{pt}\}$. Also, note that D has pure point spectrum when $\mathsf{D}^{\phi,V}$ is invertible. Then on the LHS we have the true index of D and the eta-term on the RHS vanishes. An extreme instance of this is the case $B = \{\mathrm{pt}\}$, when only the local term appears on the RHS.

5.4 The Signature Operator S

The signature operator on X for the metric g_d is the operator

$$\mathsf{S}^d = \mathsf{D}^d = d_d + d_d^* \quad \text{on} \quad E = \Lambda\,{}^dT^*X.$$

It has the property that its square, $(\mathsf{S}^d)^2$, is equal to the Laplacian on forms $\triangle_{\Lambda\, {}^dT^*X}$. Even if X is not spin, there is a *local* isomorphism $E \cong \mathrm{End}(S({}^dT^*X))$. The grading operator $\boldsymbol{\epsilon}$ on E is given by the Clifford action of the volume element on the left spinor factor. A short calculation shows that

$$\boldsymbol{\epsilon} = i^{n/2}\mathbf{c}_d(dvol_d) = i^{n/2} \star_d (-1)^{\mathsf{N}(\mathsf{N}-1)/2}.$$

For $n = \dim(X) = 4k$ we have $\star_d^2 = 1$ and the L^2_d-signature $\mathrm{sgn}_{L^2}(X, g_d)$ of X is defined as the signature of the quadratic form

$$\alpha \longmapsto (\alpha, \star_d \alpha) \quad \text{on} \quad H^{2k}_{\mathrm{Ho}}(X, g_d) := \mathrm{null}_{L^2_d}(\triangle_{\Lambda^{2k}\, {}^dT^*X}).$$

As before, the operator given by

$$\mathsf{S} = x^{v/2}\mathsf{S}^d x^{-v/2} = x^{v/2}\, d_d x^{-v/2} + x^{v/2}\, d_d^* x^{-v/2} =: \widehat{d} + \widehat{d}^*$$

is selfadjoint on $L^2_b(X, \Lambda\, {}^dT^*X)$ and its L^2_b-index equals $\mathrm{sgn}_{L^2}(X, g_d)$. The ϕ-differential operators

$$x\,\widehat{d}, x\,\widehat{d}^* : C^\infty(X, \Lambda\, {}^dT^*X) \to C^\infty(X, \Lambda\, {}^dT^*X),$$

have well defined restrictions to vertical differential operators over the boundary, which define the vertical cohomology of the boundary fibration:

Lemma 5.30 (a) $d_Z = x^{-\mathsf{N}_Z} x\,\widehat{d} x^{\mathsf{N}_Z} : C^\infty(\partial X, \Lambda V^*\partial X) \to C^\infty(\partial X, \Lambda V^*\partial X)$

(b) $d_Z^* = x^{-\mathsf{N}_Z} x\,\widehat{d}^* x^{\mathsf{N}_Z} : C^\infty(\partial X, \Lambda V^*\partial X) \to C^\infty(\partial X, \Lambda V^*\partial X)$

(c) $H(Z) = H(d_Z) \cong H_{\mathrm{Ho}}(d_Z)$ *is a vector bundle over* Y.

(d) $H(x\,\widehat{d}) \cong H_{\mathrm{Ho}}(x\,\widehat{d}) = \mathrm{null}(x\mathsf{S}|_{\partial X}) \cong C^\infty(Y, x^{\mathsf{N}_Z} H_{\mathrm{Ho}}(d_Z) \otimes \Lambda\, {}^bT^*B)$ *etc.*

Proof. Let $\alpha \in [{}^dN^*\partial X] = {}^bT^*B$. Then $d\alpha \in \Lambda^2\, {}^bT^*X$ and

$$[x\,\widehat{d}, \boldsymbol{\varepsilon}(\alpha)]|_{\partial X} = [x\,\widehat{d}, \boldsymbol{\iota}(\alpha)]|_{\partial X} = 0 \quad \text{in} \quad C^\infty(\partial X, \Lambda\, {}^dT^*X),$$

from which (a) and (b) follow. Parts (c) and (d) follow from the usual Hodge theoretic arguments. ∎

This Lemma shows that our main assumption (39) automatically holds for the signature operator. Recall the notation

$$C^\infty(X, \Lambda\, {}^dT^*X)^\circ = \{\xi \in C^\infty(X, \Lambda\, {}^dT^*X) \mid \quad \xi|_{\partial X} \in H_{\mathrm{Ho}}(x\,\widehat{d})\}.$$

The indicial operators $I_b(\widehat{d})$, $I_b(\widehat{d}^*)$, as well as the corresponding indicial families can be defined as in Section 3.5. For instance

$$\widehat{d}_Y(\xi|_{\partial X}) \equiv I_b(\widehat{d})(0)(\xi|_{\partial X}) := (\Pi_\circ \widehat{d}\xi)|_{\partial X},$$

with $(\xi|_{\partial X}) \in H_{\mathrm{Ho}}(x\,\widehat{d})$ extended to $\xi \in C^\infty(X, \Lambda\, {}^dT^*X)^\circ$ is well defined on $H_{\mathrm{Ho}}(x\,\widehat{d})$. The operators $\widehat{d}_Y$, $(\widehat{d}^*)_Y$ have the following properties

Lemma 5.31 (a) $(\widehat{d}_Y)^* = (\widehat{d}^{\,*})_Y$ *and we just write this as* $\widehat{d}^{\,*}_Y$

(b) $\widehat{d}_Y$ *is of total degree 1 on* $H_{\mathrm{Ho}}(x\,\widehat{d}) = C^\infty(Y, x^{\mathsf{N}_Z} H_{\mathrm{Ho}}(\,d_Z) \otimes \Lambda\,^bT^*B)$

(c) $(H_{\mathrm{Ho}}(x\,\widehat{d}),\ \widehat{d}_Y)$ *is an elliptical complex and* $H(\widehat{d}_Y) \cong H_{\mathrm{Ho}}(\widehat{d}_Y)$ *is finite dimensional.*

(d) $C^\infty(\partial X, \Lambda\,^dT^*X) \supset H_{\mathrm{Ho}}(x\,\widehat{d}) = C^\infty(Y, x^{\mathsf{N}_Z} H_{\mathrm{Ho}}(\,d_Z) \otimes \Lambda\,^bT^*B) \supset H_{\mathrm{Ho}}(\widehat{d}_Y)$ ∎

We have the following version of Proposition 3.15, describing the operators $\widehat{d}_Y$, $\widehat{d}^{\,*}_Y$:

Lemma 5.32 *The operators* $\widehat{d}_Y$, $\widehat{d}^{\,*}_Y$ *act on* $H_{\mathrm{Ho}}(x\,\widehat{d}) = C^\infty(Y, x^{\mathsf{N}_Z} H_{\mathrm{Ho}}(\,d_Z) \otimes \Lambda\,^bT^*B)$ *as operators of total order* 1, *resp.* -1. *We have*

$$\begin{aligned}
\widehat{d}_Y \quad = \quad \Pi_\circ \Big[& (\mathsf{h}\,\boldsymbol{\varepsilon}_d)\nabla^{\Lambda\,^dT^*X}_{\cdot} - \mathsf{v}\,\boldsymbol{\varepsilon}_d\,\lambda(x\mathsf{n}S^*_\phi(\cdot)\mathsf{v}) + \boldsymbol{\varepsilon}_d(\frac{dx}{x})(\mathsf{N}_Z - \frac{v}{2}) && (1,0)\\
& -\mathsf{v}\,\boldsymbol{\varepsilon}_d\,\nabla^{\Lambda\,^dT^*X}_{x\nabla^\phi_\cdot \nu} - (\mathsf{v}\,\boldsymbol{\varepsilon}_d, \mathsf{v}\,\boldsymbol{\varepsilon}_d, \mathsf{v}\,\boldsymbol{\iota}_d)\langle R^d(\mathsf{X}_d, \cdot)\cdot, \cdot\rangle_d && (0,1)\\
& - \frac{1}{2}(\mathsf{v}\,\boldsymbol{\varepsilon}_d, \mathsf{v}\,\boldsymbol{\varepsilon}_d, \mathsf{n}\,\boldsymbol{\iota}_d)\langle xB_\phi(\cdot,\cdot),\cdot\rangle_{\,d} \Big] && (-1,2)
\end{aligned}$$

$$\begin{aligned}
\widehat{d}^{\,*}_Y \quad = \quad \Pi_\circ \Big[& -(\mathsf{h}\,\boldsymbol{\iota}_d)\nabla^{\Lambda\,^dT^*X}_{\cdot} - \mathsf{v}\,\boldsymbol{\iota}_d\,\lambda(x\mathsf{v}S_\phi(\cdot)\mathsf{n}) + \boldsymbol{\iota}_d(\frac{dx}{x})(\mathsf{N}_Z - \frac{v}{2}) && (-1,0)\\
& +\mathsf{v}\,\boldsymbol{\iota}_d\,\nabla^{\Lambda\,^dT^*X}_{x\nabla^\phi_\cdot \nu} + (\mathsf{v}\,\boldsymbol{\iota}_d, \mathsf{v}\,\boldsymbol{\varepsilon}_d, \mathsf{v}\,\boldsymbol{\iota}_d)\langle R^d(\mathsf{X}_d, \cdot)\cdot, \cdot\rangle_d && (0,-1)\\
& + \frac{1}{2}(\mathsf{v}\,\boldsymbol{\iota}_d, \mathsf{v}\,\boldsymbol{\iota}_d, \mathsf{n}\,\boldsymbol{\varepsilon}_d)\langle xB_\phi(\cdot,\cdot),\cdot\rangle_{\,d} \Big] && (1,-2)
\end{aligned}$$

where the column on the right indicates the (n,v)*- degrees of the different parts. Note once again that the contributions of type* $(-1,2)$, $(0,1)$, $(1,-2)$ *and* $(0,-1)$ *simply vanish when* g_ϕ *is a product* ϕ*-metric.*

Proof. Recall that $F^E = R^{\Lambda\,^dT^*X} = -\lambda(R^{^dTX})$. We leave it to the more enthusiastic readers to use this with Proposition 1.14 and the method of proof of Proposition 3.15 to prove the claim. The appearance of the last summands in the $(1,0)$ and $(-1,0)$-terms above is due to the fact that

$$[\widehat{d}, \boldsymbol{\iota}_d(\mathsf{X}_d)]|_{\partial X} = [\widehat{d}^{\,*}, \boldsymbol{\varepsilon}_d(\frac{dx}{x})]|_{\partial X} = \mathsf{N}_Z - \frac{v}{2} \quad \text{on} \quad \Lambda\,^dT^*X.$$

∎

As in [BiLo], the fact that $\widehat{d}^{\,2}_Y = 0$ can be used to show that

$$\nabla^{\mathcal{H}(Z)}_T := \nabla^{\Lambda\,^dT^*X}_T + \lambda(\cdot\mathsf{v}S_\phi(\mathsf{v}\cdot)xT)$$

is a flat connection on $\mathcal{H}(Z) = x^{\mathsf{N}_Z} H_{\mathrm{Ho}}(\,d_Z) \otimes \langle \frac{dx}{x} \rangle \to Y$. In the case of a ϕ-product metric the model of the signature operator then is

$$\mathsf{S}_Y = \overline{d}_Y + \overline{d}^{\,*}_Y = \boldsymbol{\varepsilon}_Y\,\nabla^{\mathcal{H}(Z)}_{\cdot} - \boldsymbol{\iota}_Y\,\nabla^{\mathcal{H}(Z)^*}_{\cdot} + (\boldsymbol{\varepsilon}_d(\frac{dx}{x}) + \boldsymbol{\iota}_d(\frac{dx}{x}))(\mathsf{N}_Z - \frac{v}{2})$$

on $\mathcal{H}(Z)$. The operator S_Y thus differs by an endomorphism from the twisted signature operator $\mathsf{S}(Y, H(d_Z))$. However, we will only be interested in calculating the associated eta invariants. We know that the eta invariant vanishes in the case of odd dimensional fibres. In the case of even dimensional fibres, an easy symmetry argument shows that only the restriction of the operator S_Y to $\mathcal{H}^{v/2}(Z)$ contributes to its eta invariant. Therefore

$$\eta(\mathsf{S}_Y) = \eta(\mathsf{S}(Y, H(d_Z))),$$

Also, the signature operator S has the usual property that its extended L^2_b-index equals its true L^2_b-index:

Proposition 5.33 $\operatorname{ind}_-(\mathsf{S}) = \operatorname{ind}_{L^2_b}(\mathsf{S}) = \operatorname{sgn}_{L^2}(X, g_d)$.

Proof. This is shown just as in the b-case. Recall that we know from Proposition 3.28 that any element $u \in \operatorname{null}_-(\mathsf{S})$ is of the form $u = u_0 + O(x^\varepsilon)$ with $u_0 \in C^\infty(X, \Lambda^d T^* X)^\circ$. Defining J to be the corresponding restriction map $J(u) := u_0|_{\partial X}$ we get the sequence

$$0 \longrightarrow \operatorname{null}_{L^2_b}(\mathsf{S}) \longrightarrow \operatorname{null}_-(\mathsf{S}) \xrightarrow{J} H_{\mathrm{Ho}}(\widehat{d}_Y) \subset H(x\widehat{d}),$$

which is exact in the left and in the middle and compatible with the grading ϵ. Thus

$$\operatorname{ind}_-(\mathsf{S}) - \operatorname{ind}_{L^2_b}(\mathsf{S}) = \operatorname{sdim}(\operatorname{null}_-(\mathsf{S})) - \operatorname{sdim}(\operatorname{null}_{L^2_b}(\mathsf{S})) = \operatorname{sdim}(\operatorname{im}(J)),$$

and we proceed to show that the RHS vanishes. The idea is to use the decomposition

$$H(x\widehat{d}) = C^\infty(Y, x^{\mathsf{N}_Z} H(d_Z) \otimes \Lambda T^* Y) \oplus C^\infty(Y, x^{\mathsf{N}_Z} H(d_Z) \otimes \frac{dx}{x} \wedge \Lambda T^* Y), \tag{105}$$

and to show that $\operatorname{im}(J)$ decomposes accordingly, i.e. the projections on the left and right summand are maps in $\operatorname{im}(J)$. Once this is shown, the fact that the grading $\boldsymbol{\epsilon}$ switches the two summands in (105) implies that the ± 1 eigenspaces of $\boldsymbol{\epsilon}$ in $\operatorname{im}(J)$ have equal dimension.

Start with $\xi \in \operatorname{null}_-(\mathsf{S})$ such that $J(\xi) \neq 0$. We may then assume that ξ pairs to 0 with $\operatorname{null}_{L^2_b}(\mathsf{S})$ under the usual L^2_b-pairing (55). It then follows from Lemma 3.29 that $\xi = \mathsf{S}\eta$ for some $\eta \in \operatorname{null}_-(\mathsf{S}^2)$. According to Lemma 3.28 this has the form

$$\eta = x^0 \eta_0 + x^0 \log(x) \eta_1 + O(x^\varepsilon), \quad \text{with} \quad \eta_0, \eta_1 \in C^\infty(X, \Lambda^d T^* X)^\circ.$$

Then $J(\xi) = J(\widehat{d}\eta) + J(\widehat{d}^*\eta)$ is a decomposition of the desired type, since

$$\widehat{d}\eta = \widehat{d}\eta_0 + \boldsymbol{\varepsilon}(\frac{dx}{x})\eta_1 + \log(x)\widehat{d}\eta_1 + O(x^\varepsilon) = \boldsymbol{\varepsilon}(\frac{dx}{x})\eta_1 + O(x^\varepsilon)$$

and analogously $\widehat{d}^*\eta = -\boldsymbol{\iota}(\frac{dx}{x})\eta_1 + O(x^\varepsilon)$. ■

It then follows immediately that Theorem 5.29 has the following form

Theorem 5.34 (L^2-Signature Theorem)

$$\operatorname{sgn}_{L^2}(X, g_d) = \frac{1}{(\pi i)^{n/2}} \int_X L(R^d) + \frac{1}{(\pi i)^{(h+1)/2}} \int_Y L(R^Y) \widehat{\eta}(\mathsf{S}_Z, \Lambda V^* M) + \frac{1}{2} \eta(\mathsf{S}(Y, H(d_Z))).$$

Proof. The LHS in Theorem 5.34 equals $\operatorname{sgn}_{L^2}(X, g_d)$ as shown in Proposition 5.33. For the terms on the RHS use the fact that for $E = \Lambda^d T^*X$ the relative curvature $F^{E/S}$ is right Clifford multiplication $\mathbf{c}_{d,R} = \boldsymbol{\varepsilon}_d + \boldsymbol{\iota}_d$ with $\langle -R^d\cdot, \cdot\rangle_d/2$, i.e. (see [BGV], Chapter 4)

$$F^{E/S} = -\frac{1}{2}\mathbf{c}_{d,R}\langle R^d\cdot, \cdot\rangle_d = -\frac{1}{4}(\mathsf{n}\mathbf{c}_{d,R}, \mathsf{n}\mathbf{c}_{d,R})\langle R^{B,b}\cdot, \cdot\rangle_d - \frac{1}{4}(\mathsf{v}\mathbf{c}_{d,R}, \mathsf{v}\mathbf{c}_{d,R})\langle R^d\cdot, \cdot\rangle_d.$$

Recalling (see again [BGV]) that $\operatorname{Ch}(F^{E/S,d}) = 2^{-n/2}\operatorname{str}(\boldsymbol{\epsilon}\, F^{E/S})$ one shows that

$$\begin{aligned} L(R^d) &= 2^{-n/2}\widehat{A}(R^d)\operatorname{Ch}(F^{E/S,d}) \\ L(R^Y) &= L(R^{B,b}) = 2^{-(h+1)/2}\widehat{A}(R^{B,b})\operatorname{tr}_{S^*}(\exp(-\frac{1}{4}\mathsf{n}\mathbf{c}_{d,R}\langle R^{B,b}\cdot, \cdot\rangle_d)) \end{aligned}$$

Now rewrite (116) in Appendix A.5 as

$$\begin{aligned} \mathbf{B}^2 &= g^{-1}_{VM/M}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}, \boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}) + \frac{\kappa_{M/Y}}{4} - \frac{1}{8}(\mathsf{m}_d, \mathsf{m}_d, \mathbf{c}_{d,R}, \mathbf{c}_{d,R})\langle\delta_h(x^2 R^d(\cdot,\cdot))\cdot, \cdot\rangle_d \\ &=: \mathbf{B}^2_Z - \frac{1}{8}(\mathsf{h}\,\boldsymbol{\varepsilon}, \mathsf{h}\,\boldsymbol{\varepsilon}, \mathbf{c}_{d,R}, \mathbf{c}_{d,R})\langle\delta_h(x^2 R^{B,b}(\cdot,\cdot))\cdot, \cdot\rangle_d, \end{aligned}$$

and define $\widehat{\eta}(\mathsf{S}_Z, \Lambda V^*M)$ as in Appendix A.5, but using $\mathbf{B}^2_Z$ instead of $\mathbf{B}^2$. ■

Collapsing Cylinders

Fix a product structure,

$$C(M) = M \times [0, \varepsilon[_x \xrightarrow{\varphi} Y \times [0, \varepsilon[_x,$$

in a collar neighborhood of the boundary and consider the family of exact b-metrics given by

$$g_{b,\varepsilon} = \left(\frac{dx}{x}\right)^2 + \varphi^* g_Y + (x^2 + \varepsilon^2) g_Z,$$

on $C(M)$ and fixed on the interior of the manifold. Denote the associated curvature by $R^{b,\varepsilon} \in \Lambda^2\, {}^bT^*X \otimes \operatorname{End}({}^bTX)$ and the associated signature operator on $\Lambda\, {}^bT^*X$ by $\mathsf{S}_{b,\varepsilon}$. Then the b-signature theorem (or our Theorem 5.34 in the case with no fibre) tells us that

$$\operatorname{sgn}_{L^2}(X, g_{b,\varepsilon}) = \frac{1}{(\pi i)^{n/2}}\int_X L(R^{b,\varepsilon}) + \frac{1}{2}\eta(\mathsf{S}_{b,\varepsilon}|_{\partial X}). \tag{106}$$

It is known that

$$\operatorname{sgn}_{L^2}(X, g_{b,\varepsilon}) = \operatorname{sgn}(X, \partial X),$$

is independent of ε. As ε approaches 0, the metric $g_{b,\varepsilon}$ approaches the d-metric g_d and $L(R^{b,\varepsilon})$ approaches $L(R^d)$. Also, it follows from Dai's adiabatic limit formula for the signature operator in [Da1] that

$$\lim_{\varepsilon\searrow 0}\frac{1}{2}\eta(\mathsf{S}_{b,\varepsilon}|_{\partial X}) = \frac{1}{(\pi i)^{(h+1)/2}}\int_Y L(R^Y)\widehat{\eta}(\mathsf{S}_Z, \Lambda V^*M) + \frac{1}{2}\eta(\mathsf{S}(Y, H(\,d_Z))) + \boldsymbol{\tau} \tag{107}$$

The invariant $\boldsymbol{\tau}$ is defined in [Da1], section 4. It is a topological invariant associated to the Leray spectral sequence of the fibration $\phi : M \to Y$. Comparing the RHS in (107) with the RHS in our signature formula, we find the following relation for the L^2_d-signature, generalizing another formula in [Da1]:

$$\operatorname{sgn}_{L^2}(X, g_d) = \operatorname{sgn}(X, \partial X) - \boldsymbol{\tau}.$$

It is very likely that this result can be proved directly by analyzing the L^2_d-cohomology of X in more detail and establishing its relationship with the spectral sequence.

We leave considerations of this kind to the future.

Remark 5.35 The proof of the index formula, Theorem 5.29, presented in this text is far from "minimal". We therefore want finish this work with a brief discussion of possible shortcuts to the proof.

Recall that via the McKean-Singer argument, the LHS of the index formula corresponds to the large time limit of the regularized heat supertrace, whereas the RHS corresponds to the small time limit, plus the difference of the two in the form of the eta invariant. It is obvious from Section 5.3 that a full construction of the heat kernel is not necessary for the calculation of the RHS. All that is needed is a *parametrix*, solving the heat equation to first order at the faces tf, tff and hbf within the *rescaled* heat calculus. This means that we only need to solve the rescaled model problems at these faces. Especially, for the index result itself, no composition formulas for the heat calculus are needed.

On the LHS, it is easily seen that the parametrix for the resolvent constructed in Proposition 3.19 is compact for Λ close to 0, when D_Y is invertible. Thus D is Fredholm in that case, i.e. its spectral measure has a gap at 0 and D has a true index. The calculation of this index as the large time limit of the heat supertrace is then straightforward. Note that in the case of the signature operator S this assumption means that the flat cohomology bundle $H(Z) \to Y$ has to be *acyclic*, which seems somewhat too restrictive. However, given the index formula for Fredholm operators, we can easily deduce the general index formula by using a weight shifting argument, see [Me3] or [Va].

A Appendix

A.1 Clifford Algebra Conventions

Let (W, g_W) be a euclidean vectorspace of dimension n (usualy W will be the fibre of some tangent vector bundle) and $(W^*, g_{W^*} = g_W^{-1})$ its dual . We denote by $\mathrm{Cl}(W^*)$ the associated Clifford algebra over W^*. This is defined as the quotient of the tensor algebra over W^* by the relation

$$[\mathbf{c}(\alpha), \mathbf{c}(\beta)] = -2g_{W^*}(\alpha, \beta), \qquad \alpha, \beta \in W \subset \mathrm{Cl}(W^*).$$

Here, $\mathbf{c}$ is the inclusion of W^* into $\mathrm{Cl}(W^*)$ which can be regarded as an element $\mathbf{c} \in W \otimes \mathrm{Cl}(W^*)$. Elements $\alpha \in W^*$ act on the exterior algebra by

$$\lambda(\alpha)\omega := \boldsymbol{\varepsilon}(\alpha)\omega - \boldsymbol{\iota}(g_{W^*}(\alpha))\omega.$$

and it is straightforward to see that this induces an action λ of $\mathrm{Cl}(W^*)$ on ΛW^*. We will also use the action $\mathbf{c} \in W^* \otimes \mathrm{Cl}(W^*)$ by writing

$$\mathbf{c}(A) := \mathbf{c}(g_W(A, \cdot)), \qquad \text{for} \quad A \in W.$$

We will include the background metric into the notation, whenever this seems necessary. The *symbol map* is the the isomorphism of vector bundles given by

$$\sigma : \mathrm{Cl}(W^*) \longrightarrow \Lambda W^*, \qquad \sigma(\xi) = \lambda(\xi)1.$$

The inverse of this map (sometimes called “quantization” map) is a map

$$\mathbf{c} : \Lambda W^* \to \mathrm{Cl}(W^*),$$

which extends the above inclusion of W^* into $\mathrm{Cl}(W^*)$. On elements $\omega \in \Lambda^2 W^*$ this looks like

$$\mathbf{c}(\omega) = \sum_{i<j} \mathbf{c}(e_i^*)\mathbf{c}(e_j^*)\omega(e_i, e_j) = \frac{1}{2}\sum_{i,j} \mathbf{c}(e_i^*)\mathbf{c}(e_j^*)\omega(e_i, e_j) = \frac{1}{2}(\mathbf{c}, \mathbf{c})(\omega).$$

The notation on the RHS will also be used for general tensors $\eta \in W^* \otimes W^*$:

$$(\mathbf{c}, \mathbf{c})(\eta) := \sum_{i,j} \mathbf{c}(e_i^*)\mathbf{c}(e_j^*)\eta(e_i, e_j).$$

Beware of the following common source for sign errors: For an *endomorphism* $\Phi \in \mathrm{End}(W)$ we write

$$(\mathbf{c}, \mathbf{c})(\Phi) := (\mathbf{c}, \mathbf{c})(g_W(\Phi\cdot, \cdot)) \quad (= 2\mathbf{c}(\Phi) \quad \text{if } \Phi \text{ is skew}),$$

i.e. the “intuitive” order of the Clifford operations is reversed! Whenever the endomorphism Φ is skew, we have

$$[\frac{1}{2}\mathbf{c}(\Phi), \mathbf{c}(A)] = \mathbf{c}(\Phi(A)).$$

The Supertrace

Fix an orientation on W. Then the involution

$$\boldsymbol{\epsilon}_W := i^{[\frac{n+1}{2}]}\mathbf{c}(dvol_W), \qquad \boldsymbol{\epsilon}_W^2 = 1, \tag{108}$$

defines a (possibly trivial) grading on every representation space of $\mathrm{Cl}(W)$. In the following, let $n = \dim(W)$ be even and denote by $S(W)$ the spinor space for $\mathrm{Cl}(W)$ (a possible construction of $S(W)$ is described briefly in the next Appendix). This is the unique irreducible representation of $\mathrm{Cl}(W)$. The action of $\mathrm{Cl}(W)$ on $S(W)$ defines an isomorphism

$$\mathrm{Cl}(W) \cong \mathrm{End}(S(W)). \tag{109}$$

This representation is truly graded by the operator $\boldsymbol{\epsilon}_W$ and the identification (109) allows the corresponding supertrace of an element Q in $\mathrm{End}(S(W))$ to be calculated as follows

$$\mathrm{str}_{S(W)}(Q) = \mathrm{tr}(\boldsymbol{\epsilon}_W\, Q) = (\frac{2}{i})^{n/2}\mathrm{top}_{\Lambda W^*}(\sigma(Q))/\; dvol_W\,. \tag{110}$$

We will use this formula for the first factor in the decomposition

$$\mathrm{End}(E) = \phi^*\ \mathrm{Cl}({}^{\mathrm{b}}T^*B) \otimes \mathrm{END}_{\mathrm{Cl}({}^{\mathrm{b}}T^*B)}(E),$$

described in Lemma 5.1. Write $\boldsymbol{\epsilon} = \boldsymbol{\epsilon}_{{}^dT^*X}$, $\boldsymbol{\epsilon}_B = \boldsymbol{\epsilon}_{{}^bT^*B}$ and $\boldsymbol{\epsilon}_Z = \boldsymbol{\epsilon}_{{}^dV^*\partial X}$. Then, depending on the parity of the dimensions $n = \dim({}^dT^*X)$, $h+1 = \dim({}^bT^*B)$ and $v = \dim(V^*\partial X)$, with $n = h+1+v$ even, we have

$$\boldsymbol{\epsilon} = \boldsymbol{\epsilon}_B\, \boldsymbol{\epsilon}_Z \quad \text{if } h+1 \text{ is even,} \qquad \boldsymbol{\epsilon} = -i\, \boldsymbol{\epsilon}_B\, \boldsymbol{\epsilon}_Z \quad \text{if } h+1 \text{ is odd.}$$

For an element $Q_Z \in \mathrm{END}_{\mathrm{Cl}({}^{\mathrm{b}}T^*B)}(E)$ we define the *B-relative* supertrace as

$$\mathrm{str}_{E/B}(Q_Z) := \mathrm{C}(h)2^{-(h+1)/2}\ \mathrm{str}_E(\boldsymbol{\epsilon}_B\, Q_Z), \quad \text{with} \quad \mathrm{C}(h) = \begin{cases} \sqrt{i}^3 & h \quad \text{even} \\ 1 & h \quad \text{odd} \end{cases} \tag{111}$$

We can then show the following formula for the supertrace on E.

Lemma A.1 *For $Q_B \otimes Q_Z \in \mathrm{Cl}(\phi^{*b}T^*B) \otimes \mathrm{END}_{\mathrm{Cl}({}^{\mathrm{b}}T^*B)}(E)$ we have*

$$\mathrm{str}_E(Q_B \otimes Q_Z) = (\frac{2}{i})^{(h+1)/2}\mathrm{top}_{\Lambda^bT^*B}(\sigma(Q_B))/\; dvol_{{}^bT^*B} \cdot \mathrm{str}_{E/B}(Q_Z).$$

Proof. It suffices to prove this for $E = S({}^dT^*X)$ and $Q_Z \in \mathrm{Cl}({}^dV^*\partial X)$. If both v and $h+1$ are *even*, the Lemma is a direct consequence of (110):

$$\begin{aligned} \mathrm{str}_{S({}^dT^*X)}(Q_B \otimes Q_Z) &= \mathrm{str}_{S({}^bT^*B)}(Q_B)\ \mathrm{str}_{S({}^dV^*\partial X)}(Q_Z) \\ &= (\frac{2}{i})^{(h+1)/2}\mathrm{top}_{\Lambda^bT^*B}(\sigma(Q_B))/\; dvol_{{}^bT^*B} \cdot 2^{-(h+1)2)}\ \mathrm{tr}_{S({}^dT^*X)}(\boldsymbol{\epsilon}_{{}^dV^*\partial X}\, Q_Z). \end{aligned}$$

When v and $h+1$ are *odd*, we write $Q_B = \widehat{Q}_B \mathbf{c}_d(\frac{dx}{x})$ to see that

$$
\begin{aligned}
&\mathrm{str}_{S({}^dT^*X)}(Q_B \otimes Q_Z) \\
&\quad = (\frac{2}{i})^{h/2} \mathrm{top}_{\Lambda T^*Y}(\sigma(\widehat{Q}_B)) /\, dvol_{T^*Y} \cdot \mathrm{tr}_{S(\langle \frac{dx}{x} \rangle \oplus {}^dV^*\partial X)}(\boldsymbol{\epsilon}_{\langle \frac{dx}{x} \rangle \oplus {}^dV^*\partial X}\, \mathbf{c}_d(\frac{dx}{x}) Q_Z) \\
&\quad = (\frac{2}{i})^{h/2} \mathrm{top}_{\Lambda {}^bT^*B}(\sigma(Q_B)) /\, dvol_{{}^bT^*B}\, 2^{-h/2}\ \mathrm{tr}_{S({}^dT^*X)}(\boldsymbol{\epsilon}_Z\, Q_Z) \\
&\quad = (\frac{2}{i})^{(h+1)/2} \mathrm{top}_{\Lambda {}^bT^*B}(\sigma(Q_B)) /\, dvol_{{}^bT^*B}\, 2^{-(h+1)/2}\ \mathrm{tr}_{S({}^dT^*X)}(i^{3/2}\, \boldsymbol{\epsilon}\, \boldsymbol{\epsilon}_B\, Q_Z)
\end{aligned}
$$

where we have used $\boldsymbol{\epsilon}_{\langle \frac{dx}{x} \rangle \oplus {}^dV^*\partial X} = \mathbf{c}_d(\frac{dx}{x})\, \boldsymbol{\epsilon}_{{}^dV^*\partial X}$. ■

A.2 Dirac Operator and Conformal Transformations

In this Appendix we consider the behavior of Dirac operators under a conformal transformation of the metric.

Let M be an evendimensional manifold with metric g and $E \to M$ a Clifford bundle over M. Since our considerations will be purely local, we may assume right away that M is an open ball and E is trivial over M. Then E is of the form

$$E \cong S \otimes E', \quad \nabla^E \cong \nabla^S \otimes \nabla^{E'}$$

The spinor bundle S can be described as follows: There exists a polarization $T^*M_{\mathbf{C}} = W \oplus \overline{W}$ of the complexified cotangent bundle, i.e. $g(W, W) = 0$. Then $S = \Lambda W$ with metric g^S induced by g and Clifford action of an element $\alpha \in T^*M$ given by

$$\mathbf{c}(\alpha) = \mathbf{c}(\alpha_1 \oplus \overline{\alpha}_2) = \frac{1}{\sqrt{2}}(\boldsymbol{\varepsilon}(\alpha_1) - \boldsymbol{\iota}_g(\overline{\alpha}_2)).$$

Choosing a local orthonormal frame $e^1, \ldots, e^n$ for T^*M, the Levi Civita connection on T^*M is of the form $d + \omega$ with $\omega \in \Gamma(T^*M \otimes \mathbf{so}(T^*M))$. The spinor connection ∇^S is then given by $d + \frac{1}{2}\mathbf{c}(\omega)$.

Now, for a positive function $f \in C^\infty(M)$ consider the conformal metric $g_f = f^2 g$. The corresponding Levi Civita connection on T^*M is given for $T \in \Gamma(TM)$ and $\alpha \in \Gamma(T^*M)$ by

$$\nabla^f_T \alpha = \nabla_T \alpha - \frac{Tf}{f}\alpha + g^{-1}(\alpha, \frac{df}{f}) g(T) - \alpha(T)\frac{df}{f} =: \nabla_T \alpha + G_f(T)\alpha. \tag{112}$$

As before, the spinor bundle is $S_f = S = \Lambda W$ but the Clifford action is now given by

$$\mathbf{c}_f(\alpha) = \frac{1}{\sqrt{2}}(\boldsymbol{\varepsilon}(\alpha_1) - \boldsymbol{\iota}_f(\overline{\alpha}_2)) = \frac{1}{\sqrt{2}}(\boldsymbol{\varepsilon}(\alpha_1) - f^{-2}\, \boldsymbol{\iota}_g(\overline{\alpha}_2)).$$

The map

$$f : (T^*M, g) \ni \alpha \mapsto f \cdot \alpha \in (T^*M, g_f)$$

is an isometry and thus induces an isometry of the spinor bundles $f^{\mathsf{N}} : S \to S_f$, where N is the number operator in ΛW. We now have

$$f^{-\mathsf{N}}\mathbf{c}_f(\alpha)f^{\mathsf{N}} = f^{-\mathsf{N}}(\boldsymbol{\varepsilon}(\alpha_1) - f^{-2}\,\boldsymbol{\iota}_g(\overline{\alpha}_2))f^{\mathsf{N}} = f^{-1}\mathbf{c}(\alpha),$$

and from the description of ∇^S, ∇^{S_f} we deduce

$$f^{-\mathsf{N}}\nabla_T^{S_f} f^{\mathsf{N}} = \nabla_T^S + \frac{1}{2}\mathbf{c}(\boldsymbol{\varepsilon}(\frac{df}{f})\,\boldsymbol{\iota}(T) - \boldsymbol{\varepsilon}_g(T)\,\boldsymbol{\iota}_g(\frac{df}{f})) = \nabla_T^S + \widehat{G}(T),$$

where we have defined $G(T)$ to be $\frac{1}{2}(\mathbf{c}_g(T)\mathbf{c}(\frac{df}{f}) + \frac{df}{f}(T))$. This just means that the following diagrams are commutative

$$\begin{tikzcd} S \otimes T^*M \arrow[r, "f^{\mathsf{N}}\otimes\mathrm{id}"] \arrow[d, "f^{-1}\mathbf{c}"] & S_f \otimes T^*M \arrow[d, "\mathbf{c}_f"] \\ S \arrow[r, "f^{\mathsf{N}}"] & S_f \end{tikzcd} \qquad \begin{tikzcd} \Gamma(S) \arrow[r, "f^{\mathsf{N}}"] \arrow[d, "\nabla^S + \widehat{G}"] & \Gamma(S_f) \arrow[d, "\nabla^{S_f}"] \\ \Gamma(S \otimes T^*M) \arrow[r, "f^{\mathsf{N}}"] & \Gamma(S_f \otimes T^*M), \end{tikzcd}$$

and with these isomorphisms the corresponding Dirac operator is of the form

$$\mathsf{D}^f := \mathbf{c}_f \circ \nabla^{S_f} \overset{\wedge}{=} f^{-1}\mathsf{D} + \frac{n-1}{2}\mathbf{c}(\frac{df}{f^2}) = f^{-\frac{n-1}{2}} f^{-1}\mathsf{D} f^{\frac{n-1}{2}}.$$

Note that for some Clifford bundles like $E = \Lambda T^*M$ the bundle E' in the local decomposition $E = S \otimes E'$ might also contain copies of S or S^*. The above formulas remain true in that case, at the cost of a nonuniform treatment of the different copies of S in E.

Since it will be of some relevance in applications, see Section 5.4, we describe the case of the signature operator S on the bundle $E = \Lambda T^*M$ in detail. Clifford action is given $\mathbf{c}(\alpha) = (\boldsymbol{\varepsilon}(\alpha) - \boldsymbol{\iota}_g(\alpha))$ and our identifications above would give $\mathbf{c}_f(\alpha) = f^{-1}\mathbf{c}(\alpha)$. However, the natural isometry,

$$f^{\mathsf{N}} : (\Lambda T^*M, g) \longrightarrow (\Lambda T^*M, g_f)$$

yields an identification

$$f^{\mathsf{N}}\,\boldsymbol{\varepsilon}(\alpha)f^{-\mathsf{N}} = \boldsymbol{\varepsilon}(f\alpha) \quad \text{and} \quad f^{\mathsf{N}}\,\boldsymbol{\iota}(\alpha)f^{-\mathsf{N}} = \boldsymbol{\iota}(f^{-1}\alpha) = f\,\boldsymbol{\iota}_f(\alpha)).$$

Using the generalization to differential forms of (112) one can also show that

$$f^{\mathsf{N}}(\,d + \,d^*)f^{-\mathsf{N}} = f(\,d_f + \,d_f^*) - f\,\boldsymbol{\varepsilon}(\frac{df}{f})\mathsf{N} - f\,\boldsymbol{\iota}_f(\frac{df}{f})(\mathsf{N} - n). \tag{113}$$

A.3 Geometry of Fibre Bundles

In this Appendix we give a brief overview of the main notions in the geometry of riemannian fibre bundles. Our exposition follows [BGV] closely.

Let $\phi : M \to Y$ be a fibre bundle with fibre Z, with $m := \dim M, v := \dim Z$. The vertical bundle is denoted by $VM := \ker(\phi_*)$. By choosing a horizontal subbundle $HM \subset TM$ we get a decomposition

$$TM = VM \oplus HM,$$

and associated projections $\mathsf{v} : TM \to VM$, $\mathsf{h} : TM \to HM$. This decomposition is defined to be orthogonal with respect to a riemannian metric on M, of the form

$$\langle .,. \rangle_M = \phi^* \langle .,. \rangle_Y + \langle .,. \rangle_{M/Y}, \tag{114}$$

where $\langle .,. \rangle_Y$ and $\langle .,. \rangle_{M/Y}$ are given metrics on the base and on the fibres respectively.

Recall that the map $\phi_* : HM \to TY$ is a pointwise isomorphism. This allows us to freely identify HM with the lifted bundle ϕ^*TY. Denoting by ∇^M the Levi-Civita connection on M we then define the induced connections

$$\nabla^V \equiv \nabla^{M/Y} := \mathsf{v} \circ \nabla^M \circ \mathsf{v} : \Gamma(M, VM) \to \Gamma(M, T^*M \otimes VM)$$

$$\nabla^H := \phi^* \nabla^Y \circ \mathsf{h} : \Gamma(M, HM) \to \Gamma(M, T^*M \otimes HM).$$

These connections are invariant under multiplication of $\langle .,. \rangle_Y$ and $\langle .,. \rangle_{M/Y}$ with positive constants. The connection $\nabla^{M,\oplus} := \nabla^V \oplus \nabla^H$ is thus also invariant under this operation and compatible with the metric, but it can have torsion.

We now define the geometric tensors associated to the fibration:

Definition A.2 *Let X, Y be vector fields in TM*

(a) *The* difference tensor $\omega_M \in \Gamma(M, T^*M \otimes \mathbf{so}(TM))$ *is just*

$$\omega_M(X)Y := \nabla^M_X Y - \nabla^{M,\oplus}_X Y.$$

(b) *The* second fundamental form $S_M \in \Gamma(M, TM \otimes T^*M \otimes H^*M)$ *of the fibres Z is given by*

$$S_M(X)Y := \mathsf{v}\nabla^M_{\mathsf{v}X}\mathsf{h}Y.$$

(c) *The* curvature $\Omega_M \in \Gamma(\Lambda^2 T^*M, VM)$ *of the horizontal space is defined as*

$$\Omega_M(X,Y) := \mathsf{v}[\mathsf{h}X, \mathsf{h}Y].$$

The connection $\nabla^{M,\oplus}$ commutes with the projections h, v. Therefore

$$S_M(X)Y = \mathsf{v}\omega_M(\mathsf{v}X)\mathsf{h}Y, \qquad \Omega_M(X,Y) = \mathsf{v}(\omega_M(\mathsf{h}X)\mathsf{h}Y - \omega_M(\mathsf{h}Y)\mathsf{h}X).$$

By definition the term $\mathsf{v}\omega_M\mathsf{v}$ vanishes. Using the Koszul formula, we find furthermore that $\langle \omega_M(X)\mathsf{h}Y, \mathsf{h}Z \rangle = -\frac{1}{2}\langle \Omega(\mathsf{h}Y, \mathsf{h}Z), \mathsf{v}X \rangle$. With these formulas it is now easy to show:

Lemma A.3 *Let X, Y, Z be vector fields in TM. Then the following formula for ω_M holds:*

$$\begin{aligned}\langle \omega_M(X)Y, Z \rangle_M &= \langle S_M(\mathsf{v}X)\mathsf{h}Y, \mathsf{v}Z \rangle_M - \langle S_M(\mathsf{v}X)\mathsf{h}Z, \mathsf{v}Y \rangle_M \\ &-\frac{1}{2}\left(\langle \Omega_M(\mathsf{h}X, \mathsf{h}Z), \mathsf{v}Y \rangle_M - \langle \Omega_M(\mathsf{h}X, \mathsf{h}Y), \mathsf{v}Z \rangle_M + \langle \Omega_M(\mathsf{h}Y, \mathsf{h}Z), \mathsf{v}X \rangle_M \right). \qquad \blacksquare\end{aligned}$$

Let now $\mathcal{E} \to M$ be a Clifford bundle over M. The connection on $\mathcal{E}$ is denoted by $\nabla^{\mathcal{E}}$, the hermitian metric by $\langle .,. \rangle_{\mathcal{E}}$ and Clifford multiplication by $\mathbf{c}$. We assume that there is another connection $\nabla^{\mathcal{E},\oplus}$ on $\mathcal{E}$ which is a Clifford connection with respect to $\nabla^{M,\oplus}$, i.e.

$$[\nabla^{\mathcal{E},\oplus}_X, \mathbf{c}(\alpha)] = \mathbf{c}(\nabla^{M,\oplus}_X \alpha).$$

We then require that $\nabla^{\mathcal{E}}$ be of the form

$$\nabla^{\mathcal{E}}_X = \nabla^{\mathcal{E},\oplus}_X + \frac{1}{2}\mathbf{c}(\omega_M(X)).$$

This is clearly a Clifford connection with respect to ∇^M, as

$$[\nabla^{\mathcal{E}}_X, \mathbf{c}(Y)] = [\nabla^{\mathcal{E},\oplus}_X, \mathbf{c}(Y)] + [\frac{1}{2}\mathbf{c}(\omega_M(X)), \mathbf{c}(Y)] = \mathbf{c}(\nabla^{M,\oplus}_X Y + \omega_M(X)Y),$$

and the last term is equal to $\mathbf{c}(\nabla^M_X Y)$.

As an example, these requirements are satisfied for the bundle $\mathcal{E} = \phi^*\Lambda T^*Y \otimes \mathcal{E}'$, where $\mathcal{E}'$ is a vertical Clifford bundle, the Clifford action of $\alpha \in T^*Y$ is given by $\mathbf{c}(\alpha) = \boldsymbol{\varepsilon}(\alpha) - \boldsymbol{\iota}(\alpha)$, and $\nabla^{\mathcal{E},\oplus} = \phi^*\nabla^{\Lambda T^*Y} \otimes \nabla^{\mathcal{E}'}$. For these bundles we can also examine the degenerate Clifford action m given by

$$\mathsf{m}(\mathsf{h}\alpha) = \boldsymbol{\varepsilon}(\mathsf{h}\alpha), \qquad \mathsf{m}(\mathsf{v}\alpha) = \mathbf{c}(\mathsf{v}\alpha), \qquad [\mathsf{m}(\alpha), \mathsf{m}(\alpha)] = -2\langle \mathsf{v}\alpha, \mathsf{v}\alpha \rangle_M.$$

This degenerate Clifford action can be extended to more general tensors as in Appendix A.1. The associated 'degenerate' connection is then given by

$$\widehat{\nabla}^{\mathcal{E}}_X := \nabla^{\mathcal{E},\oplus}_X + \frac{1}{2}\mathsf{m}(\omega_M(X)).$$

The identity $[\frac{1}{2}\mathsf{m}(\omega(X)), \mathsf{m}(\alpha)] = \mathsf{m}(\alpha(\mathsf{v}\omega(X)))$ implies that this connection is a m-Clifford connection with respect to the connection

$$\widehat{\nabla}^{T^*M} := \nabla^{M,\oplus} + (\mathsf{v}\omega_M)^* \quad \text{or} \quad \widehat{\nabla}^{TM} := \nabla^{M,\oplus} + \mathsf{v}\omega_M.$$

This connection can be described more explicitly:

Lemma A.4 *Let $W \in VM$, $A \in HM$*

(a) $\widehat{\nabla}^{\mathcal{E}}_W = \nabla^{\mathcal{E},\oplus}_W + \frac{1}{2}(\boldsymbol{\varepsilon}, \mathsf{v}\mathbf{c})(\langle S_M(W)\mathsf{h}\cdot, \mathsf{v}\cdot \rangle_M) - \frac{1}{4}\boldsymbol{\varepsilon}(\langle \Omega_M(\mathsf{h}\cdot, \mathsf{h}\cdot), W \rangle_M)$

(b) $\widehat{\nabla}^{\mathcal{E}}_A = \phi^*\nabla^{\Lambda T^*Y}_A + \frac{1}{4}(\boldsymbol{\varepsilon}, \mathsf{v}\mathbf{c})(\langle \Omega_M(A,\cdot),\cdot \rangle_M)$.

Proof. For (a) note that

$$\begin{aligned} \mathsf{m}(\omega(W)) &= \boldsymbol{\varepsilon}(\mathsf{h}\omega(W)\mathsf{h}) + (\boldsymbol{\varepsilon}, \mathsf{v}\mathbf{c})(\mathsf{v}\omega(W)\mathsf{h}) \\ &= -\frac{1}{2}\boldsymbol{\varepsilon}(\langle \Omega(\mathsf{h}\cdot, \mathsf{h}\cdot), W \rangle_M) + (\boldsymbol{\varepsilon}, \mathsf{v}\mathbf{c})(\langle S_M(W)\mathsf{h}\cdot, \mathsf{v}\cdot \rangle_M) \end{aligned}$$

Analogously in (b), using

$$\langle \omega_M(A)Y, Z \rangle_M = -\frac{1}{2}(\langle \Omega_M(A, \mathsf{h}Z), \mathsf{v}Y \rangle_M - \langle \Omega_M(A, \mathsf{h}Y), \mathsf{v}Z \rangle_M),$$

we get

$$\mathsf{m}(\omega(A)) = \frac{1}{2}(\mathsf{m},\mathsf{m})(\langle\omega(A)\cdot,\cdot\rangle_M) = \frac{1}{2}(\boldsymbol{\varepsilon},\mathsf{v}\mathbf{c})(\langle\Omega_M(A,\cdot),\cdot\rangle_M),$$

and dividing everything by 2 gives the result. ■

Analogous to the definition of the Dirac operator $\mathsf{D} = \mathbf{c}\circ\nabla^{\mathcal{E}}$ on $\mathcal{E}$ we can define the superconnection corresponding to $\widehat{\nabla}^{\mathcal{E}}$ by $\mathbf{A}_M := \mathsf{m}\circ\widehat{\nabla}^{\mathcal{E}}$. In the case of a fibre bundle with base space Y a point this is again just the Dirac operator. Using Lemmas A.3 and A.4 we get

Lemma A.5 *Write* $\operatorname{tr}(S_M)(A) = \operatorname{tr}(\cdot S_M(\mathsf{v}\cdot)A) \in HM$. *Then*

$$\mathbf{A}_M := \mathsf{m}\circ\widehat{\nabla}^{\mathcal{E}} = d_H + \mathsf{D}^{\mathcal{E},V} + \frac{1}{2}\boldsymbol{\varepsilon}(\operatorname{tr}(S_M)) + \frac{1}{8}(\boldsymbol{\varepsilon},\boldsymbol{\varepsilon},\mathbf{c})\langle\Omega_M(\cdot,\cdot),\cdot\rangle_M.$$

Finally, using the fact that $\mathsf{h}\omega(A)B$ is symmetric in A and B it is easy to see that $\widehat{\nabla}^{TM}$ is torsionfree! Thus, the following generalization of the Lichnerowicz formula holds (cf. [BGV]):

Proposition A.6 *Set* $F^{\mathcal{E}/S}(X,Y) := F^{\mathcal{E}}(X,Y) - \frac{1}{2}\mathsf{m}(R^M(X,Y))$. *Then the following analogue of the Lichnerowicz formula holds:*

$$\mathbf{A}_M^2 = \widehat{\triangle}^{\mathcal{E},v} + \mathsf{m}(\widehat{F}^{\mathcal{E}}) = \widehat{\triangle}^{\mathcal{E},v} + \frac{\kappa_{M/Y}}{4} + \mathsf{m}(F^{\mathcal{E}/S}(\cdot,\cdot)).$$

Proof. The proof proceeds as usual.

$$2\mathbf{A}_M^2 = [\mathsf{m}\widehat{\nabla},\mathsf{m}\widehat{\nabla}] = \mathsf{m}\mathsf{m}[\widehat{\nabla},\widehat{\nabla}] + \mathsf{m}[\widehat{\nabla},\mathsf{m}]\widehat{\nabla} - \mathsf{m}[\mathsf{m},\widehat{\nabla}]\widehat{\nabla} + [\mathsf{m},\mathsf{m}]\widehat{\nabla}\widehat{\nabla}$$

The first summand on the RHS gives $2\mathsf{m}(\widehat{F}^{\mathcal{E}})$ since $\widehat{\nabla}^{TM}$ is torsionfree. The second and third summand vanish since $(\widehat{\nabla}^{\mathcal{E}},\, \widehat{\nabla}^{TM})$ is a Clifford connection. The last summand gives $-2g_M^{-1}(\mathsf{v}\widehat{\nabla},\mathsf{v}\widehat{\nabla})$, which is just two times the vertical Laplacian in the above formula. This proves the first equation in the Proposition, for the proof of the second equation we refer to [BGV] Chapter 10. ■

A.4 Mehler's Formula and the Simple Heat Space

In this Appendix we adapt the well known Mehler formula to our setting. For this, let (W,g_W) be a euclidean vector space and $\mathcal{A}$ a commutative algebra (usually this will be the space of even forms on some vector space). Let $R \in \operatorname{End}(W)\otimes\mathcal{A}$ such that $g_W(R) \in \Lambda^2W^*\otimes\mathcal{A}$. Also let $q \in W^*\otimes\mathcal{A}$. We consider $g_W(R\mathbf{y},\cdot)$ as a linear map $W \to W^*\otimes\mathcal{A}$ and q as a constant map of the same type. Then

Theorem A.7 *The operator*

$$H = g_W^{-1}\Big(\, d - \frac{1}{4}g_W(R) - \frac{1}{2}q,\; d - \frac{1}{4}g_W(R) - \frac{1}{2}q\Big) \tag{115}$$

on $C^\infty(W,\mathcal{A})$ *has the heat kernel* $[e^{-tH}](0,\mathbf{y}) = p(R,\mathbf{y},t)\exp(q(\mathbf{y})/2)\,dt\,dvol_W$ *given by the formula*

$$\begin{aligned} p(R,\mathbf{y},t) &= \sqrt{4\pi t}^{-n/2}\det\left(\frac{tR/2}{e^{tR/2}-e^{-tR/2}}\right)\exp\left(-\frac{1}{4t}\left\langle \mathbf{y}\left|\frac{tR}{2}\coth\left(\frac{tR}{2}\right)\right|\mathbf{y}\right\rangle_W\right), \\ p(R,0,t) &= \sqrt{4\pi t}^{-n/2}\det\left(\frac{tR/2}{e^{tR/2}-e^{-tR/2}}\right) =: \sqrt{4\pi t}^{-n/2}\widehat{A}(tR). \end{aligned}$$

Proof. The Mehler formula for the operator H_0 with $q = 0$ is well known and proved for example in [BGV] Chapter 4. For a general operator H just note that

$$H = e^{q(\mathbf{y})/2}H_0 e^{-q(\mathbf{y})/2},$$

from which the result follows immediately. ■

To be able to transfer this result to the model problems at a specific "temporal front face" we define the heat space, on which the above solution lives.



Figure 8: The Simple Heat Space $\overline{W}_H$

Denote by

$$W_H = [W \times [0,\infty[_t; \{0,0\},\ dt]$$

the "simple heat space" associated to W and by $\overline{W}_H$ its radial compactification at spatial and temporal infinity. As before, the blown up face at $t \to 0$ will be denoted by tf. The face at infinity will be called ti. Away from $t = 0$ the coordinate $t^{1/2}$ is a defining function for tf, and away from spatial infinity the coordinate $t^{-1/2}$ is a defining function for ti. We can identify the interior of *both* faces tf and ti with W such that for a constant vector field V in TW the restriction

$$t^{1/2}V|_{\mathrm{tf}} \quad \text{corresponds to} \quad V \in T\mathrm{tf} \quad \text{and} \quad t^{1/2}V|_{\mathrm{ti}} \quad \text{corresponds to} \quad V \in T\mathrm{ti}.$$

Equivalently the linear coordinate $\mathbf{Y}$ on tf or ti corresponds to $\mathbf{y}/t^{1/2}$ on the interior $W\times]0,\infty[$.

For $l, p > 0$ the associated "calculus" is defined by

$$\Psi^{l,p}_{H,\mathrm{s,cl}}(W) := \rho^l_{\mathrm{tf}}\rho^p_{\mathrm{ti}}\,\dot{C}^\infty_{\mathrm{tf,ti}}(\overline{W}_H, t^{-n/2}\frac{dt}{t}\Omega(W)).$$

This will be a recipient space for solutions of the heat equation. However these solutions cannot necessarily be composed with one another – at least we will not say how. Nevertheless we have normal maps

$$0 \longrightarrow \Psi^{l-1,p}_{H,\mathrm{s},\mathrm{cl}}(W) \longrightarrow \Psi^{l,p}_{H,\mathrm{s},\mathrm{cl}}(W) \overset{N'_{H,\mathrm{tf},l}}{\longrightarrow} \mathcal{S}(W, \Omega(W)\,dt) \to 0,$$

$$0 \longrightarrow \Psi^{l,p-1}_{H,\mathrm{s},\mathrm{cl}}(W) \longrightarrow \Psi^{l,p}_{H,\mathrm{s},\mathrm{cl}}(W) \overset{N'_{H,\mathrm{ti},p}}{\longrightarrow} \mathcal{S}(W, \Omega(W)\,dt) \to 0,$$

given by $N'_{H,\mathrm{tf},l}(A) := t^{1-l/2}A|_{\mathrm{tf}}$ and $N'_{H,\mathrm{tf},p}(A) := t^{p/2}A|_{\mathrm{ti}}$.

Also, a rescaled version of the simple heat "calculus" with coefficients in an algebra $\mathcal{A}$ can be defined. Assuming that the above algebra is of the form $\mathcal{A} = \Lambda^{ev}U^*$ for some vector space U we can introduce the heat calculus $\Psi_{G,\mathrm{s},\mathrm{cl}}(W)$ with coefficients in ΛU^* and rescaled w.r.t the corresponding filtration (which already is a grading) at tf. Using $dt^{1/2}$ to trivialize the conormal bundle $N^*\mathrm{tf}$ we get the rescaled normal operator

$$0 \longrightarrow \Psi^{l-1,p}_{G,\mathrm{s},\mathrm{cl}}(W) \longrightarrow \Psi^{l,p}_{G,\mathrm{s},\mathrm{cl}}(W) \overset{N'_{G,\mathrm{tf},l}}{\longrightarrow} \mathcal{S}(W, \Lambda^*U \otimes \Omega(W)\,dt) \longrightarrow 0.$$

We then have

Proposition A.8 (a) *The operator H maps*

$$\Psi^{l-2,p}_{G,\mathrm{s},\mathrm{cl}}(W) \xrightarrow{H} \Psi^{l,p}_{G,\mathrm{s},\mathrm{cl}}(W)$$

with rescaled normal operator $N_{G,\mathrm{tf},l-2}(H \circ A) = HN_{G,\mathrm{tf},l}(A)$.

(b) *The heat kernel for H is*

$$\mathsf{K}_H(\mathbf{y}, t) := p(R, \mathbf{y}, t)\exp(q(\mathbf{y})/2)\,dt\,dvol_W \in \Psi^{2,0}_{G,\mathrm{s},\mathrm{cl}}(W)$$

with normal operators at tf *and* ti *given by*

$$N_{G,\mathrm{tf},2}(\mathsf{K}_H)(\mathbf{Y}) = p(R, \mathbf{Y}, 1)\exp(q(\mathbf{Y})/2)\,dt\,dvol_W,$$

$$N_{G,\mathrm{ti},0}(\mathsf{K}_H)(\mathbf{Y}) = \sqrt{4\pi}^{-n/2}\exp(-\|\mathbf{Y}\|^2/4)\,dt\,dvol_W\,.$$

Proof. This follows from Mehler's formula once one has shown that $N_{G,\mathrm{tf},l-2}(H \circ A) = HN_{G,\mathrm{tf},l}(A)$. This is done as in Section 5.2 by calculating the normal action of the expressions appearing in (115), g_W^{-1}, d, $g_W(R)$ and q, using their mapping properties as described at the beginning of this Appendix. ■

We will also use the obvious generalization of these results to families, where W is really the fibre of some vector bundle, and apply it to the operators $H = \mathcal{H}_X, \mathcal{H}_B$ in Section 5.3.

A.5 Heat Kernels of Vertical Families

In this Appendix we introduce the heat calculus for vertical families of operators over the boundary fibration $M \to Y$. Define the heat space for vertical families as the parabolic blow up

$$M^2_{H,\mathrm{fib}} := [M \times_Y M \times [0,\infty[_\tau; \triangle_M \times \{0\},\ d\tau].$$

The interior of the blown up space $\mathrm{tf} \overset{\triangle}{=} \mathrm{tf}(M^2_{H,\mathrm{fib}})$ can be identified with the vector bundle $VM \to M$ in the usual manner. Also, we introduce the space $\overline{M}^2_{H,\mathrm{fib}}$, compactified at temporal infinity by introducing $1/\tau^{1/2}$ as a coordinate. The corresponding face will be denoted by ti. Define the density bundle

$$\mathrm{KD}_{M/Y,H} := \tau^{-v/2}\frac{d\tau}{\tau}\beta_L^*\, dvol_{M/Y}\,.$$

For $l > 0$ the associated family heat calculus is then defined by

$$\Psi^{l,p}_{H,\mathrm{fib,cl}}(M/Y) := \rho^l_{\mathrm{tf}}\rho^{p-v}_{\mathrm{ti}}\ \dot{C}^\infty_{\mathrm{tf,ti}}(\overline{M}^2_{H,\mathrm{fib}}, \mathrm{KD}_{M/Y,H}).$$

In the case $l = 0$ we have to add the mean value condition as in Section 4.1:

$$\Psi^{0,p}_{H,\mathrm{fib,cl}}(M/Y) := \dot{C}^\infty_{\mathrm{tf}}(\overline{M}^2_{H,\mathrm{fib}}, \mathrm{KD}_{M/Y,H})^{\mathrm{tf}} \otimes \mathbf{C}.$$

Leaving aside the behavior at temporal infinity for the moment, it can be shown as before that the composition formula holds

Theorem A.9 *Composition gives a map* $\Psi^l_{H,\mathrm{fib,cl}}(M) \times \Psi^r_{H,\mathrm{fib,cl}}(M) \xrightarrow{\circ} \Psi^{l+r}_{H,\mathrm{fib,cl}}(M)$. ■

It is also straightforward to describe the rescaled version of these constructions. The coefficient bundle from Section 5.1 is

$$\Lambda(\mathrm{tff}) = \Lambda\phi^{*b}T^*B \otimes \mathrm{END}_{\mathrm{Cl}(\phi^{*b}T^*B)}(E),$$

where we have used dx to trivialize $N^*\mathrm{tff}$. Writing ν_{tff} for the normal vector $P_{\mathrm{tff}}/\,dx$ the connection on $\Lambda(\mathrm{tff})$ for $T \in \beta_L^*TM/Y$ is then given by

$$\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}_T = \boldsymbol{\nabla}_T - \mathbb{K}(T,\nu_{\mathrm{tff}}) - \frac{1}{2}\nabla_{\nu_{\mathrm{tff}}}\mathbb{K}(T,\nu_{\mathrm{tff}}) : \Gamma(M^2_{H,\mathrm{fib}}, \Lambda(\mathrm{tff})) \to \Gamma(M^2_{H,\mathrm{fib}}, \Lambda(\mathrm{tff})).$$

It is calculated more explicitly in Corollary 5.17(b). As before, the filtration of $\Lambda(\mathrm{tff})$ over the front face tf is given by the total Clifford filtration. Using this data, we can define the rescaled bundle $\mathrm{Gr}_{\mathrm{tf}}(\Lambda(\mathrm{tff}))$ as in Section 5.1. Restriction to tf is defined using the connection $\nabla^{\Lambda(\mathrm{tff})}$:

$$|_{\mathrm{Gr,tf}} : \Gamma(\mathrm{Gr}_{\mathrm{tf}}(\Lambda(\mathrm{tff}))) \longrightarrow C^\infty(VM, \mathcal{N}\Lambda(\mathrm{tf},\mathrm{tff})).$$

For lack of (obvious) better alternatives we have used the notation on the RHS to denote the bundle $\mathcal{N}\mathcal{N}\Lambda(\mathrm{tf},\mathrm{tff})$ in (93) with factors $(N^*\mathrm{tff})^l$ trivialized by dx, i.e. we have written

$$\mathcal{N}\Lambda(\mathrm{tf},\mathrm{tff}) = \textstyle\bigoplus_k N^*\mathrm{tf}(M^2_{H,\mathrm{fib}})^k \otimes \Lambda^k({}^dT^*X) \otimes \mathrm{END}_{\mathrm{Cl}({}^dT^*X)}(E).$$

The normal operator

$$\begin{aligned} N'_{G,\mathrm{tf},l>0}(A) &:= \tau^{1-l/2} A|_{\mathrm{tf},\mathrm{Gr}} \in \mathcal{S}_{\mathrm{fib}}(VM, \Omega(VM/M)\, d\tau \otimes \mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff})) \\ N'_{G,\mathrm{tf},0}(A \oplus c) &:= \tau A|_{\mathrm{tf},\mathrm{Gr}} \oplus c \in \mathcal{S}_{\mathrm{fib}}(VM, \Omega(VM/M)\, d\tau \otimes \mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff}))^{\mathrm{tf}} \oplus \mathbf{C} \end{aligned}$$

fits into the short exact sequence for $l \geq 1$:

$$0 \to \Psi^{l-1}_{G,\mathrm{fib},\mathrm{cl}}(M/Y) \to \Psi^{l}_{G,\mathrm{fib},\mathrm{cl}}(M/Y) \xrightarrow{N_{H,\mathrm{tf},l}} \mathcal{S}_{\mathrm{fib}}(VM, \Omega(VM/M)\, d\tau \otimes \mathcal{N}\Lambda(\mathrm{tf}, \mathrm{tff})) \to 0,$$

and similarly for $l = 1$. The notation N'_G is used to remind the reader that these normal operators are defined using the temporal variable τ instead of t.

Our aim in the rest of this Appendix will be to construct the heat kernel of the operator

$$\mathbf{B}^2 := g^{-1}_{VM/M}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}, \boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}) + \frac{\kappa_{M/Y}}{4} + \mathsf{m}_d(\delta_h(x^2 F^{E/S,d}(\cdot,\cdot)) \tag{116}$$

as an element in $\Psi^{2,0}_{G,\mathrm{fib},\mathrm{cl}}(M/Y)$. The following Proposition calculates the normal operator of $\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}$ at tf.

Proposition A.10 *Let $T \in \beta_L^* T\, M/Y$ as before. Then the maps*

$$\Psi^l_{G,\mathrm{cl}}(M) \xrightarrow{\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}_{\tau^{1/2}T}} \Psi^l_{G,\mathrm{cl}}(X), \qquad \Psi^l_{G,\mathrm{cl}}(M) \xrightarrow{\frac{\partial}{\partial \tau}} \Psi^{l-2}_{G,\mathrm{cl}}(M)$$

have the following general form when restricted to the face tf

(a) $N'_{G,\mathrm{tf},l-1}(\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}_{\tau^{1/2}T} A) = (\boldsymbol{\nabla}^{\Lambda(\mathrm{tff})}_{\tau^{1/2}T} - \frac{1}{2} P_{\mathrm{tf}} \nabla^{\Lambda(\mathrm{tff})}_{\cdot} \mathbb{K}^{\Lambda(\mathrm{tff})}(\tau^{1/2}T, \cdot)) N'_{G,\mathrm{tf},l}(A)$

(b) $N'_{G,\mathrm{tf},l-2}(\frac{\partial}{\partial\tau} A) = \begin{cases} -\frac{1}{2}(L_{R^{VM}} + 2 - l - \mathsf{N}) N'_{G,\mathrm{tf},l}(A) & \textit{for} \quad l > 2 \\ -\frac{1}{2}(L_{R^{VM}} + 2 - l - \mathsf{N}) N'_{G,\mathrm{tf},2}(A) \oplus -\int_{VM/M} N'_{H,\mathrm{tf},2}(A) & \textit{for} \quad l = 2 \end{cases}$

Here N denotes the number operator in $\Lambda(\mathrm{tf})$. Note also that $\boldsymbol{\nabla}^{N^*\mathrm{tf}}_{R^{\phi N \partial X}}\, d\tau^{1/2} = 0$. ■

As in Section 5.2 the form of the normal operator of the connection in (a) can be calculated more explicitly:

Proposition A.11 *Write $\nu_{\mathrm{tf}} := P_{\mathrm{tf}} /\, d\tau^{1/2}$. Then*

$$P_{\mathrm{tf}} \nabla^{\Lambda(\mathrm{tff})}_{\cdot} \mathbb{K}^{\Lambda(\mathrm{tff})}(\tau^{1/2}T, \cdot)|_{U \in VM} = \frac{1}{2}(\, d\tau^{1/2})^2\, \boldsymbol{\varepsilon}(\delta_h \langle x^2 R^{d,V}(\cdot,\cdot)[\nu_{\mathrm{tf}} + \beta_L^* U], \beta_L^* T\rangle_\phi).$$

Proof. The proof proceeds just as in Proposition 5.16.

$$\begin{aligned} P_{\mathrm{tf}} \nabla^{\Lambda(\mathrm{tff})}_{\cdot} \mathbb{K}^{\Lambda(\mathrm{tff})}(\tau^{1/2}T, \cdot)|_{U \in VM} &= (\, d\tau^{1/2})^2 \mathbb{K}^{\Lambda(\mathrm{tff})}(\beta_L^* T, \beta_L^* U + \nu_{\mathrm{tf}})|_{\mathrm{tf}} \\ &= (\, d\tau^{1/2})^2 (\mathbb{K}(\beta_L^* T, \beta_L^* U + \nu_{\mathrm{tf}}))|_{\mathrm{tf},\mathrm{Gr}} = (\, d\tau^{1/2})^2 \delta_h(\mathbb{K}(\beta_L^* T, \beta_L^* U + \nu_{\mathrm{tf}}))|_{\mathrm{tf}} \\ &= \frac{1}{2}(\, d\tau^{1/2})^2\, \boldsymbol{\varepsilon}(\delta_h \langle x^2 R^{d,V}(\cdot,\cdot)[\nu_{\mathrm{tf}} + \beta_L^* U], \beta_L^* T\rangle_\phi)|_{\mathrm{tf}} \end{aligned}$$

Here we have used Corollary 5.20 and Proposition 5.6(c). ■

Multiplication with the function $\tau\kappa_{M/Y}$ does not contribute anything at tf, since the function vanishes there. Finally, the partial Clifford action

$$\mathsf{m}_d(\alpha) = \boldsymbol{\varepsilon}_d(\mathsf{n}\alpha) + \mathbf{c}_d(\mathsf{v}\alpha), \quad \alpha \in {}^dT^*X,$$

becomes exterior multiplication at tf:

Proposition A.12 *The map*

$$\Psi^{l}_{G,\mathrm{fib,cl}}(M/Y) \overset{\mathsf{m}_d(\alpha)}{\longrightarrow} \Psi^{l-1}_{G,\mathrm{fib,cl}}(M/Y)$$

has normal operator $N_{G,\mathrm{tf},l-1}(\mathsf{m}_d(\alpha)A) = \boldsymbol{\varepsilon}_d(\alpha)N_{G,\mathrm{tf},l}(A)$. ■

Using the methods described in the body of the text it is now easy to show the following

Theorem A.13 *The heat kernel for* $\mathbf{B}^2$ *is an element*

$$\mathsf{K}_{\mathbf{B}^2} \in \Psi^{2,0}_{G,\mathrm{fib,cl}}(M/Y).$$

It has normal operators at tf *and* ti *given by*

(a) $N'_{G,\mathrm{tf},2}(\mathsf{K}_{\mathbf{B}^2})(\mathbf{z}) =$

$$\delta_h \left[p(\mathbf{z}, x^2R^{d,V}, 1)\exp(\tfrac{1}{2}\langle x^2R^{d,V}\nu_{\mathrm{tf}}, \mathbf{z}\rangle_{VM/M})\exp(-x^2F^{E/S,d})\right] dvol_{VM/M}\ d\tau$$

(b) $N'_{G,\mathrm{ti},0}(\mathsf{K}_{\mathbf{B}^2}) = [\Pi_\circ]\ d\tau$*, the projection onto the null space of* $\mathsf{D}^{\phi,V}$

We emphasize once more that all these normal operators commute with $\boldsymbol{\varepsilon}(\frac{dx}{x})$.

Proof. The first part is obtained using the specialization of the methods in Chapters 4 and 5 based on the series of Propositions above. For part (b) we just mention that

$$\mathbf{B}^2 = (\mathsf{D}^{\phi,V})^2 + \quad \text{a vertical operator of order} \geq 1 \text{ in } \Lambda\,\phi^{*b}T^*B\ .$$

This is just of the type used in the Berline-Vergne theorem (Theorem 9.19 in [BGV]). ■

Denote by $\mathrm{str}_{E/B}$ the relative supertrace in $\mathrm{End}_{\mathrm{Cl}(^bT^*B)}(E)$ as defined in (111), and by N_B the number operator on $\Lambda\,\phi^{*b}T^*B$. Also, write T for the time axis $[0,\infty]_{\tau^{1/2}}$, compactified by $\tau^{-1/2}$ at infinity. Then

Corollary A.14 (a) $\tau^{-\mathsf{N}_B/2}\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau)|_{\triangle_M}) \in C^\infty(M \times \mathsf{T}, \Lambda\,\phi^{*b}T^*B \otimes \Omega(VM/M)\ d\tau)$.

(b) $\boldsymbol{\iota}(x\frac{\partial}{\partial x})\tau^{-\mathsf{N}_B/2}\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau)|_{\triangle_M})$ *vanishes in* $\tau^{1/2} = 0$ *and* $\tau^{-1/2} = 0$. ■

This Corollary allows us to define the ("twisted") family eta invariant as the form on the base Y given by

$$\widehat{\eta}(\mathsf{D}^{\phi,V}, E) := \int_0^\infty \int_{M/Y} \boldsymbol{\iota}(x\frac{\partial}{\partial x})\tau^{-\mathsf{N}_B/2}\ \mathrm{str}_{E/B}(\mathsf{K}_{\mathbf{B}^2}(\tau)|_{\triangle_M}). \tag{117}$$

This is an even (resp. odd) form on Y, whenever v, $h+1$ are odd (resp. even).

Remark A.15 By Remark 5.22 this is just the classical family eta invariant, whenever $B_\phi = 0$. Note however that in the literature ([BiCh1], [BiCh2], [Da1], (111)) authors hide of 2, π, and i in their definitions of the family eta invariant in many different ways.

A.6 Conormal Functions, Pullbacks and Pushforwards

In this Appendix we explain some of the essential notions introduced in [Me0], see also [Me3] or [Lo], and adapt them to our needs.

Let N be a manifold with corners. Let $\mathcal{M}(N) = \{H_1, \dots, H_f\}$ be the set of boundary faces of N and choose a boundary defining function $\rho_{H_j} \equiv \rho_j$ for each face H_j. As usual, denote by ${}^b\mathcal{V}(N)$ the space of vector fields on N tangent to the boundaries.

The space of functions conormal to the faces of N is defined as

$$\mathcal{A}(N) := L^\infty H_b^\infty(N) = \{u \in L^\infty(N) \quad | \quad {}^b\mathcal{V}(N) \cdot u \subset L^\infty(N)\}.$$

Elements in $\mathcal{A}(N)$ are C^∞ in the interior and bounded, but control at the boundary is only very moderate. One might think of such functions as being "C^∞ along the boundary". Choosing a f-tuple $\mathbf{a} := ((a_1, m_1), \dots, (a_f, m_f))$ with a pair $(a_j, m_j) \in \mathbf{C} \times \mathbf{N}$ for each face H_j of N, we can also define the space of conormal functions on N with *conormal bounds* $\mathbf{a}$ as

$$\mathcal{A}^{\mathbf{a}}(N) := \rho_1^{a_1} \log(\rho_1)^{m_1} \cdots \rho_f^{a_f} \log(\rho_f)^{m_f} \cdot \mathcal{A}(N). \tag{118}$$

A special class of conormal functions is given by functions which possess "generalized Taylor expansions" into powers of the type

$$\rho_j^z \log(\rho_j)^m, \qquad (z, m) \in \mathbf{C} \times \mathbf{N} \tag{119}$$

at the boundary faces H_j of N. To describe these expansions, we list the admissible powers (119) in an "index set". To explain this concept, introduce a partial ordering on $\mathbf{C} \times \mathbf{N}$, describing the comparative vanishing order at H_j of expressions of type (119), by declaring

$$(z_1, m_1) \leq (z_2, m_2) \quad \Longleftrightarrow \quad [\mathrm{Re}(z_1) < \mathrm{Re}(z_2) \quad \text{or} \quad \mathrm{Re}(z_1) = \mathrm{Re}(z_2) \quad \text{and} \quad m_1 \geq m_2].$$

Now, a (C^∞-)*index set* is a countable subset $I \subset \mathbf{C} \times \mathbf{N}$ such that

- I is bounded from below in the above ordering, i.e. there is a "worst" power in the expansion,
- $I \cap B_N(0) \times \{0, \dots, N\}$ is finite for any $N \in \mathbf{N}$, i.e. powers in the expansion "improve" steadily,
- $(z, m) \in I$ implies $(z + n, m - l) \in I$ for any $l, n \in \mathbf{N}$ and $l \leq m$. This is the C^∞-condition.

Often, a complex number z will be interpreted as the index element $(z, 0)$.

Now given a collection $\mathcal{I} = (I_{H_1}, \dots, I_{H_f})$ of (C^∞-) index sets associated to the faces of N, we can recursively define the space of (classical) conormal functions associated to $\mathcal{I}$ as follows: A section $f \in \mathcal{A}^{\mathcal{I}}(N)$ has an expansion at each face H of the form

$$f \sim \sum_{(z,m) \in I_H} \rho_H^z \log(\rho_H)^m f_{H,z,m}, \qquad f_{H,z,m} \in \mathcal{A}^{\mathcal{I}'}(N), \quad I'_F = \left\{ \begin{array}{ll} E_F & F \neq H \\ \mathbf{N} & F = H \end{array} \right. . \tag{120}$$

Obviously, for $\mathbf{a} = ((a_1, m_1), \ldots, (a_f, m_f))$ as above, we have

$$\mathcal{A}^{\mathcal{I}}(N) \subset \mathcal{A}^{\mathbf{a}}(N), \quad \text{if} \quad (a_j, m_j) \le I_{H_j}.$$

We leave it to the reader to define spaces of conormal functions when one or several index sets are finite. The largest or "best" index is then to be interpreted as a conormal bound in the sense of (118), and the C^∞-condition is assumed to hold up to that best index.

It will be important to keep track of the behavior of the index sets of conormal functions w.r.t. different operations.

Definition A.16 *Let I, J be two C^∞-index sets. Besides the simple set-theoretic union $I \cup J$ we will use the following other operations on I and J:*

(a) $I + J := (\,(z + z', m + m')\,|(z, m) \in I, (z', m') \in J\,)$

(b) $I \overline{\cup} J := (\,(z, j)\,|\,j \le m + m' + 1,\ (z, m) \in I, (z, m') \in J\,)$

(c) $\widehat{I} := \overline{\bigcup}_{l \in \mathbf{N}} (l + I)$

It should be clear that

$$f + g \in \mathcal{A}^{\mathcal{I} \cup \mathcal{J}}(N) \quad \text{and} \quad f \cdot g \in \mathcal{A}^{\mathcal{I} + \mathcal{J}}(N) \quad \text{for} \quad f \in \mathcal{A}^{\mathcal{I}}(N), \quad g \in \mathcal{A}^{\mathcal{J}}(N).$$

Especially, the space of conormal functions $\mathcal{A}^{\mathcal{I}}(N)$ is $C^\infty(N)$-module for any C^∞-index set $\mathcal{I}$.

The reader should verify that the coefficients $f_{H,z,m}$ in the expansion (120) of $f \in \mathcal{A}^{\mathcal{I}}(N)$ are not unique. However, if we have $(z, m) \in I_H$ but $(z - n, m) \notin I_H$ for any $n \in \mathbf{N}_+$, then the coefficient

$$f_{H_j,z,m}|_{H_j} \in \mathcal{A}^{(I_{H_1}, \ldots, I_{H_{j-1}}, I_{H_{j+1}}, \ldots)}(H_j)$$

is well defined. We therefore define the "leading part" of an index set I as

$$\operatorname{lead}(I) := \{(z, m) \in I \quad | \quad (z - n, m) \notin I \quad \text{for any} \quad n \in \mathbf{N}_+\}.$$

Note the rule

$$I - \operatorname{lead}(I) = 1 + I \quad !$$

Information associated to the leading coefficients of a conormal function is more persistent, in a sense that we describe now.

In the case of the Dirac operator D^ϕ over the "manifold with corners" $N = X$ an important piece of information is given by the decomposition of the space of sections over the boundary into zero modes and nonzero modes: By our main assumption (39) the null space of the vertical Dirac family $\mathsf{D}^{\phi,V}$ over the boundary ∂X is the space of sections of a vector bundle $\mathcal{K} \to Y$ and the projections $\Pi_\circ$, $\Pi_\perp$ onto this null space and its orthogonal complement give a decomposition

$$C^\infty(\partial X, E) = \Pi_\circ C^\infty(\partial X, E) \oplus \Pi_\perp C^\infty(\partial X, E) \triangleq \Gamma(Y, \mathcal{K}) \oplus \Gamma(Y, \mathcal{K})^\perp.$$

For a section $\xi \in \mathcal{A}^I(X,E)$ it now clearly makes sense to ask, whether the *leading* coefficients $\xi_{(z,n)}$ in its asymptotic expansion of type (120) restrict to the zero modes over the boundary or not. To keep track of this information, we agree to let an index $(z,m)^\circ$ denote a coefficient in the zero modes and $(z,n)^\perp$ denote a coefficient perpendicular to the zero modes:

$$\xi_{(z,n)^\circ}|_{\partial X} \in \mathcal{K}, \qquad \xi_{(z,n)^\perp}|_{\partial X} \perp \mathcal{K}.$$

More generally, given an index set I, we we define $[\circ]I$ to be the same index set, but carrying the additional information that the leading coefficients lie in the zero modes:

$$[\circ]I := \mathrm{lead}(I)^\circ \cup (I - \mathrm{lead}(I)),$$

and $[\perp]I$ is defined analogously.

Of course, these notations also make sense for the manifold $N = X^2_\phi$. Here the null space of the vertical family $\mathsf{D}^{\phi,V}$ defines vector bundles over the bases of lf, ϕbf, ff and we can use the above notation at each of these faces. We can now describe the mapping properties of the Dirac operator D^d (lifted from the left to X^2_ϕ):

Lemma A.17 (a) $\mathsf{D}^d : C^\infty(X,E)^\circ \to C^\infty(X,E)$

(b) $\mathsf{D}^d : \mathcal{A}^{[\circ]I}(X,E) \to \mathcal{A}^I(X,E)$.

(c) $\mathsf{D}^d : C^\infty(X^2_\phi, \mathrm{END})^\circ \to \rho_{\mathrm{ff}}^{-1} C^\infty(X^2_\phi, \mathrm{END})$

(d) $\mathsf{D}^d : \mathcal{A}^{(I_{\mathrm{rf}},[\circ]I_{\mathrm{lf}},[\circ]I_{\phi\mathrm{bf}},I_{\mathrm{ff}})}(X^2_\phi, \mathrm{END}) \to \mathcal{A}^{(I_{\mathrm{rf}},I_{\mathrm{lf}},I_{\phi\mathrm{bf}},I_{\mathrm{ff}}-1)}(X^2_\phi, \mathrm{END})$ ■

b-Fibrations, Pullback and Pushforward

Let N, N' be two manifolds with corners. The corresponding sets of boundary faces are $\mathcal{M}(N)$, $\mathcal{M}(N')$. A C^∞-map $F : N_1 \to N_2$ will be called a *b-map*, if it maps boundary faces to boundary faces in the sense that for every face $H \in \mathcal{M}(N')$ we require

$$F^*\rho'_H = a_H \prod_{G\in\mathcal{M}(N)} \rho_G^{e(G,H)}, \quad \text{or} \quad F^*\rho'_H \equiv 0,$$

with $e(G,H) \in \mathbf{N}$, $0 < a_H \in C^\infty(N)$. If no $F^*\rho'_H$ is identically 0, F will be called an *interior* b-map. The matrix $\mathbf{e} = (e(G,H))$ will be called the *coefficient matrix* of F. The readers can convince themselves, that composition of two b-maps is again a b-map, and that the coefficient matrix of the composition is the product of the coefficient matrices of the components.

The following result about the pullback of conormal functions under b-maps will be used several times in the text

Theorem A.18 (Pullback) *Let $F : N \to N'$ be an interior b-map, with coefficient matrix $\mathbf{e}$. Let $\mathcal{I}'$ a (possibly finite) index set for N'. Then*

$$F^* : \mathcal{A}(N')^{\mathcal{I}'} \longrightarrow \mathcal{A}(N)^{F^\sharp\mathcal{I}'},$$

where the index set $F^\sharp \mathcal{I}'$ *is defined as*

$$(F^\sharp \mathcal{I}')_G = (\,(\sum_{H \in \mathcal{M}(N')} e(G,H) z'_H, \sum_{H \in \mathcal{M}(N')} m'_H) \,|\, (z'_H, m'_H) \in I'_H\,),$$

which we read as $\mathbf{N}$*, if* G *is mapped to the interior of* N' *by* F*.* ■

The corresponding result for the pushforward of functions is not quite as easy to formulate. Also we will have to restrict the class of admissible maps F further. First, note that an interior b-map $F : N \to N'$ maps

$$F_* : {}^bT_pN \to {}^bT_{F(p)}N' \quad \text{and} \quad F_* : {}^bN_pN \to {}^bN_{F(p)}N'$$

for all $p \in N$. If the first of these (b-) differentials is always surjective, F will be called a *b-submersion.* It will be called b-normal, if the second (b-) differential is always surjective. An interior b-map F, which is both, a b-submersion and b-normal, will be called a b-fibration. A more intuitive description of b-fibrations is stated in [Me0]:

Proposition A.19 *Let* $F : N \to N'$ *be a b-submersion. Then* F *is a b- fibration* if and only if *for each face in the preimage* $G \in \mathcal{M}(N)$ *there is* at most one *face in the image* $H \in \mathcal{M}(N')$*, such that* $e(G,H) \neq 0$*.* ■

We can now formulate the pushforward theorem in the version given in [Lo]:

Theorem A.20 (Pushforward) *Let* $F : N \to N'$ *be a b-fibration, and let* $\mathcal{I}$ *be an index set for* N *such that* $I_G > 0$ *for all faces* G *of* N *which are mapped to the interior of* N'*. Then*

$$F_* : \mathcal{A}^{\mathcal{I}}(N, \Omega(N)) \longrightarrow \mathcal{A}^{F_\sharp \mathcal{I}}(N', \Omega(N')),$$

$$\textit{with} \quad (F_\sharp \mathcal{I})_H := \overline{\bigcup}_{G \in \mathcal{M}(N), F(G) \subset H} e(G,H)^{-1} I_G,$$

for all $H \in \mathcal{M}(N')$*.* ■

References

[APS] M.F. Atiyah, V.K. Patodi, I.M. Singer : “Spectral Asymmetry in Riemannian Geometry I”, Math. Proc. Camb. Phil. Soc. **77** (1975), 43-69

[BGV] N. Berline, E. Getzler, M. Vergne : “Heat Kernels and Dirac Operators”, Grundl. Math. Wiss. **297**, Springer (1992)

[BiCh1] J.M. Bismut, J. Cheeger :“η-Invariants and their Adiabatic Limits”, Jour. AMS **2** (1989), 33–70

[BiCh2] J.M. Bismut, J. Cheeger :“ Transgressed Euler Classes of SL$(2n, \mathbf{Z})$- Vector Bundles, Adiabatic Limits of Eta Invariants and Special Values of L-Functions”, Ann. Sc. ENS **25** (1992), 335–391

[BiLo] J.M. Bismut, J. Lott :“ Flat Vector Bundles, Direct Images and Higher Real Analytic Torsion”, Jour. AMS **8** (1995), 291–362

[Bu] U. Bunke : “Spektraltheorie von Diracoperatoren auf offenen Mannigfaltigkeiten”, Doktorarbeit, Uni Greifswald (1991)

[Ch] T. Christiansen : “Scattering Theory for Manifolds with Asymptotically Cylindrical Ends”, JFA **131** (1995), 499-530

[Da1] X. Dai : “Adiabatic Limits, Nonmultiplicativity of the Signature and the Leray Spectral Sequence”, Jour. AMS **4** (1991), 265–321

[Da2] X. Dai : “Geometric Invariants and their Adiabatic Limits”, Proc. of Symp. **54** (1993), 145–176

[DaMe] X. Dai, R.B. Melrose : “Adiabatic Limit of the Torsion” (manuscript, unpublished)

[LaMo1] R. Lauter, S. Moroianu : “Fredholm Theory for Degenerate Pseudodifferential Operators on Manifolds with Fibred Boundaries” (Preprint 23/99, University of Mainz, to appear in Comm. Part. Diff. Eq.)

[LaMo2] R. Lauter, S. Moroianu : “Homology of Pseudodifferential Operators on Manifolds with Fibred Boundaries”

[Lo] P. Loya : “On the b-Pseudodifferential Calculus on Manifolds with Corners”, Thesis, MIT (1998)

[Ma] R. Mazzeo : “Elliptic Theory of Differential Edge Operators I”, Comm. Part. Diff. Eq. **16** (1991), 1615–1664.

[MaMe1] R. Mazzeo, R.B. Melrose : “Meromorphic Extension of the Resolvent on Complete Spaces with Asymptotically Constant Negative Curvature”, J. Funct. Anal. **75** (1987), 260–310.

[MaMe2] R. Mazzeo, R.B. Melrose: "The Adiabatic Limit, Hodge Cohomology and Leray's Spectral Sequence for a Fibration", J. Diff. Geom. **31** (1990), 185–213.

[MaMe3] R. Mazzeo, R.B. Melrose: "Pseudodifferential Operators on Manifolds with Fibred Boundaries", Asian J. Math **2** (1998), 833–866

[Me0] R.B. Melrose : "Differential Analysis on Manifolds with Corners", Preprint, MIT, http://www-math.mit.edu/ rbm/

[Me1] R.B. Melrose : "Calculus of Conormal Distributions on Manifolds with Corners", Int. Math. Res. Notices (1992), 51–61

[Me2] R.B. Melrose : "Spectral and Scattering Theory for the Laplacian on Asymptotically Euclidean Spaces", Lecture Notes in Pure and Appl. Mat. **161**, (1992), 85–130

[Me3] R.B. Melrose : "The Atiyah-Patodi-Singer Index Theorem", Research Notes in Mathematics vol.**4** (1993), A.K. Peters

[Me4] R.B. Melrose : "Geometric Scattering Theory", Stanford Lectures, Cambridge University Press (1995)

[Mü1] W. Müller : "Signature Defects of Cusps of Hilbert Modular Varieties and Values of L-Series at $s = 1$", Jour. Diff. Geom. **20** (1984), 55-119

[Mü2] W. Müller : "Manifolds with Cusps of Rank one. Spectral Theory and L^2-Index Theorem", LNM **1244** (1987), Springer

[Mü3] W. Müller : "Eta Invariants and Manifolds with Boundary", Jour. Diff. Geom. **40** (1994), 311-377

[Mü4] W. Müller : "On the Index of Dirac Operators on Manifolds with Corners of Codimension Two I.", MPI-Preprint **95-23** (1995)

[Ro] J. Roe : "Elliptic operators, topology and asymptotic methods", Pitman Research Notes in Mathematics **179** (1988)

[Sh] M.A. Shubin : "Spectral Theory of Elliptic Operators on Non-Compact Manifolds", Asterisque **207** (1992), 37-108

[Va] B. Vaillant : "Indextheorie für Überlagerungen", Diploma thesis, University of Bonn (1996)